

Is there a risk of prion-like disease transmission by Alzheimer- or Parkinson-associated protein particles?

Michael Beekes · Achim Thomzig ·
Walter J. Schulz-Schaeffer · Reinhard Burger

Received: 9 May 2014 / Revised: 18 July 2014 / Accepted: 18 July 2014 / Published online: 30 July 2014
© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract The misfolding and aggregation of endogenous proteins in the central nervous system is a neuropathological hallmark of Alzheimer's disease (AD), Parkinson's disease (PD), as well as prion diseases. A molecular mechanism referred to as “nucleation-dependent aggregation” is thought to underlie this neuropathological phenomenon. According to this concept, disease-associated protein particles act as nuclei, or seeds, that recruit cellular proteins and incorporate them, in a misfolded form, into their growing aggregate structure. Experimental studies have shown that the aggregation of the AD-associated proteins amyloid- β (A β) and tau, and of the PD-associated protein α -synuclein, can be stimulated in laboratory animal models by intracerebral (i.c.) injection of inocula containing aggregated species of the respective proteins. This has raised the question of whether AD or PD can be transmitted, like certain human prion diseases, between individuals by self-propagating protein particles potentially present on medical instruments or in blood or blood products. While the i.c. injection of inocula containing AD- or PD-associated protein aggregates was found to cause neuronal damage and clinical abnormalities (e.g., motor impairments) in some animal models, none of the studies published so far

provided evidence for a transmission of severe or even fatal disease. In addition, available epidemiological data do not indicate a transmissibility of AD or PD between humans. The findings published so far on the effects of experimentally transmitted AD- or PD-associated protein seeds do not suggest specific precautionary measures in the context of hemotherapy, but call for vigilance in transfusion medicine and other medical areas.

Keywords Alzheimer's disease · Parkinson's disease · Prion · Prion protein · Seed · Amyloid- β (A β) · Tau · α -Synuclein · Transmission · Risk

Introduction

Many neurodegenerative diseases such as Alzheimer's disease (AD), Parkinson's disease (PD), Huntington's disease, amyotrophic lateral sclerosis, or frontotemporal dementias share as a defining neuropathological feature the aggregation and deposition of misfolded endogenous proteins in the central nervous system (CNS). Although their amino acid sequences and native folds differ from each other, these disease-associated proteins form extracellular amyloid deposits or intracellular amyloid-like inclusions with common structural characteristics. Amyloid or amyloid-like protein deposits typically show a fibrillar morphology and an ordered “cross- β ” assembly of the aggregated proteins. Comprehensive evidence suggests that a general mechanism of “nucleated growth” underlies the aggregation of endogenous proteins in the neurodegenerative diseases mentioned above (for a review see [16]).

A similar mechanism is thought to govern the self-propagation and cerebral deposition of amyloid-like beta-sheet-rich aggregates of misfolded prion protein (PrP) in

M. Beekes (✉) · A. Thomzig · R. Burger
Robert Koch-Institut, Nordufer 20, 13353 Berlin, Germany
e-mail: beekesm@rki.de

A. Thomzig
e-mail: thomziga@rki.de

R. Burger
e-mail: burgerr@rki.de

W. J. Schulz-Schaeffer
University Medical Center Goettingen, Robert Koch-Str. 40,
37099 Goettingen, Germany
e-mail: wjschulz@med.uni-goettingen.de

transmissible spongiform encephalopathies (TSEs) [20, 87]. Yet, these fatal neurodegenerative diseases in animals [e.g., scrapie, bovine spongiform encephalopathy (BSE)] and humans [e.g., Creutzfeldt–Jakob disease (CJD)] were considered for many years unique. This is because they are caused, and can be transmitted between individuals, by unconventional pathogens devoid of coding nucleic acids designated as “proteinaceous infectious particles”, or “prions” [71, 72]. However, during the past few years an increasing number of prion-like phenomena have been detected in the context of AD, PD, and other non-prion neurodegenerative diseases [12, 32, 43, 52]. This has raised concerns that the transmission of pathological protein aggregates from common neurodegenerative diseases (which are generally presumed to be non-transmissible) may possibly pose a previously unrecognized risk to patient safety, e.g., in surgery or transfusion medicine.

To address this topical question for the two most frequent neurodegenerative disorders, AD and PD, this article reviews similarities and differences shown by pathological protein particles of these diseases in comparison to prions. After a brief introduction to prions and prion diseases, we provide an overview of animal experiments and epidemiological studies that have examined the transmission of protein aggregation, neurodegeneration or clinical disease between individuals by AD- or PD-associated protein particles. Finally, we discuss the results from these studies with respect to both implications for transmission risks and practical conclusions for transfusion medicine and other medical areas.

The example of prions and prion diseases

Molecular basis of prion formation, replication and transmission

Human TSEs, or prion diseases, occur in idiopathic, hereditary and acquired forms as exemplified by sporadic, familial and variant Creutzfeldt–Jakob disease (sCJD, fCJD and vCJD), respectively [73]. According to current knowledge, all of these diseases are caused and are transmissible by proteinaceous infectious particles (prions) that accumulate in the brain [71, 72, 87]. Prions are thought to be essentially composed of misfolded and aggregated conformational isomers of the cellular prion protein (PrP^C), referred to as PrP^{Sc} or PrP^{TSE} [10, 72] (the acronym PrP^{TSE} will be used in this article). While PrP^C contains a high proportion of α -helical structure and only 3 % β -sheets, PrP^{TSE} has a profoundly different, amyloid-like conformation that is substantially made up of β -sheets and exhibits a markedly reduced amount of α -helical structure [15, 29, 66].

The prevailing mechanistic model of prion replication involves nucleation-dependent aggregation of the prion protein (Fig. 1) [20, 35, 87]. According to this concept, prions of sporadic or familial CJD are thought to be initially formed by a self-assembly of PrP monomers that occurs spontaneously (sCJD) or in association with stochastic (sCJD) or inherited (fCJD) amino acid exchanges in the prion protein which facilitate a structural PrP conversion. The de novo aggregation, or nucleation, of PrP constitutes a high kinetic reaction barrier [20, 35]. Thus, nucleation is considered as a slow and rate-determining step in the genesis of sporadic and hereditary prion diseases. However, once PrP nuclei have been formed the conversion of PrP^C to PrP^{TSE} is no longer kinetically inhibited. Then, the growth of PrP aggregates proceeds swiftly by the successive association of further PrP molecules, in an aberrant folding structure that reproduces the conformation of the PrP^{TSE} template, with the nucleus. This mechanism of nucleated growth is strongly reminiscent of the acceleration of crystallization by “seeding” (i.e., the addition of preformed “seeds” to a salt or protein solution). Accordingly, “PrP seeding” provides a straightforward explanation for why prions are agents that can transmit the misfolding and aggregation of PrP between individuals: Upon introduction into a new host they simply act as exogenous proteinaceous seeds which bypass the slow nucleation step of PrP conversion. When PrP^{TSE} aggregates fragment into smaller units this increases the number of prion protein particles with seeding-active growth surfaces. Thus, while the de novo formation of prions depends on nucleation, prion replication is mediated by an alternating growth and fragmentation of PrP^{TSE} aggregates. In seeded PrP^C to PrP^{TSE} conversion, the conformation of the PrP^{TSE} template is typically reproduced with high fidelity. On this basis, phenotypically distinct prion agents that are made up of PrP with an identical amino acid and referred to as prion “types” or “strains” can exist and propagate in vivo in the form of different PrP conformers [14, 45, 94].

Based on their replication mechanism prions can transmit misfolding and aggregation of PrP (i) at the molecular level, (ii) within or at the surface of cells, (iii) from one cell to another, (iv) in and between tissues [e.g., along neuroanatomical pathways in the peripheral and central nervous system (PNS, CNS), or between the lymphoreticular system and the PNS], and (v) also between individuals. The formation and/or replication of prions trigger pathogenic processes that are still poorly understood and cause neuronal degeneration in the brain. This results in neurological or mental symptoms and eventually fatal disease in sporadic, hereditary and acquired TSEs. Prions are able to transmit TSEs between animals, from animal to man (or vice versa),




Fig. 1 Mechanistic model of prion formation and replication by nucleation-dependent aggregation of the prion protein. The prion protein (PrP) can adopt different spatial structures. Cellular prion protein (PrP^{C}) has a relatively high content of α -helical structure elements but may convert into isoforms with an aberrant conformation. Under certain conditions those PrP conformers can assemble into β -sheet-rich aggregates that constitute self-replicative proteinaceous infectious particles (prions). The misfolded and aggregated PrP in prions is referred to as PrP^{TSE} . Primary seed formation, or nucleation, of PrP underlying the initial formation of prions in sporadic or hereditary prion diseases is controlled by a high kinetic barrier. However, once

prions have been endogenously formed, or exogenously introduced into the body by infection, they behave as seeds that swiftly recruit and attach further PrP molecules. When prions eventually fragment into smaller PrP aggregates, progeny seeds enter the prion replication cycle and further propagate the pathological protein state. The NMR solution structure of recombinant human prion protein hPrP(23–230) was reproduced, in accordance with the copyright policies of PNAS, from Zahn et al. [97] (copyright by PNAS). Nucleation-dependent protein aggregation is thought to similarly underlie the formation and propagation of AD- and PD-associated protein particles

and between humans. Thus, they are genuinely infectious agents.

Notably, prion replication by the described propagation of PrP^{TSE} seeds is in principle indefinitely self-perpetuating. This holds true in cell-free conversion assays [6, 13, 93], cell assays [21, 48] and bioassays using transgenic or wild-type animals [31, 47] as evidenced by sustained PrP^{C} to PrP^{TSE} conversion and/or replication of infectivity in multiple serial passages in which the original seeds were gradually diluted out.

Iatrogenic transmission of human prions: occurrence and prevention

A total of more than 450 cases of iatrogenic CJD have occurred worldwide after application of cadaver-derived human growth hormone (c-hGH) or gonadotrophin, transplantation of dura mater or cornea, and neurosurgery or EEG recording using invasive medical devices [11]. In addition, iatrogenic transmission of vCJD by blood transfusion is presumed for three clinically diseased recipients [36, 53, 95] and for one asymptomatic recipient [68] of non-leucodepleted erythrocytes. Furthermore, asymptomatic vCJD was diagnosed in a patient with hemophilia who had received factor VIII concentrate prepared from

plasma pools that included donations from a vCJD-infected donor [67].

The BSE epidemic and resulting emergence of vCJD in the United Kingdom, and the subsequent occurrence of these novel TSEs in other countries, have substantially increased awareness of the zoonotic and iatrogenic risk potential of proteinaceous infectious particles. Together with the lessons learned from iatrogenic CJD this prompted comprehensive measures aiming at the prevention of iatrogenic prion transmissions to humans. An important element of these measures are the specifically devised safety guidelines on the reprocessing of surgical, diagnostic or other medical devices and on the supply of blood or blood products with a view to CJD and vCJD [39].

Prions and Alzheimer- or Parkinson-associated protein aggregates: commonalities and dissimilarities

Pathological protein aggregates in Alzheimer's and Parkinson's disease

The neuropathological hallmarks of AD are extracellular senile plaques and intraneuronal neurofibrillary tangles (NFT) primarily composed of aggregated amyloid- β [$\text{A}\beta$,

a peptide processed from the cellular amyloid precursor protein (APP)] and partially hyperphosphorylated tau protein, respectively [60]. In about 80 % of patients, AD is associated with cerebral amyloid angiopathy, i.e., deposits of A β in blood vessels of the brain [84]. Tau aggregates in the brain do not occur exclusively in AD, but also in other neurodegenerative diseases which are collectively referred to as tau aggregation diseases and include frontotemporal dementias [41].

Brains of patients with PD typically show cell loss in the substantia nigra (particularly in the ventral part of the pars compacta) and contain fibrillar protein inclusions in cell bodies and processes of neurons. These inclusions are designated as Lewy bodies and Lewy neurites, respectively, and essentially consist of aggregated α -synuclein, a proportion of which is phosphorylated [23]. Such Lewy pathology also occurs in “dementia with Lewy bodies” (DLB), and pathological deposits of misfolded and aggregated α -synuclein primarily in oligodendrocytes are pathognomonic of multiple system atrophy (MSA), another α -synuclein aggregation disease [16]. Recently, it has been shown that in DLB and PD most α -synuclein aggregates occur in the presynapse of nerve cells [50, 81] and are linked with a synaptic failure [83]. It is under debate, whether Lewy bodies are a compartmentalization of protein aggregates to protect the cell from harmful (synaptic) α -synuclein aggregates [65, 82].

The deposition of pathological protein aggregates in the nervous system of humans with AD and PD shows partly stereotypic temporal–spatial spreading patterns, particularly for tau and α -synuclein [12, 43, 52], which is reminiscent of the spreading characteristics of prions. Studies in animals with an experimental or natural prion infection have shown that the deposition of PrP^{TSE} in the nervous system proceeds in highly defined temporal–spatial patterns, apparently transsynaptically, along neuronal projections and circuits from initial foci to adjacent and distant target areas [4, 5, 90]. Accordingly, stereotypic spatiotemporal patterns of pathological protein deposition in AD and PD can be explained by the formation of initial protein seeds early in pathogenesis, and a subsequent prion-like spread of misfolding and aggregation along neuroanatomical pathways. Alternatively, protein aggregation may start and proceed, in a relatively uniform temporal–spatial pattern, independently at different sites in the brain of AD and PD patients, and thereby produce the observed stereotypic courses [52]. The former hypothesis, however, seems to be supported by an observed spread of α -synuclein aggregation from affected brain tissue into fetal neuronal transplants in PD patients [43], and by further findings from in vitro and in vivo studies described in the following sections.

Seeded aggregation of A β , tau and α -synuclein in vitro

The aggregation mechanisms of A β , tau and α -synuclein have been extensively explored in comprehensive in vitro studies which involved natural and recombinant protein forms [16]. Results from these studies indicate that the aggregation of A β , tau and α -synuclein occurs by a mechanism of nucleated growth which basically resembles that of PrP aggregation. However, it is still unclear to what extent AD- or PD-associated protein aggregates are self-propagating in a prion-like manner, i.e., how efficiently they fragment into progeny seeds, and how efficiently such progeny seeds then replicate in further cycles of growth and fragmentation.

During the past few years, it has become possible to mimic an indefinite self-perpetuation of the seeded aggregation of PrP (which is accompanied by a replication of infectivity) in vitro using cell-free serial protein misfolding cyclic amplification (sPMCA) [6, 13, 79, 93]. This technique reproduces putative key steps of prion replication in an accelerated manner in the test tube, and demonstrated the auto-seeding activity of newly generated PrP aggregates in multiple serial passages (i.e., after the initial seeding material had been successively diluted out). A similar, virtually indefinite, self-perpetuation of protein aggregation in cell-free conversion assays has not been shown so far for A β and tau aggregates, but was recently reported for α -synuclein based on findings in PMCA and aggregation assays that used in vitro generated fibrillar α -synuclein as seeding material [38, 77].

In cell assays, AD- and PD-associated protein aggregates show similarities to prions in terms of (i) binding to or uptake by cells of A β [96], tau [28, 63] or α -synuclein aggregates [63, 91], (ii) seeding in or on cells of A β [96], tau [63, 80] or α -synuclein aggregates [54, 63, 91], and (iii) intercellular spread of tau [28] and α -synuclein deposition [34]. In contrast with prions, aggregated A β and α -synuclein species have not yet been reported to show sustained self-propagation in multiple serial cell culture passages. However, Sanders et al. [80] recently demonstrated that different species of aggregated tau templated themselves with high fidelity through serial passages of HEK cell lines that stably expressed the aggregation-competent repeat domain of this protein.

The fact that AD- and PD-associated protein particles share nucleation-dependent protein aggregation as the molecular mechanism of their origin and other properties with prions, but not the latter’s obvious infectiousness (i.e., communicability or even contagiousness), is reflected in the term “prionoid” that has been introduced to designate prion-like seeding-active forms of A β , tau or α -synuclein [1, 2]. A substantial body of data obtained during the past few years [12, 32, 43, 52] has raised the question as to what extent A β -, tau- or α -synuclein aggregates behave like prions—and whether they can possibly transmit disease.

Experimental transmission of A β -, tau- or α -synuclein aggregates to laboratory animals

Fatal prion infections can be experimentally transmitted from affected individuals to laboratory animals. Subsequently, they can be further passaged, in principle indefinitely, in vivo. During the past few years, similarly designed animal experiments have been increasingly performed to examine whether protein aggregation, neurodegeneration or clinical disease can result from an in vivo transmission of sample materials containing A β -, tau- or α -synuclein aggregates. In the following text, such sample materials are designated as “A β -, tau or α -synuclein inocula”, or as “A β ”, “tau” or “ α -synuclein” with subscripted indices to indicate the origin of the respective inoculum.

For the experimental transmission of A β -, tau- or α -synuclein inocula, different types of wild-type animals were used, as well as transgenic rodents that homo-, hetero- or hemizygotously expressed normal or mutated human APP-, tau- or α -synuclein. The mutated protein variants were associated with familial forms of AD, frontotemporal dementia, or PD. Many transgenic models used in these transmission studies showed a considerable overexpression of the respective proteins. A proportion (but not all) of the transgenic rodent models developed during their natural lifespan a congenital phenotype with (i) protein aggregation detectable in the CNS as A β deposits, inclusions of filamentous or NFT-like tau, or accumulations of α -synuclein, (ii) neurodegeneration in the CNS, and/or (iii) cognitive or neurological symptoms. Among the rodents challenged with A β -, tau- or

α -synuclein inocula were the following animals (for study design, references and results see Table 1):

- Transgenic mice expressing mutated forms of human
 - APP (Tg2576 mice; APP23 mice; APP23:*Gfap*-luc mice; APP_{Swe}/PSEN $_{\Delta E9}$ mice; APPPS1 mice),
 - tau (B6/P301L mice; P301S mice; P301S PS19 mice), or
 - α -synuclein (homozygous A53T M83^{+/+} mice, Fig. 2)

that developed during their natural lifespan A β -, tau- or α -synuclein aggregates in the CNS, respectively. P301S PS19 and M83^{+/+} mice also developed neurodegeneration as well as severe motor dysfunction and fatal neurological disease.

- Transgenic mice hemizygotously expressing mutated human α -synuclein (A53T M83^{+/-}: *Gfap*-luc mice) which were reported to show no protein aggregation or clinical disease until at least 629 days of age. It is, however, currently unclear whether these mice may develop protein aggregates, neurodegeneration or clinical disease at higher ages.
- Transgenic mice expressing normal forms of human APP (HuAPPwt mice) or tau (ALZ17 mice), and transgenic rats expressing a mutated form of human APP (APP21 rats). These transgenic rodent models did not show protein aggregates in the CNS during their (median) natural lifespan.
- Wild-type (wt) mice.




Fig. 2 Congenital phenotype of cerebral α -synuclein aggregation in TgM83^{+/+} mice used for transmission studies with different α -synuclein inocula. TgM83^{+/+} mice expressing human A53T mutated α -synuclein begin to show α -synuclein aggregates in the CNS from an age of about 8 months. At the same age, few TgM83^{+/+} mice display signs of neurodegeneration and onset of eventually fatal motor dysfunction. The percentage of neuropathologically and clinically affected mice increases with age, and by 16 months of age all mice have developed the described phenotype [30]. The photomicrographs show pathological α -synuclein deposits in cerebral perikarya (**a**, arrows) and neurites (**b**, arrowheads) of a female TgM83^{+/+}

mouse at an age of 257 days [immunostaining was performed using an anti- α -synuclein (phospho S129) antibody (ab51253; Abcam, Cambridge, UK)]. Using this mouse model, different studies were able to demonstrate that i.c. injected inocula containing α -synuclein aggregates from mice or humans accelerated the occurrence of intracellular α -synuclein inclusions in the CNS of the recipient animals [56, 62]. Additionally, these studies showed a premature onset of fatal disease in TgM83^{+/+} mice that had been challenged by i.c. injection of those murine inocula or fibrils that had been preformed in vitro from recombinant human α -synuclein

Two important aspects have to be considered for the interpretation of results from transmission studies in transgenic animals. First, if A β -, tau- or α -synuclein inocula stimulate the formation of cerebral protein aggregates in transgenic mice which have a congenital phenotype of protein aggregation, this leaves basically two alternative explanations: (i) that the inoculated material caused (i.e., transmitted) such aggregation in the true sense, or (ii) that the inoculum merely increased or accelerated the pathological protein deposition in specific brain regions in terms of a quantitatively stronger or earlier occurrence, respectively. Accordingly, it is deemed prudent to interpret a stimulation of protein aggregation in such bioassay animals conservatively as evidence for an “increase” or “acceleration” of the protein pathology [14].

Second, transgenic animals showing no congenital protein aggregation may theoretically still contain unnaturally high levels of different species of A β -, tau- or α -synuclein assemblies. Therefore, such mice “might be borderline with respect to the levels required for triggering higher-level aggregation. Increasing the A β (or tau- or α -synuclein—author’s note) load of a particular biochemical species could then trigger A β (or tau- or α -synuclein—author’s note) deposits” [70]. Accordingly, the occurrence of protein deposition in those transgenic animals after a challenge with A β -, tau- or α -synuclein inocula should be interpreted as a “triggering” of endogenous protein deposition.

Whether A β -, tau- or α -synuclein inocula can genuinely “cause” (i.e., actually transmit) protein aggregation in vivo, finally, requires testing in wild-type animals.

Reported findings

Key findings from transmission experiments with A β -, tau- and α -synuclein inocula are described in the following paragraphs and summarized, for the studies that were performed in rodents, in Table 1.

A β

When inocula containing A β aggregates from brain tissue of AD patients, non-demented elderly donors or transgenic mice with cerebral A β -amyloidosis (A β _{hu-AD}, A β _{hu-nonAD}, or A β _{mu}, respectively), or inocula which contained A β aggregates that had been preformed in vitro from A β peptides (A β _{preformed}) were

- intracerebrally injected into Tg2576- (A β _{hu-AD} [46]), APP23-(A β _{hu-AD} [59], A β _{mu} [26, 37, 59]), APP23:*Gfap*-luc-(A β _{mu}, A β _{preformed} [88]), APP_{Swe}/PSEN $_{\Delta E9}$ - (A β _{hu-nonAD} [25]), APPPS1-(A β _{mu} [37, 59]), or B6/P301L mice (A β _{mu} [8]),
- intraperitoneally injected into APP23 mice (A β _{mu} [27]),

- or administered to APP23 mice by i.c. implantation of stainless steel wires that had been contaminated with A β _{mu} [26]

this resulted in a stimulation of the cerebral A β deposition in the recipient animals which were all congenitally predisposed for protein aggregation. Specifically, an increase of the cerebral A β -burden was found in two mouse models (APP23:*Gfap*-luc mice injected with A β _{mu} or A β _{preformed} [88], and APP_{Swe}/PSEN $_{\Delta E9}$ mice injected with A β _{hu-nonAD} [25]), while Tg2576 mice (injected with A β _{hu-AD} [46]), APP23 mice (injected with A β _{hu-AD} [59] or A β _{mu} [26, 37, 59]) and APPPS1 mice (injected with A β _{mu} [37, 59]) showed an acceleration of cerebral A β deposition.

Furthermore, the study in APP23:*Gfap*-luc mice demonstrated that preformed A β aggregates were sufficient to stimulate A β deposition in vivo [88]. In B6/P301L mice injected with A β _{mu} an acceleration of the formation of hyperphosphorylated intraneuronal tau inclusion was detected [8], which was indicative of cross-seeding. Also, intraperitoneally administered A β _{mu} was found to accelerate cerebral A β -amyloidosis [27], and increased or accelerated A β deposition in the brain was often detected beyond the site of inoculation in all studies. These latter finding resembled to some extent prion-like spreading phenomena. Finally, different morphotypes of A β aggregates were detected and differentially maintained upon inoculation into recipient animals [37]. Such morphotypes may provide yet another analogy to prions. However, the replication of distinct prion conformers can be maintained in vivo over multiple serial passages which has not been shown so far for A β .

Aliquots of A β _{hu-AD} were i.c. also injected into transgenic HuAPPwt mice [61] and APP21 rats [78] that had no congenital phenotype of protein aggregation. This triggered A β deposition, which would otherwise not have occurred, in brain tissue near the injection site, and, in HuAPPwt mice, also in distant brain areas.

However, none of the transmission studies with A β inocula provided evidence for cerebral neurodegeneration, pronounced cognitive decline or development of fatal disease in the challenged rodent models.

Primates appear as the most suited model animals for assessing potential hazards of A β inocula due to their close phylogenetic relationship to humans. Monkeys (marmosets) that had been i.c. inoculated with brain homogenates from nine patients with sporadic AD and five patients with familial AD were found in a long-term study (duration >20 years) to show more frequently and earlier cerebral A β deposition than control animals [3, 57, 76]. However, cerebral A β deposits also occurred in a proportion of control animals. In a conservative approach these findings would be interpreted as an indication that

Table 1 Effects of the transmission of sample materials containing A β -, tau- or α -synuclein aggregates to rodents

References	Study design		Effect on protein aggregation in the CNS				Spread of observed aggregation effect	Neurodegeneration in the CNS	Non-transient clinical abnormalities	Severe or fatal disease
	Inoculum	Route	Recipients	Increase	Acceleration	Triggering				
<i>Aβ</i>										
Kane et al. [46]	A β _{hu-AD}	i.c.	Tg2576 mice	+			+	- ^{la}	- ^{la}	- ^{la}
Meyer-Luehmann et al. [59]	A β _{hu-AD}	i.c.	APP23 mice	+			+	n.d.	n.d.	n.d.
	A β _{mu}	i.c.	APP23 mice	+			+	n.d.	n.d.	n.d.
	A β _{mu}	i.c.	APPS1 mice	+			+	n.d.	n.d.	n.d.
Bolmont et al. [8]	A β _{mu}	i.c.	P301L mice ^{2a}	+			+	n.d.	n.d.	n.d.
Eisele et al. [26]	A β _{mu}	i.c.	APP23 mice ³	+			+	n.d.	n.d.	n.d.
Eisele et al. [27]	A β _{mu}	i.p.	APP23 mice	+			+	n.d.	n.d.	n.d.
Rosen et al. [78]	A β _{hu-AD}	i.c.	APP21 rats		+		-	n.d.	n.d.	n.d.
Morales et al. [61]	A β _{hu-AD}	i.c.	HuAPPwt mice		+		+	n.d.	n.d.	n.d.
Stöhr et al. [88]	A β _{mu}	i.c.	APP23;G β -luc mice	+			+	n.d.	n.d.	n.d.
	A β _{preformed}	i.c.	APP23;G β -luc mice	+			+	n.d.	n.d.	n.d.
Duran-Aniotz et al. [25]	A β _{hu-non AD}	i.c.	APP ^{swe} /PSEN Δ E9 mice	+			+	n.d.	n.d.	n.d.
Heilbronner et al. [37]	A β _{mu}	i.c.	APP23 mice	+			+	n.d.	n.d.	n.d.
	A β _{mu}	i.c.	APPS1 mice	+			+	n.d.	n.d.	n.d.
<i>Tau</i>										
Clavaguera et al. [17]	tau _{mu}	i.c.	ALZ17 mice			+	+	- ^{lb}	n.d.	n.d.
Clavaguera et al. [18]	tau _{hu}	i.c.	ALZ17 mice			+	+	- ^{lc}	n.d.	n.d.
	tau _{hu}	i.c.	C57BL/6 wt mice				+	n.d.	n.d.	n.d.
	tau _{mu}	i.c.	ALZ17 mice			+	+	n.d.	n.d.	n.d.
Lasagna-Reeves et al. [51]	tau _{hu}	i.c.	C57BL/6 wt mice				+	n.d.	- ^{ld}	- ^{ld}
Iba et al. [40]	tau _{preformed}	i.c.	P301S PS19 mice	+			+	- ^{le}	n.d.	- ^{lf}
Clavaguera et al. [19]	tau _{mu}	i.p.	P301S mice	+			+	n.d.	n.d.	n.d.
Sanders et al. [80]	tau _{mu}	i.c.	P301S PS19 mice		+		n.d.	n.d.	n.d.	n.d.
	tau _{preformed}	i.c.	P301S PS19 mice	+			n.d.	n.d.	n.d.	n.d.
	tau _{HEK cells}	i.c.	P301S PS19 mice	+			+ ⁴	n.d.	n.d.	n.d.
<i>α-Synuclein</i>										
Mougenot et al. [62]	α -syn _{mu}	i.c.	TgM83 ^{+/+} mice	+			+	n.d.		+(Acceleration)
Luk et al. [55]	α -syn _{preformed/mu}	i.c.	C57BL/6 wt mice				+	+(Causation)	+(Causation)	- ^{lj}
Luk et al. [56]	α -syn _{mu}	i.c.	TgM83 ^{+/+} mice	+			+	+(Acceleration)	+(Acceleration)	+(Acceleration)
	α -syn _{preformed/mu}	i.c.	TgM83 ^{+/+} mice	+			+	+(Acceleration)	+(Acceleration)	+(Acceleration)

Table 1 continued

References	Study design		Effect on protein aggregation in the CNS				Spread of observed aggregation effect	Neurodegeneration in the CNS	Non-transient clinical abnormalities	Severe or fatal disease
	Inoculum	Route	Recipients	Increase	Acceleration	Triggering				
Masuda-Suzukake et al. [58]	α -syn _{hu-DLB}	i.c.	C57BL/6 wt mice			+	+	n.d.	n.d.	n.d.
	α -syn _{preformed/mu}	i.c.	C57BL/6 wt mice			+	+	-lg	-lh	-lh
	α -syn _{preformed/hu}	i.c.	C57BL/6 wt mice			+	+	-lg	-lh	-lh
Watts et al. [92]	α -syn _{hu-MSA}	i.c.	TgM83 ^{+/-} mice	+			+	n.d.		+(Acceleration)
Guo et al. [33]	α -syn _{preformed/hu}	i.c.	P301S P19 mice ^{2b}	+		+	+	n.d.	n.d.	n.d.
Recasens et al. [75]	α -syn _{hu-PD}	i.c.	C57BL/6 wt mice			+	+	+(Causation)	+(Causation)	-li

A β , *tau*- or *α -syn* inocula containing A β -, tau- or α -synuclein aggregates, respectively (subscripted indices indicate the following origins: HEK cells HEK cell lines that stably expressed the aggregation-competent repeat domain of tau, *hu-AD* humans with AD, *hu-DLB* humans with DLB, *hu-non-AD* humans without AD, *hu-MSA* humans with MSA, *hu-PD* humans with PD, *mu* mice, *preformed* in vitro preformed peptide or protein aggregates, *preformed/hu* or *preformed/mu* in vitro preformed aggregates of human or murine proteins, respectively), *n.d.* not determined, + positive finding reported, - negative finding reported, *i.c.* intracerebral administration to recipients, *i.p.* intraperitoneal administration to recipients

^{1a-i} Negative up to 5^a, 15^{b,c,g}, 11^d, 6^{e,h,j}, 9^f or 17ⁱ months post-inoculation (g—negative finding refers to absence of dopaminergic degeneration)

^{2a,b} Indication of cross-seeding by A β _{mu} and α -syn_{preformed/hu} of mutated human tau, respectively

³ Intracerebral inoculation with A β _{mu} was performed by injection of liquid sample materials or implantation of steel wires

⁴ Positive finding with one of the tested HEK cell clones

the injected inocula accelerated A β amyloidosis. Whether a triggering or causation of cerebral A β deposition had occurred in a proportion of the challenged primates could not be resolved by the experiment. In any case, none of the challenged monkeys showed the development of neurofibrillary tau deposits or AD typical signs of disease. The findings from this long-term study in nonhuman primates are consistent with the notion that cerebral A β deposits are an essential but not sufficient neuropathological criterion for AD, and with reports that the brain of elderly people who are not affected by neurodegenerative dementias may contain substantial deposits of aggregated A β [74, 85, 86, 89].

Similarly, the inoculation of sample material from more than 100 cases of AD into primates at the US National Institutes of Health (NIH) did not result in motor or behavioral abnormalities in the recipients that would have indicated a transmission of disease [9]. Additionally, neuropathological postmortem examinations of brain specimens from primates of the NIH study did not provide evidence for a pre- or subclinical disease transmission, according to a personal communication by members of the former NIH research team (P. Brown, D. Asher) reported by Irwin et al. [42]. In contrast to these negative findings with AD inocula, more than 300 cases of prion diseases were transmitted to primates at the NIH during the same period of time [9].

Tau

The administration of inocula which had been prepared from brain tissue of transgenic mice with cerebral tau deposits (tau_{mu}) or lysates of transgenic HEK cell lines that propagated tau inclusions (tau_{HEK cells}), as well as the administration of inocula which contained tau aggregates that had been preformed in vitro from recombinant tau protein (tau_{preformed}), by

- i.c. injection into P301S PS19 mice (tau_{mu} and tau_{HEK cells}) [80]; tau_{preformed} [40, 80] or
- intraperitoneal injection into P301S mice (tau_{mu} [19])

accelerated or increased, respectively, the cerebral deposition of filamentous tau. The increase of tau aggregation in the brain of P301S mice which had been intraperitoneally injected with tau_{mu} indicated a centrifugal spread of tau aggregates or aggregation.

In addition, tau_{mu} [17, 18] or inocula containing aggregated tau from brain tissue of human patients with tau aggregation diseases (tau_{hu} [18]) were intracerebrally injected into transgenic ALZ17 mice. This triggered the initiation and spread of filamentous tau deposition in the brain of these recipient animals. The deposition of filamentous

tau eventually extended from the injection sites to distant, anatomically connected brain regions.

I.c. injection of tau_{hu} into C57BL/6 wild-type mice caused inclusions of murine tau that were detected 6 months after inoculation at the injection site and subsequently increased over time [18]. When purified tau oligomers from AD brain tissue were intracerebrally injected into wild-type C57BL/6 mice, this induced widespread deposits of hyperphosphorylated murine tau filaments in the injected area and neighboring brain regions [51]. According to the authors of the latter study, AD-associated tau oligomers were thus shown to cause tau deposition in wild-type mice “via a mechanism reminiscent of that used by prions”.

Finally, serial transmission of tau seeding was observed in ALZ17 mice following the i.c. injection of homogenates prepared from ALZ17- or C57BL/6 brains that had been originally injected with tau_{mu} or tau_{hu}, respectively [18]. Taken together, these findings suggested according to Clavaguera et al. [18] that “once tau aggregates have formed in discrete brain areas, they become self-propagating and spread in a prion-like manner”.

Most recently, Sanders et al. [80] reported that distinct species of tau aggregates could be stably propagated through multiple generations of transgenic P301S PS19 mice. These authors concluded that tau “encodes self-catalyzing conformational information that it propagates indefinitely with high fidelity”.

Despite the ability of tau inocula to increase, accelerate, trigger and cause tau aggregation in vivo, none of the transmission studies provided evidence for cerebral neurodegeneration, clinical abnormalities or severe disease in tau-challenged mice.

α -Synuclein

When inocula containing aggregated α -synuclein from brain homogenates of aged TgM83^{+/+} mice that had developed cerebral protein aggregates and neurodegeneration as well as fatal clinical disease (α -synuclein_{mu}) were i.c. injected into young asymptomatic TgM83^{+/+} mice, this accelerated the occurrence of intracellular α -synuclein inclusions in the brains of the recipient animals [56, 62]. The deposition of α -synuclein seemed to propagate through the CNS in a time-dependent manner along neural pathways and also occurred in the substantia nigra [56]. This suggested a parallel to both the “apparent spread of α -synuclein deposits in the human brain” [44], and the propagation of prions along neuroanatomical projections. After i.c. injection of α -synuclein_{mu}, recipient animals eventually showed a reduction of tyrosine hydroxylase staining in nigral neurons with α -synuclein inclusions (suggesting impaired dopamine production in

these cells) [56]. Additionally, TgM83^{+/+} mice that had been inoculated with α -synuclein_{mu} showed an accelerated development of severe motor dysfunction and premature death [56, 62]. After i.c. injection of inocula containing fibrils that had been preformed in vitro from recombinant human α -synuclein (α -synuclein_{preformed/hu}), similar results were found in TgM83^{+/+} mice with respect to accelerated α -synuclein aggregation and premature onset of clinical disease [56].

Recently, Watts et al. [92] reported the experimental transmission of protein aggregation and fatal clinical disease for multiple system atrophy (MSA), an α -synuclein aggregation disease with α -synuclein inclusions in oligodendrocytes. In this study, inocula containing human α -synuclein aggregates from brain homogenates of MSA patients (α -synuclein_{hu-MSA}) were intracerebrally injected into M83^{+/-}: *Gfap*-luc mice [92]. Control M83^{+/-}: *Gfap*-luc mice that had not been i.c. injected with α -synuclein_{hu-MSA} showed no protein aggregation or clinical disease (neurodegeneration was not examined) until at least 629 days of age. However, it is currently unclear whether such control mice may have developed a congenital phenotype of protein aggregation in the CNS and clinical disease at higher ages. Hemizygous TgM83^{+/-} mice that do not carry the *Gfap*-luc transgene develop pathological α -synuclein deposition in the CNS at 22–28 months of age. At the same age (660–840 days), many TgM83^{+/-} mice also develop the disease phenotype of homozygous TgM83^{+/+} mice [30]. Thus, a prudent approach would take the data reported by Watts et al. [92] either as an acceleration, or—if the absence of protein aggregation and clinical disease were shown to persist in M83^{+/-}: *Gfap*-luc control mice during their natural lifespan—as a triggering of protein aggregation and clinical disease by the injected α -synuclein_{hu-MSA}.

In another study, inocula containing two types of preformed fibrils produced in vitro from recombinant, C-terminally truncated human α -synuclein (α -synuclein_{preformed/hu}) were intracerebrally injected into transgenic P301S PS19 mice [33]. While one type of the fibrils substantially accelerated cerebral tau deposition, possibly by cross-seeding, the other type of fibrils was much less able to do so. The two types of fibrils apparently represented distinct conformers of aggregated α -synuclein, and their different effects in terms of the stimulation of tau aggregation may be indicative of analogies to phenomena observed in the context of prion strains or prion types.

Furthermore, inocula containing fibrils that had been preformed in vitro from recombinant murine or human α -synuclein (α -synuclein_{preformed/mu} and α -synuclein_{preformed/hu}, respectively), and inocula containing α -synuclein aggregates from enriched extracts of brain tissue from a patient with DLB (α -synuclein_{hu-DLB}) or from patients

with PD (α -synuclein_{hu-PD}) caused cerebral deposition of endogenous α -synuclein in different types of C57BL/6 wild-type mice after injection into the striatum (α -synuclein_{preformed/mu} [55]) or administration to the substantia nigra (α -synuclein_{preformed/mu}, α -synuclein_{preformed/hu}, α -synuclein_{hu-DLB} [58] and α -synuclein_{hu-PD} [75]). The subsequent spread of α -synuclein deposition seemed to follow neuronal connections and circuits. While the three studies in wild-type mice by Luk et al. [55], Masuda-Suzukake et al. [58] and Recasens et al. [75] produced similar results in these respects, they partly differed with regard to the occurrence of neurodegeneration and clinical abnormalities in the challenged animals.

Luk et al. [55] observed in their study a gradual loss of tyrosine hydroxylase immunoreactivity and neurons in the substantia nigra pars compacta, and reduced striatal dopamine levels in the injected hemisphere. This was accompanied by a performance deterioration in rotarod tests (latency to fall: ~200 s for α -synuclein_{preformed/mu}-injected animals vs. ~325 s for control animals), and a poorer performance in wire-hang tests. However, gross motor or behavioral abnormalities were not observed, and overall motor activity was not significantly altered. In the study by Recasens et al. [75] the i.c. injection of α -synuclein_{hu-PD} caused a progressive nigrostriatal dopaminergic neurodegeneration. At 4 months post-inoculation these authors observed impaired motor ability when mice were examined in the pole test which measures motor coordination and balance. Yet, again, no gross motor or behavioral abnormalities were observed up to 17 months post-inoculation. Finally, Masuda-Suzukake et al. [58] reported findings suggesting that dopaminergic neurons were retained in the substantia nigra after injection of α -synuclein_{preformed/hu}, while they observed a prominent reduction of the neurotransmitter enkephalin in some brain regions. In the study by Masuda-Suzukake et al. [58], no significant differences were observed in rotarod- or wire-hang tests between α -synuclein_{preformed/hu}-injected and control mice at 6 months post-inoculation. At this time-point prominent cerebral α -synuclein deposition was already detectable in α -synuclein_{preformed/hu}-injected mice.

Despite these partly divergent results, the studies by Luk et al. [55], Masuda-Suzukake et al. [58] and Recasens et al. [75] consistently failed to detect typical PD symptoms (such as tremor, rigor, slowness of movement or postural instability) in wild-type mice after the transmission of α -synuclein inocula.

Recasens et al. [58] also injected α -synuclein_{hu-PD} into the striatum or substantia nigra of rhesus monkeys. Striatal inoculation produced a widespread increase of α -synuclein immunolabeling and also an increase of phosphorylated α -synuclein species in the brain, while injection into the substantia nigra essentially resulted in diffuse

local intracellular accumulations of α -synuclein. These α -synuclein pathologies were accompanied by a slowly progressing dopaminergic nigrostriatal neurodegeneration. At 14 months post-inoculation, when the study was terminated, the α -synuclein_{hu-PD}-injected monkeys showed neither behavioral changes nor signs of clinical disease.

When samples from 24 cases of PD with dementia were inoculated into primates at the NIH, this produced exclusively negative results. The challenged primates did not show clinical signs of disease transmission [9]. Neuropathological postmortem examinations also failed to reveal indications for a pre- or subclinical disease transmission according to a personal communication by members of the former NIH research team (P. Brown, D. Asher) reported by Irwin et al. [42].

Epidemiological studies on the transmissibility of Alzheimer's and Parkinson's disease

Available epidemiological data do not provide indications that transmission of AD or PD between humans occurs in real life. Observational and cohort studies failed to identify sufficient evidence for firm conclusions on the association of any modifiable risk factors for the development of AD [24, 69]. A limited number of studies specifically examined the epidemiological association between AD and blood transfusions. Findings of a study by Kokmen et al. [49] suggested that blood transfusions do not constitute a risk factor for AD in transfusion recipients. Two case-control studies by O'Meara et al. [64] and Bohnen et al. [7] which examined more than 550 cases and corresponding controls also did not find an association between blood transfusions and AD. This result was not influenced by the latency period between transfusion and occurrence of AD in transfusions recipients, the recipient's apolipoprotein E epsilon 4 (APOE-e4) status, or the number of received transfusions. Finally, an epidemiological analysis in the United Kingdom of the causes of death of more than 6,000 hemophiliacs who had received blood products via plasma protein therapy throughout their whole life did not detect an elevated level of diseases of the nervous system in this group of patients [22].

A recent study by Irwin et al. [42] examined the risk of AD or PD transmission in recipients of cadaveric human growth hormone that had been prepared from the pituitary gland of deceased donors. Medical treatments with c-hGH caused more than 200 cases of iatrogenic Creutzfeldt–Jakob disease (iCJD) worldwide due to inadvertent contamination with infectious PrP^{TSE} seeds (i. e. prions) from donors who pre- or subclinically had incubated a prion disease [11]. Against this background, Irwin et al. analyzed whether AD or PD occurred, similarly

to iCJD, more frequently in recipients of c-hGH than in control persons. The authors reported that they found mild amounts of aggregated A β -, tau- and α -synuclein in the pituitary glands of normal individuals as well as of patients with neurodegenerative diseases. At the same time, the incidence of PD and AD exceeds that of human prion diseases in the general population (and thus probably also in c-hGH donors) by more than two or three orders of magnitude, respectively. Accordingly, recipients of c-hGH should have been exposed to potentially seeding-active A β -, tau- or α -synuclein particles with a much higher likelihood than to prions, provided that AD- or PD-associated protein particles passed like PrP^{TSE} seeds through the extraction procedure of c-hGH. Yet, Irwin et al. found in their epidemiological analysis that of 796 deceased c-hGH recipients none had developed AD or PD (mean age at death: 27.2 years [range 0–77 years]; mean duration from first treatment to death: 16.3 years [range 0–45 years]). Basically, this finding suggested two alternative conclusions: (i) That AD- or PD-associated protein particles were present in c-hGH but not able to transmit AD or PD to the hormone recipients, or (ii) that A β -, tau- or α -synuclein aggregates were more effectively removed or inactivated by the procedure for c-hGH purification than PrP^{TSE} seeds. In any case, no indications for a prion-like disease transmission by A β -, tau- or α -synuclein aggregates were detected in recipients of c-hGH, a group of patients that provide a “unique in vivo model of human-to-human transmission” [42].

Discussion

Transmission studies in animals

A substantial body of experimental data shows that the administration of inocula containing A β -, tau- or α -synuclein aggregates of animal, human or in vitro origin can increase, accelerate, trigger or cause the cerebral aggregation and deposition of endogenous A β -, tau- and α -synuclein in laboratory animals. As to the negative findings that have been published [9] or otherwise communicated [42] from the NIH’s primate study it should be noted that these experiments were performed before the modern tools to analyze neuropathological lesions of AD and PD were available. Thus, it appears tempting to speculate whether a re-examination of brain tissue specimens from these studies using current tools and antibodies (if possible at all) would, or would not, reveal the transmission of AD- or PD-like protein aggregation.

In contrast to the efficient stimulation of protein aggregation in vivo by A β -, tau- and α -synuclein inocula these sample materials have not been found to transmit severe

or fatal diseases after i.c. injection into rodents and primates. Thus, in terms of disease transmission A β -, tau- and α -synuclein particles seem to fundamentally differ from prions. However, the failure to detect any transmission of a severe cognitive or neurological disease after i.c. injection of A β -, tau- and α -synuclein inocula may be due to the inability of the used animal models to reproduce cardinal symptoms of complex human neurodegenerative diseases such as AD or PD. Additionally, many transgenic mouse models were primarily engineered for the purpose of developing morphologically identifiable A β -, tau- or α -synuclein aggregates. Those mouse models often do not show significant CNS neurodegeneration (in proportion to the burden of protein aggregates), and whether they represent models of disease as opposed to protein aggregation is unclear. Although such transgenic models can be very sensitive to protein seeding by injected A β -, tau- or α -synuclein samples they may be rather insensitive to clinical disease transmission. In contrast to such transgenic mouse models of neuropathological AD or PD features, PrP transgenic mice were often specifically designed to increase the susceptibility to prion infections. This potential caveat has to be considered for AD or PD transmission studies in transgenic animals but does not apply to wild-type rodents or primates. Still, transmission studies in these latter animals also revealed a conspicuous discrepancy between efficient experimental acceleration (by A β inocula) or transmission (by tau- and α -synuclein inocula) of protein aggregation, and a lack of concomitant acceleration or transmission, respectively, of severe cognitive or neurological disease. Thus, the consistently failed disease transmissions by A β -, tau- and α -synuclein inocula observed so far may suggest an alternative conclusion: That A β -, tau- and α -synuclein aggregates, also when being seeding-active and partly able to produce neurotoxic effects, cannot transmit neurodegenerative diseases such as AD or PD.

In any case, the seeding effects that A β -, tau- or α -synuclein inocula have been shown to exert on endogenous proteins of laboratory animals have raised concerns of potential infectiological risks. A transfer of AD or PD protein particles between humans could hypothetically occur via contaminated and insufficiently reprocessed surgical instruments, or via blood or blood products.

Epidemiological studies

Epidemiological studies can help to assess whether there are tangible risks for the transmission of AD or PD under real-life conditions. Available data from such studies do not provide indications for the transmissibility or transmission of AD or PD between humans via blood, blood products

or other routes. However, AD and PD have a much higher prevalence than human prion diseases, and the interpretation of epidemiological studies on highly prevalent and multifactorial diseases such as AD and PD is complicated. Therefore, it may be difficult to epidemiologically track down any infectious origin if such cause underlay a subgroup of AD or PD cases.

Conclusion and outlook

So far, neither experimental nor epidemiological studies provided evidence for a transmission of severe or even fatal disease by AD- or PD-associated protein particles. However, in the light of findings in TgM83 mice [56, 62, 92] it appears conceivable that exogenous α -synuclein seeds may accelerate the pathogenesis of genetically predisposed α -synuclein aggregation diseases. Furthermore, inocula that contained α -synuclein aggregates caused neurotoxic effects and neurological impairments in some studies with wild-type mice [55, 75]. Thus, a thorough decontamination of surgical instruments and other medical devices from aggregated A β -, tau- and α -synuclein by effective and routinely applicable reprocessing procedures may possibly add to patient safety. In contrast, the current data reviewed in this article do not suggest specific precautionary measures for hemotherapy.

Certainly, further epidemiological and experimental research is necessary to clarify the hazards possibly emanating from transmissible A β -, tau- or α -synuclein seeds in more depth. Yet, for the time being, prions of transmissible spongiform encephalopathies seem to remain unique pathogens.

Acknowledgments The German Federal Ministry of Health financially supports a study at the Robert Koch-Institute that addresses the safety of blood and medical devices in view of prion-like phenomena possibly associated with Alzheimer's disease, Parkinson's disease and other non-prion proteopathies (IIA5-2512NIK004//321-4471-02).

Conflict of interest The authors declare that no competing interests exist.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Aguzzi A (2009) Cell biology: beyond the prion principle. *Nature* 459:924–925
- Ashe KH, Aguzzi A (2013) Prions, prionoids and pathogenic proteins in Alzheimer disease. *Prion* 7:55–59
- Baker HF, Ridley RM, Duchon LW, Crow TJ, Bruton CJ (1994) Induction of beta (A4)-amyloid in primates by injection of Alzheimer's disease brain homogenate. Comparison with transmission of spongiform encephalopathy. *Mol Neurobiol* 8:25–39
- Beekes M, McBride PA, Baldauf E (1998) Cerebral targeting indicates vagal spread of infection in hamsters fed with scrapie. *J Gen Virol* 79:601–607
- Beekes M, McBride PA (2007) The spread of prions through the body in naturally acquired transmissible spongiform encephalopathies. *FEBS J* 264:588–605
- Bieschke J, Weber P, Sarafoff N, Beekes M, Giese A, Kretzschmar H (2004) Autocatalytic self-propagation of misfolded prion protein. *Proc Natl Acad Sci* 101:12207–12211
- Bohnen NI, Warner MA, Kokmen E, Beard CM, Kurland LT (1994) Prior blood transfusions and Alzheimer's disease. *Neurology* 44:1159–1160
- Bolmont T, Clavaguera F, Meyer-Luehmann M et al (2007) Induction of tau pathology by intracerebral infusion of amyloid-beta-containing brain extract and by amyloid-beta deposition in APP x Tau transgenic mice. *Am J Pathol* 171:2012–2020
- Brown P, Gibbs CJ Jr, Rodgers-Johnson P et al (1994) Human spongiform encephalopathy: the National Institutes of Health series of 300 cases of experimentally transmitted disease. *Ann Neurol* 35:513–529
- Brown P, Cervenakova L (2005) A prion lexicon (out of control). *Lancet* 365:122
- Brown P, Brandel JP, Sato T et al (2012) Iatrogenic Creutzfeldt-Jakob disease, final assessment. *Emerg Infect Dis* 18:901–907
- Brundin P, Melki R, Kopito R (2010) Prion-like transmission of protein aggregates in neurodegenerative diseases. *Nat Rev Mol Cell Biol* 11:301–307
- Castilla J, Saa P, Hetz C, Soto C (2005) In vitro generation of infectious scrapie prions. *Cell* 121:195–206
- Caughey B, Baron GS, Chesebro B, Jeffrey M (2009) Getting a grip on prions: oligomers, amyloids, and pathological membrane interactions. *Annu Rev Biochem* 78:177–204
- Caughey BW, Dong A, Bhat KS, Ernst D, Hayes SF, Caughey WS (1991) Secondary structure analysis of the scrapie-associated protein PrP 27–30 in water by infrared spectroscopy. *Biochemistry* 30:7672–7680
- Chiti F, Dobson CM (2006) Protein misfolding, functional amyloid, and human disease. *Annu Rev Biochem* 75:333–366
- Clavaguera F, Bolmont T, Crowther RA et al (2009) Transmission and spreading of tauopathy in transgenic mouse brain. *Nat Cell Biol* 11:909–913
- Clavaguera F, Akatsu H, Fraser G et al (2013) Brain homogenates from human tauopathies induce tau inclusions in mouse brain. *Proc Natl Acad Sci* 110:9535–9540
- Clavaguera F, Hench J, Lavenir I et al (2014) Peripheral administration of tau aggregates triggers intracerebral tauopathy in transgenic mice. *Acta Neuropathol* 127:299–301
- Come JH, Fraser PE, Lansbury PT (1993) A kinetic model for amyloid formation in the prion diseases: importance of seeding. *Proc Natl Acad Sci* 90:5959–5963
- Courageot MP, Daude N, Nonno R et al (2008) A cell line infectible by prion strains from different species. *J Gen Virol* 89:341–347
- Darby SC, Kan SW, Spooner RJ et al (2007) Mortality rates, life expectancy, and causes of death in people with hemophilia A or B in the United Kingdom who were not infected with HIV. *Blood* 110:815–825
- Davie CA (2008) A review of Parkinson's disease. *Br Med Bull* 86:109–127
- Davignus ML, Plassman BL, Pirzada A et al (2011) Risk factors and preventive interventions for Alzheimer disease: state of the science. *Arch Neurol* 68:1185–1190
- Duran-Aniotz C, Morales R, Moreno-Gonzalez I, Hu PP, Soto C (2013) Brains from non-Alzheimer's individuals containing

- amyloid deposits accelerate Abeta deposition in vivo. *Acta Neuropathol Commun* 1:76
26. Eisele YS, Bolmont T, Heikenwalder M et al (2009) Induction of cerebral beta-amyloidosis: intracerebral versus systemic Abeta inoculation. *Proc Natl Acad Sci* 106:12926–12931
 27. Eisele YS, Obermuller U, Heilbronner G et al (2010) Peripherally applied Abeta-containing inoculates induce cerebral beta-amyloidosis. *Science* 330:980–982
 28. Frost B, Ollesch J, Wille H, Diamond MI (2009) Conformational diversity of wild-type Tau fibrils specified by templated conformation change. *J Biol Chem* 284:3546–3551
 29. Gasset M, Baldwin MA, Lloyd DH et al (1992) Predicted alpha-helical regions of the prion protein when synthesized as peptides form amyloid. *Proc Natl Acad Sci* 89:10940–10944
 30. Giasson BI, Duda JE, Quinn SM, Zhang B, Trojanowski JQ, Lee VM (2002) Neuronal alpha-synucleinopathy with severe movement disorder in mice expressing A53T human alpha-synuclein. *Neuron* 34:521–533
 31. Groschup MH, Gretzschel A, Kuczus T (2006) Prion strains. In: Hörlmann B, Riesner D, Kretzschmar H (eds) *Prions in animals and humans*. de Gruyter, Berlin, pp 166–183
 32. Guest WC, Silverman JM, Pokrishevsky E, O'Neill MA, Grad LI, Cashman NR (2011) Generalization of the prion hypothesis to other neurodegenerative diseases: an imperfect fit. *J Toxicol Environ Health A* 74:1433–1459
 33. Guo JL, Covell DJ, Daniels JP et al (2013) Distinct alpha-synuclein strains differentially promote tau inclusions in neurons. *Cell* 154:103–117
 34. Hansen C, Angot E, Bergstrom AL et al (2011) Alpha-synuclein propagates from mouse brain to grafted dopaminergic neurons and seeds aggregation in cultured human cells. *J Clin Invest* 121:715–725
 35. Harper JD, Lansbury PT Jr (1997) Models of amyloid seeding in Alzheimer's disease and scrapie: mechanistic truths and physiological consequences of the time-dependent solubility of amyloid proteins. *Annu Rev Biochem* 66:385–407
 36. Health Protection Agency (2007) Health protection report (19 January 2007): Fourth case of transfusion-associated variant-CJD infection
 37. Heilbronner G, Eisele YS, Langer F et al (2013) Seeded strain-like transmission of beta-amyloid morphotypes in APP transgenic mice. *EMBO Rep* 14:1017–1022
 38. Herva ME, Zibae S, Fraser G, Barker RA, Goedert M, Spillantini MG (2014) Anti-amyloid compounds inhibit alpha-synuclein aggregation induced by protein misfolding cyclic amplification (PMCA). *J Biol Chem* 289:11897–11905
 39. Hörlmann B, Pauli G, Lemmer K, Beekes M, Mielke M (2006) Prevention of the transmission of prion diseases in healthcare settings. In: Hörlmann B, Riesner D, Kretzschmar HA (eds) *Prions in humans and animals*. de Gruyter, Berlin, pp 546–560
 40. Iba M, Guo JL, McBride JD, Zhang B, Trojanowski JQ, Lee VM (2013) Synthetic tau fibrils mediate transmission of neurofibrillary tangles in a transgenic mouse model of Alzheimer's-like tauopathy. *J Neurosci* 33:1024–1037
 41. Iqbal K, Liu F, Gong CX, Grundke-Iqbal I (2010) Tau in Alzheimer disease and related tauopathies. *Curr Alzheimer Res* 7:656–664
 42. Irwin DJ, Abrams JY, Schonberger LB et al (2013) Evaluation of potential infectivity of Alzheimer and Parkinson disease proteins in recipients of cadaver-derived human growth hormone. *JAMA Neurol* 70:462–468
 43. Jucker M, Walker LC (2011) Pathogenic protein seeding in Alzheimer disease and other neurodegenerative disorders. *Ann Neurol* 70:532–540
 44. Jucker M, Walker LC (2013) Self-propagation of pathogenic protein aggregates in neurodegenerative diseases. *Nature* 501:45–51
 45. Kabir ME, Safar JG (2014) Implications of prion adaptation and evolution paradigm for human neurodegenerative diseases. *Prion* 8:111–116
 46. Kane MD, Lipinski WJ, Callahan MJ et al (2000) Evidence for seeding of beta-amyloid by intracerebral infusion of Alzheimer brain extracts in beta-amyloid precursor protein-transgenic mice. *J Neurosci* 20:3606–3611
 47. Kimberlin RH, Walker CA (1978) Evidence that the transmission of one source of scrapie agent to hamsters involves separation of agent strains from a mixture. *J Gen Virol* 39:487–496
 48. Klöhn PC, Stoltze L, Flechsig E, Enari M, Weissmann C (2003) A quantitative, highly sensitive cell-based infectivity assay for mouse scrapie prions. *Proc Natl Acad Sci* 100:11666–11671
 49. Kokmen E, Beard CM, O'Brien PC, Kurland LT (1996) Epidemiology of dementia in Rochester, Minnesota. *Mayo Clin Proc* 71:275–282
 50. Kramer ML, Schulz-Schaeffer WJ (2007) Presynaptic alpha-synuclein aggregates, not Lewy bodies, cause neurodegeneration in dementia with Lewy bodies. *J Neurosci* 27:1405–1410
 51. Lasagna-Reeves CA, Castillo-Carranza DL, Sengupta U et al (2012) Alzheimer brain-derived tau oligomers propagate pathology from endogenous tau. *Sci Rep* 2:700
 52. Lee SJ, Lim HS, Masliah E, Lee HJ (2011) Protein aggregate spreading in neurodegenerative diseases: problems and perspectives. *Neurosci Res* 70:339–348
 53. Llewellyn CA, Hewitt PE, Knight RS et al (2004) Possible transmission of variant Creutzfeldt–Jakob disease by blood transfusion. *Lancet* 363:417–421
 54. Luk KC, Song C, O'Brien P et al (2009) Exogenous alpha-synuclein fibrils seed the formation of Lewy body-like intracellular inclusions in cultured cells. *Proc Natl Acad Sci* 106:20051–20056
 55. Luk KC, Kehm V, Carroll J et al (2012) Pathological alpha-synuclein transmission initiates Parkinson-like neurodegeneration in nontransgenic mice. *Science* 338:949–953
 56. Luk KC, Kehm VM, Zhang B, O'Brien P, Trojanowski JQ, Lee VM (2012) Intracerebral inoculation of pathological alpha-synuclein initiates a rapidly progressive neurodegenerative alpha-synucleinopathy in mice. *J Exp Med* 209:975–986
 57. Maclean CJ, Baker HF, Ridley RM, Mori H (2000) Naturally occurring and experimentally induced beta-amyloid deposits in the brains of marmosets (*Callithrix jacchus*). *J Neural Transm* 107:799–814
 58. Masuda-Suzukake M, Nonaka T, Hosokawa M et al (2013) Prion-like spreading of pathological alpha-synuclein in brain. *Brain* 136:1128–1138
 59. Meyer-Luehmann M, Coomaraswamy J, Bolmont T et al (2006) Exogenous induction of cerebral beta-amyloidogenesis is governed by agent and host. *Science* 313:1781–1784
 60. Minati L, Edginton T, Bruzzone MG, Giaccone G (2009) Current concepts in Alzheimer's disease: a multidisciplinary review. *Am J Alzheimers Dis Other Demen* 24:95–121
 61. Morales R, Duran-Aniotz C, Castilla J, Estrada LD, Soto C (2012) De novo induction of amyloid- β deposition in vivo. *Mol Psychiatry* 17:1347–1353
 62. Mougnot AL, Nicot S, Bencsik A et al (2011) Prion-like acceleration of a synucleinopathy in a transgenic mouse model. *Neurobiol Aging* 33:2225–2228
 63. Nonaka T, Watanabe ST, Iwatsubo T, Hasegawa M (2010) Seeded aggregation and toxicity of alpha-synuclein and tau: cellular models of neurodegenerative diseases. *J Biol Chem* 285:34885–34898
 64. O'Meara ES, Kukull WA, Schellenberg GD et al (1997) Alzheimer's disease and history of blood transfusion by apolipoprotein-E genotype. *Neuroepidemiology* 16:86–93
 65. Olanow CW, Perl DP, DeMartino GN, McNaught KS (2004) Lewy-body formation is an aggresome-related process: a hypothesis. *Lancet Neurol* 3:496–503

66. Pan KM, Baldwin M, Nguyen J et al (1993) Conversion of alpha-helices into beta-sheets features in the formation of the scrapie prion proteins. *Proc Natl Acad Sci* 90:10962–10966
67. Peden A, McCordle L, Head MW et al (2010) Variant CJD infection in the spleen of a neurologically asymptomatic UK adult patient with haemophilia. *Haemophilia* 16:296–304
68. Peden AH, Head MW, Ritchie DL, Bell JE, Ironside JW (2004) Preclinical vCJD after blood transfusion in a PRNP codon 129 heterozygous patient. *Lancet* 364:527–529
69. Plassman BL, Williams JW Jr, Burke JR, Holsinger T, Benjamin S (2010) Systematic review: factors associated with risk for and possible prevention of cognitive decline in later life. *Ann Intern Med* 153:182–193
70. Prado MA, Baron G (2012) Seeding plaques in Alzheimer's disease. *J Neurochem* 120:641–643
71. Prusiner SB (1982) Novel proteinaceous infectious particles cause scrapie. *Science* 216:136–144
72. Prusiner SB (1998) Prions. *Proc Natl Acad Sci* 95:13363–13383
73. Prusiner SB (1998) The prion diseases. *Brain Pathol* 8:499–513
74. Rafalowska J, Barcikowska M, Wen GY, Wisniewski HM (1988) Laminar distribution of neuritic plaques in normal aging, Alzheimer's disease and Down's syndrome. *Acta Neuropathol* 77:21–25
75. Recasens A, Dehay B, Bove J et al (2014) Lewy body extracts from Parkinson disease brains trigger alpha-synuclein pathology and neurodegeneration in mice and monkeys. *Ann Neurol* 75:351–362
76. Ridley RM, Baker HF, Windle CP, Cummings RM (2006) Very long term studies of the seeding of beta-amyloidosis in primates. *J Neural Transm* 113:1243–1251
77. Roostae A, Beaudoin S, Staskevicius A, Roucou X (2013) Aggregation and neurotoxicity of recombinant alpha-synuclein aggregates initiated by dimerization. *Mol Neurodegener* 8:5
78. Rosen RF, Fritz JJ, Dooyema J et al (2012) Exogenous seeding of cerebral beta-amyloid deposition in betaAPP-transgenic rats. *J Neurochem* 120:660–666
79. Saborio GP, Permanne B, Soto C (2001) Sensitive detection of pathological prion protein by cyclic amplification of protein misfolding. *Nature* 411:810–813
80. Sanders DW, Kaufman SK, DeVos SL et al (2014) Distinct tau prion strains propagate in cells and mice and define different tauopathies. *Neuron* 82:1271–1288
81. Schulz-Schaeffer WJ (2010) The synaptic pathology of alpha-synuclein aggregation in dementia with Lewy bodies, Parkinson's disease and Parkinson's disease dementia. *Acta Neuropathol* 120:131–143
82. Schulz-Schaeffer WJ (2012) Neurodegeneration in Parkinson disease: moving Lewy bodies out of focus. *Neurology* 79:2298–2299
83. Schulz-Schaeffer WJ (2014) Morbus Parkinson—a synaptic disorder? *Basal Ganglia* 4:3–8
84. Serrano-Pozo A, Frosch MP, Masliah E, Hyman BT (2011) Neuropathological alterations in Alzheimer disease. *Cold Spring Harb Perspect Med* 1:a006189
85. Snowdon DA (1997) Aging and Alzheimer's disease: lessons from the Nun study. *Gerontologist* 37:150–156
86. Snowdon DA (2003) Healthy aging and dementia: findings from the Nun study. *Ann Intern Med* 139:450–454
87. Soto C (2011) Prion hypothesis: the end of the controversy? *Trends Biochem Sci* 36:151–158
88. Stöhr J, Watts JC, Mensinger ZL et al (2012) Purified and synthetic Alzheimer's amyloid beta (Aβ) prions. *Proc Natl Acad Sci* 109:11025–11030
89. Tyas SL, Snowdon DA, Desrosiers MF, Riley KP, Markesbery WR (2007) Healthy ageing in the Nun study: definition and neuropathologic correlates. *Age Ageing* 36:650–655
90. van Keulen LJ, Vromans ME, van Zijderveld FG (2002) Early and late pathogenesis of natural scrapie infection in sheep. *APMIS* 110:23–32
91. Volpicelli-Daley LA, Luk KC, Patel TP et al (2011) Exogenous alpha-synuclein fibrils induce Lewy body pathology leading to synaptic dysfunction and neuron death. *Neuron* 72:57–71
92. Watts JC, Giles K, Oehler A et al (2013) Transmission of multiple system atrophy prions to transgenic mice. *Proc Natl Acad Sci* 110:19555–19560
93. Weber P, Giese A, Piening N et al (2006) Cell-free formation of misfolded prion protein with authentic prion infectivity. *Proc Natl Acad Sci* 103:15818–15823
94. Wemheuer W, Benestad S, Wrede A et al (2009) Similarities between forms of sheep scrapie and Creutzfeldt–Jakob disease are encoded by distinct prion types. *Am J Pathol* 175:2566–2573
95. Wroe SJ, Pal S, Siddique D et al (2006) Clinical presentation and pre-mortem diagnosis of variant Creutzfeldt–Jakob disease associated blood transfusion: a case report. *Lancet* 368:2061–2067
96. Yamamoto N, Matsuzaki K, Yanagisawa K (2005) Cross-seeding of wild-type and hereditary variant-type amyloid beta-proteins in the presence of gangliosides. *J Neurochem* 95:1167–1176
97. Zahn R, Liu A, Lührs T et al (2000) NMR solution structure of the human prion protein. *Proc Natl Acad Sci* 97:145–150