Short-term results of hip revisions with a curved cementless modular stem in association with the surgical approach

Bernd Fink · Alexandra Grossmann ·
Svenja Schubring · Martin S. Schulz · Martin Fuerst

Published online: 26 November 2008
© Springer-Verlag 2008

Erratum to: Arch Orthop Trauma Surg DOI 10.1007/s00402-008-0617-7

There was an error in the column heads of Table 2, where “Trendelenburg limping” appeared instead of the correct “Trendelenburg sign”.

The corrected table is given in full here.

Table 2 Harris Hip Score und frequency of Trendelenburg signs in Revitan Curved stems implanted endofemoral and transfemoral

Time	Harris Hip Score			Trendelenburg sign		
	Endofem. <i>N</i> = 42	Transfem. <i>N</i> = 78	<i>P</i> value Mann–Whitney	Endofem. <i>N</i> = 42	Transfem. <i>N</i> = 78	<i>P</i> value Chi ² test
Preoperative	48.3 ± 15.5	44.1 ± 16.3	<i>P</i> = 0.226	23 (54.8%)	43 (55.1%)	<i>P</i> = 0.969
3 months	72.2 ± 14.9	61.0 ± 14.3	<i>P</i> < 0.001	28 (66.6%)	68 (87.2%)	<i>P</i> = 0.007
6 months	83.6 ± 19.9	72.1 ± 15.1	<i>P</i> = 0.001	21 (50%)	58 (74.3%)	<i>P</i> = 0.007
9 months	91.4 ± 11.3	78.8 ± 18.6	<i>P</i> = 0.001	18 (42.8%)	51 (65.4%)	<i>P</i> = 0.017
12 months	92.3 ± 11.6	81.4 ± 16.9	<i>P</i> = 0.002	18 (42.8%)	49 (62.8%)	<i>P</i> = 0.035
18 months	92.2 ± 12.8	84.7 ± 15.3	<i>P</i> = 0.046	15 (35.7%)	45 (57.7%)	<i>P</i> = 0.021
24 months	92.9 ± 12.1	86.3 ± 15.6	<i>P</i> = 0.067	11 (26.2%)	41 (52.6%)	<i>P</i> = 0.005

The online version of the original article can be found under
doi:[10.1007/s00402-008-0617-7](https://doi.org/10.1007/s00402-008-0617-7).

B. Fink (✉) · A. Grossmann · M. S. Schulz
Clinic of Joint Replacement,
General and Rheumatic Orthopaedics,
Orthopaedic Clinic Markgröningen gGmbH,
Kurt-Lindemann-Weg 10, 71706 Markgröningen, Germany
e-mail: b.fink@okm.de

S. Schubring · M. Fuerst
Orthopaedic Department of Rheumaklinik Bad Bramstedt,
Oskar-Alexander-Str. 26, 24576 Bad Bramstedt, Germany