Correction to: Fewer patients undergo surgery when adding a comprehensive geriatric assessment in older patients with a hip fracture

Babette C. van der Zwaard¹ · Charlotte E. Stein² · Janet E. M. Bootsma² · Hendrikus J. A. A. van Geffen³ · Conny M. Douw¹ · Carolina J. P. W. Keijsers²

Published online: 30 November 2019

© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to: Archives of Orthopaedic and Trauma Surgery
<https://doi.org/10.1007/s00402-019-03294-5>

The original version of this article unfortunately contained a mistake. The presentation of Fig. 1 was incorrect. The correct version of Fig. 1 is given below.

Fig. 1 Flowchart of inclusions in the study: all patients are ≥ 70 years with a hip fracture and received a pre-operative comprehensive geriatric assessment (CGA) or not

The original article can be found online at <https://doi.org/10.1007/s00402-019-03294-5>.

✉ Babette C. van der Zwaard
b.v.d.zwaard@jbz.nl

¹ Department of Orthopedic Surgery, Jeroen Bosch Hospital, PO Box 90153, 5200 ME 's-Hertogenbosch, The Netherlands

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

² Department of Geriatric Medicine, Jeroen Bosch Hospital, 's-Hertogenbosch, The Netherlands

³ Department of Surgery, Jeroen Bosch Hospital, 's-Hertogenbosch, The Netherlands