Correction to: Osseoperception in transcutaneous osseointegrated prosthetic systems (TOPS) after transfemoral amputation: a prospective study

Marcus Örgel¹ · Mohamed Elareibi¹ · Tilman Graulich¹ · Christian Krettek¹ · Claudia Neunaber¹ · Horst-Heinrich Aschoff¹ · Alexander Ranker² · Marcel Winkelmann¹

Published online: 17 December 2021

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Correction to: Archives of Orthopaedic and Trauma Surgery
<https://doi.org/10.1007/s00402-021-04099-1>

The original version of this article unfortunately contained a mistake. In Table 4, Männlich and Weiblich should be changed into male and female.

The corrected Table 4 is placed in the following page.

The original article can be found online at <https://doi.org/10.1007/s00402-021-04099-1>.

✉ Marcus Örgel
Oergel.Marcus@mh-hannover.de

¹ Department of Trauma, Hannover Medical School (MHH), Carl-Neuberg-Straße 1, 30625 Hannover, Germany

² Department of Physical Medicine and Rehabilitation, Hannover Medical School (MHH), Carl-Neuberg-Str. 1, 30625 Hannover, Germany

Table 4 Presentation of the subgroup analysis divided according to gender

	Male (<i>n</i> = 46)	Female (<i>n</i> = 29)	<i>p</i> value
Amputation side <i>n</i> (%) ^{##}			
Right	27 (58.7)	16 (55.2)	0.8 ^a
Left	19 (41.3)	17 (44.8)	
Reason for amputation <i>n</i> (%)			
Trauma	26 (56.5)	10 (34.5)	
Vascular	5 (10.9)	4 (13.8)	
Tumor	3 (6.5)	2 (6.9)	
No amputation	12 (26.1)	13 (48.2)	
Age mean ± SD, 95% CI, median (IQR) [#]	52.2 ± 12.0, 48.0–56.4, 51.5 (15)	49.8 ± 9.7, 44.6–54.9, 50 (12)	0.05 ^b
Years since amputation mean ± SD, 95% CI, Median (IQR) [#]	15.9 ± 11.4, 11.9–19.9, 16 (16)	16.4 ± 11.5, 10.3–22.5, 15.5 (19)	0.9 ^b
BMI [kg/m ²] mean ± SD, 95%-CI, median (IQR) [#]	26.5 ± 4.0, 25.2–27.9, 26.3 (5)	28.5 ± 5.4, 25.6–31.4, 27.4 (6)	0.7 ^b
V1 mean ± SD, 95% CI, median (IQR) ^{##}	2.0 ± 1.3, 1.6–2.5, 2.0 (2)	1.7 ± 1.1, 1.1–2.3, 2 (1)	0.9 ^a
V2 mean ± SD, 95% CI, median (IQR) ^{##}	1.4 ± 1.1, 1.0–1.7, 1.0 (2)	1.3 ± 1.3, 0.7–2.0, 1 (3)	0.6 ^a
V3 mean ± SD, 95% CI, median (IQR) ^{##}	2.9 ± 0.3, 2.8–3.1, 3.0 (0)	3 (constant)	0.4 ^a
Total V1–V3 mean ± SD, 95%-CI, median (IQR) [#]	6.2 ± 1.7, 5.6–6.8, 6 (3)	6.0 ± 1.8, 5.1–7.0, 6 (4)	0.5 ^b
PMQ mean ± SD, 95% CI, median (IQR) ^{##}	24.4 ± 21.0–27.8, 25.0 (17)	27.5 ± 9.3, 22.3–32.6, 29.0 (18)	0.3 ^b
K-Level mean ± SD, 95% CI, median (IQR) [#]	2.6 ± 0.7, 2.3–2.9, 3.0 (1)	2.6 ± 0.8, 2.1–3.1, 3.0 (1)	0.6 ^a
Prosthesis wear time mean ± SD, 95% CI, median (IQR) [#]	11.1 ± 6.1, 9.0–13.2, 11.0 (7)	11.1 ± 5.3, 8.1–14.0, 14.0 (7)	0.8 ^a

[#]Normally distributed data^{##}Non-normally distributed data^aMann–Whitney *U* test^b*T* test

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.