


CORRECTION

Correction to: Digestive system in psoriasis: an update

Daniel Pietrzak¹ · Aldona Pietrzak² · Dorota Krasowska² · Andrzej Borzęcki³ ·
Kinga Franciszkiewicz-Pietrzak⁴ · Beata Polkowska-Pruszyńska⁵  ·
Maja Baranowska⁶ · Kristian Reich⁷

Published online: 10 October 2017
© Springer-Verlag GmbH Germany 2017

Correction to: Arch Dermatol Res
DOI 10.1007/s00403-017-1775-7

In the original publication, the data labels are incorrect in Fig. 3. The corrected Fig. 3 is given here.

The online version of the original article can be found under
doi:[10.1007/s00403-017-1775-7](https://doi.org/10.1007/s00403-017-1775-7).

✉ Beata Polkowska-Pruszyńska
polkowska.beata@gmail.com

¹ Department of Anaesthesiology and Intensive Care, Medical University of Lublin, Lublin, Poland

² Chair and Department of Dermatology, Venereology and Paediatric Dermatology, Medical University of Lublin, Lublin, Poland

³ Chair and Department of Hygiene, Medical University of Lublin, Lublin, Poland

⁴ Department of Surgical Oncology, Medical University of Lublin, Lublin, Poland

⁵ Student's Scientific Association at the Department of Dermatology, Venereology and Pediatric Dermatology, Medical University of Lublin, Lublin, Poland

⁶ Institute of Fine Arts University of Warmia and Mazury in Olsztyn, Olsztyn, Poland

⁷ Dermatologikum Hamburg, Hamburg, Germany

Fig. 3 Mechanisms enhancing an increase of intestinal permeability in inflammatory bowel disease IBD involve several group of cells—dendritic cells “in situ in the intestinal wall”, T-regulatory lymphocytes (T-reg), Th17 cells, Th1 cells, which are responsible for producing cytokines such as IFN γ , INF α , IL-6, IL-12, IL-23 and IL-17. Based on Vlachos et al. [167]

