Correction to: Combination of early pushing with extended second stage increases the rates of spontaneous vaginal deliveries, but might be associated with adverse maternal and neonatal outcomes

Asmaa Masri¹ · Lelia Abu-Nasra² · Reuven Kedar² · Amit Damti² · Moti Bardicef² · Nir Kugelman² · Lena Sagi-Dain^{1,2}

Published online: 10 July 2021

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Correction to: Archives of Gynecology and Obstetrics
<https://doi.org/10.1007/s00404-021-06111-w>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

In the original article published, the given name and family name of all the authors are interchanged.

The correct given name and family name are given below:

Asmaa Masri, Lelia Abu-Nasra, Reuven Kedar, Amit Damti, Moti Bardicef, Nir Kugelman and Lena Sagi-Dain

The original article can be found online at <https://doi.org/10.1007/s00404-021-06111-w>.

✉ Lena Sagi-Dain
lana2303@gmail.com

¹ The Ruth and Bruce Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel

² Department of Obstetrics and Gynecology, Carmel Medical Center, 7 Michal St, Haifa, Israel