Ericksonian hypnosis in tinnitus therapy: effects of a 28-day inpatient multimodal treatment concept measured by Tinnitus-Questionnaire and Health Survey SF-36

U. H. Ross · O. Lange · J. Unterrainer · R. Laszig

Published online: 15 March 2007
© Springer-Verlag 2007

Erratum to: Eur Arch Otorhinolaryngol
DOI 10.1007/s00405-006-0221-9

Table 1 unfortunately contained minor errors regarding the references in brackets. The correct version is shown below.

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s00405-006-0221-9>.

U. H. Ross (✉)
Practice for Otorhinolaryngology and Psychotherapy,
Luisenstrasse 6, 79098 Freiburg, Germany
e-mail: Dr-Ross@web.de

O. Lange
Institute of Environmental Medicine and Hospital Epidemiology,
University of Freiburg, Freiburg, Germany

J. Unterrainer
Department of Neuropsychology,
University of Freiburg, Freiburg, Germany

R. Laszig
Department of Otorhinolaryngology,
University of Freiburg, Freiburg, Germany

Table 1 Components of the treatment concept

Procedure	Sessions	Total time
Education Tinnitus etiology and maintenance, bio-psycho-social concept of salutogenesis, sleep	6 × 90 min (group)	540 min
Ericksonian Hypnosis - therapeutic metaphors and symbolization [7]	4 × 90 min/week (group)	1440 min
- anchoring and self-relation techniques [8, 23]	2 × 60 min/week	480 min
- systematic change of time focus (time-line techniques) [6]	(individual)	
- ideodynamic techniques [22]		
Jacobson relaxation training	2 × 60 min/week (group)	480 min
Music therapy	1 × 60 min/week (group)	240 min