Correction to: Increased incidence of Warthin tumours of the parotid gland: a 42-year evaluation

Achim M. Franzen¹ · Christiane Kaup Franzen² · Thomas Guenzel³ · Anja Lieder¹

Published online: 27 August 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to:
European Archives of Oto-Rhino-Laryngology
<https://doi.org/10.1007/s00405-018-5092-3>

It should read “265” instead of the “235”. The corrected Table 2 is given below.

In the original publication, in Table 2, the last value of third column, referring to ‘Number of cases’ has been misprint.

Table 2 Gender distribution of Warthin tumours

Author and year	Country	Number of cases	Male-to-female ratio	Development of male-to-female ratio within 40-year period
Thompson 1950 [10]	United States of America	160	7–1	
Chaudhry 1959 [26]	United States of America	306	5–1	
Seifert 1980 [27]	Germany	275	3–1	
Eveson 1986 [28]	United Kingdom	278	1.6–1	
Yoo 1994 [15]	United States of America	119	2.2–1	3.8 to 1–1.6–1
Mortensen 2008 [4]	Denmark		4.9–1	
Tian 2010 [5]	China (Eastern)	957	11.2–1	
Chulam 2013 [29]	Brazil	100	2.6–1	
Witt 2015 [21]	United States of America	50	2.1–1	
Patel 2016 [13]	New Zealand	41	2–1	
This study	Germany	265	2.6–1	5.3–1–2.1–1

The original article can be found online at <https://doi.org/10.1007/s00405-018-5092-3>.

✉ Anja Lieder
anja_lieder@hotmail.com

¹ Department of Otorhinolaryngology and Head and Neck Surgery, Brandenburg Medical University-Theodor Fontane, Campus Ruppiner Kliniken, Fehrbelliner Strasse 38, 16816 Neuruppin, Germany

² Academic Teaching practice of the Brandenburg Medical School-Theodor Fontane, Dorfstrasse 34, Walsleben, Germany

³ Department of Otorhinolaryngology and Head and Neck Surgery, Borromaeus-Hospital, Leer, Germany