



# Correction to: Preoperative risk assessment for ambulatory sinonasal surgery

Hans Rudolf Briner<sup>1</sup> · Andreas Leunig<sup>2</sup> · Christoph Schlegel<sup>3</sup> · Daniel Simmen<sup>1</sup>

© The Author(s) 2021

**Correction to: European Archives of Oto-Rhino-Laryngology**  
<https://doi.org/10.1007/s00405-020-06435-4>

In the original publication of the article, under the abstract section, the subsections “Ethical consideration” and “Study design and patients” was included incorrectly. The correct abstract includes the subsections Objectives, Design, Setting, Participants, Main outcome measures, Results and Conclusions.

In addition, under the material and methods section, the subsections “Ethical consideration” and “Study design and patients” was published incorrectly. The correct text should read as below,

## Material and methods

### Ethical considerations

The study was approved by the Ethics Committee of the Canton Zurich, Switzerland.

### Study design and patients

In a prospective multicentric interventional study, patients where endoscopic sinus procedures were planned at the Center for Otorhinolaryngology, Head and Neck Surgery, Klinik Hirslanden, Zurich, the Department of

Otolaryngology, Head and Neck Surgery, Luzerner Kantonsspital, Lucerne and HNO-Klinik München-Bogenhausen, Munich, in the period between January 1st 2017 and December 31st 2018 were evaluated. Due to the structure of the health care system in these centers, patients were admitted with a minimal hospital stay of 24 h. Patients below the age of 18 were excluded. Patient specific data (patient age, gender), the type of the procedure, the preoperative risk score or “day surgery risk score”, postoperative complications during the first 14 days and the length of hospital stay were analyzed. Data collection was performed using “ENTstatistics” software (ENTstatistics, Innoforce, LI-9491 Ruggel, Liechtenstein).

The original article has been updated.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

The original article can be found online at <https://doi.org/10.1007/s00405-020-06435-4>.

✉ Hans Rudolf Briner  
briner@orl-zentrum.com

<sup>1</sup> Center for Otorhinolaryngology, Head and Neck Surgery, Klinik Hirslanden, Witellikerstrasse 40, 8032 Zurich, Switzerland

<sup>2</sup> Rhinology Center Munich and ENT-Clinic Dr. Gaertner GmbH, Possartstr. 27-31, 81679 Munich, Germany

<sup>3</sup> Department of Otorhinolaryngology, Head and Neck Surgery, Luzerner Kantonsspital, 6000 Lucerne 16, Switzerland

**Publisher’s Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.