



Correction to: Tumor-host interface in oral squamous cell carcinoma: Impact on nodal metastasis and prognosis

Rishikesh Thakur¹ · Alok Thakar¹ · Rajeev K. Malhotra² · Atul Sharma³ · Aanchal Kakkar⁴ 

Published online: 4 May 2021

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

**Correction to: European Archives of
Oto-Rhino-Laryngology**
<https://doi.org/10.1007/s00405-021-06756-y>

In the original publication of the article, under the Table 2, values of Recurrences and Loco-regional was published incorrectly. The correct Table 2 is provided below,

The original article can be found online at <https://doi.org/10.1007/s00405-021-06756-y>.

✉ Alok Thakar
drathakar@gmail.com

✉ Aanchal Kakkar
aanchalkakkar@gmail.com

¹ Department of Otorhinolaryngology and Head and Neck Surgery, All India Institute of Medical Sciences, Ansari Nagar, New Delhi 110029, India

² Delhi Cancer Registry, BR Ambedkar Institute Rotary Cancer Hospital, All India Institute of Medical Sciences, New Delhi, India

³ Department of Medical Oncology, All India Institute of Medical Sciences, New Delhi, India

⁴ Department of Pathology, All India Institute of Medical Sciences, Ansari Nagar, New Delhi 110029, India

Table 2 Comparison of Brandwein-Gensler histological risk score groups with clinicopathological parameters and oncologic outcome

	Low risk (<i>n</i> = 25)	Intermediate risk (<i>n</i> = 93)	High risk (<i>n</i> = 60)	<i>p</i> value ^a
Demographic parameters				
Mean age	53 years	48 years	44 years	0.030^b
Male: Female	23: 2	82: 11	50: 10	0.57
Tumor subsite	Group I: 3 Group II: 22	Group I: 32 Group II: 61	Group I: 32 Group II: 28	0.001
Tobacco consumption Yes: No	23: 2	86:7	53:7	0.440
Histopathological parameters				
Pathological T stage	T1: 4 T2: 9 T3: 6 T4: 6	T1: 17 T2: 30 T3: 19 T4: 27	T1: 10 T2: 22 T3: 9 T4: 19	0.95
Pathological node stage	N0: 24 N1: 1 N2: 0	N0: 59 N1: 14 N2: 20	N0: 4 N1: 35 N2: 21	0.000
Pathological TNM stage group (AJCC 8th edition)	I: 4 II: 9 III: 6 IV: 6	I: 13 II: 18 III: 22 IV: 40	I: 3 II: 1 III: 27 IV: 29	0.000
Histological grade	WDSCC: 20 MDSCC: 5 PDSCC: 0	WDSCC:36 MDSCC: 55 PDSCC: 2	WDSCC: 12 MDSCC: 44 PDSCC: 4	0.001
Margin status	Clear: 12 Close: 13 Involved: 0	Clear: 44 Close: 46 Involved: 3	Clear: 31 Close: 26 Involved: 3	0.760
Oncology Outcomes				
Recurrences	2 (8%)	32 (34.4%)	35 (58.3%)	0.001
Loco-regional	2	21	23	
Distant	Nil	11	12	
2-year DFS	96%	51.6%	15%	< 0.001^c
Early stage	100%	81.7%	25%	< 0.001^c
Late stage	91.7%	33.9%	14.3%	0.001^c
2-year OS	96%	66.7%	31.3%	< 0.001^c
Early stage	100%	90.3%	75%	0.214 ^c
Late stage	91.7%	54.8%	28.1%	< 0.001^c
2-year LRC	96%	57.3%	18.7%	< 0.001^c
2-year DSS	96%	67%	33%	< 0.001^c

WD well differentiated, MD moderately differentiated, PD poorly differentiated, DFS disease-free survival, LRC Locoregional Control, DSS disease specific survival, OS overall survival, Group I subsites Tongue, Floor of mouth, Group II subsites Alveolus, Buccal mucosa, retromolar trigone, Hard palate, lip

^a*p* values by Chi Square test except where indicated

^bANOVA

^clog-rank test; significant *p* values are in bold