



Correlations between IL-6 serum level and olfactory dysfunction severity in COVID-19 patients: a preliminary study

Luigi Angelo Vaira^{1,2} · Andrea De Vito³ · Giovanna Deiana^{2,4} · Chiara Pes^{5,6} · Federica Giovanditto^{1,7} · Vito Fiore³ · Jerome R. Lechien^{8,9} · Serge-Daniel Le Bon¹⁰ · Sven Saussez^{8,9} · Giordano Madeddu³ · Sergio Babudieri³ · Antonio Pazzola⁷ · Franco Bandiera^{5,6} · Alessandro Giuseppe Fois¹¹ · Andrea Fausto Piana^{2,4} · Claire Hopkins¹² · Giacomo De Riu¹

Received: 28 February 2021 / Accepted: 3 May 2021 / Published online: 13 May 2021
© The Author(s) 2021

Abstract

Background Interleukin 6 (IL-6) is a proinflammatory cytokine that is secreted by cells infected with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and it is widely recognized as a negative prognostic factor. The purpose of this study was to analyze the correlations between the olfactory scores determined by psychophysical tests and the serum levels of IL-6 in patients affected by coronavirus disease 2019 (COVID-19)

Methods Patients underwent psychophysical olfactory assessment with Connecticut Chemosensory Clinical Research Center test and IL-6 plasma level determination within 10 days of the clinical onset of COVID-19.

Results Seventy-four COVID-19 patients were included in this study. COVID-19 staged as mild in 34 patients, moderate in 26 and severe in 14 cases. There were no significant differences in olfactory scores across the different COVID-19 severity groups. In the patient series, the median plasma level of IL-6 was 7.7 pg/mL (IQR 3.7–18.8). The concentration of IL-6 was found to be significantly correlated with the severity of COVID-19 with a directly proportional relationship. The correlation between IL-6 plasma concentrations and olfactory scores was weak ($r_s = 0.182$) and not significant ($p = 0.12$).

Conclusions In COVID-19 patients, psychophysical olfactory scores did not show significant correlations with the plasma levels of a well-recognized negative prognostic factor such as IL-6. This observation casts some shadows on the positive prognostic value of olfactory dysfunctions.

Keywords COVID-19 · IL-6 · Interleukin 6 · Smell · Anosmia · Cytokine · SARS-CoV-2 · Coronavirus · Cytokine storm

Introduction

Olfactory dysfunctions are one of the most frequent clinical manifestations of coronavirus disease 2019 (COVID-19), affecting more than 70% of patients infected with SARS-CoV-2 [1–5].

Recently, the prognostic value of olfactory disorders has been the subject of heated debate. Although some

researchers have found no correlation between the prevalence of loss of smell and severity of COVID-19 [6–9], many report that olfactory dysfunction is more frequent in mild forms [10–13] and some postulate that this is the price to pay for a more effective immune response against the virus in the olfactory epithelium [14].

Interleukin 6 (IL-6) is a proinflammatory cytokine that is secreted by cells infected with SARS-CoV-2 [15]. IL-6 is one of the factors behind the cytokine storm that occurs in the most severe cases of COVID-19 and serum levels of this cytokine are therefore a widely recognized negative prognostic factor [16–18]. Before the pandemic, only a few authors have investigated the correlation between the severity of chronic olfactory disorders and serum levels of IL-6, noting a statistically significant directly proportional correlation [19, 20]. These results support a hypothesis that IL-6 may play a role in biochemical pathological process underlying

Luigi Angelo Vaira and Andrea De Vito equally contributed to this work and should be regarded as joint first authors.

Claire Hopkins and Giacomo De Riu equally contributed to this work and should be regarded as joint senior authors.

✉ Luigi Angelo Vaira
lavaira@uniss.it

Extended author information available on the last page of the article

these chronic olfactory dysfunctions [19]. However, this directly proportional correlation conflicts with what has been proposed in COVID-19 patients in whom the severity of olfactory dysfunction appears to be inversely proportional to the severity of COVID-19 [10–13] and therefore likely inversely proportional to serum IL-6 levels, assuming that serum IL-6 levels largely reflect the severity of COVID-19 pneumonia and other major organ involvement.

The purpose of this study was therefore to analyze the correlations between the olfactory scores determined by psychophysical tests and the serum levels of IL-6 in patients affected by COVID-19 and admitted to the University Hospital of Sassari. We hypothesize that severity of olfactory dysfunction will be inversely proportional to serum IL-6 levels, which are a marker of COVID-19 disease severity.

Materials and methods

This cohort observational study was conducted in the COVID departments of the University Hospital of Sassari (Infectious and Tropical Disease, Pneumology, Onco-COVID and Neuro-COVID operative units) between January 10 and February 1, 2021.

To be enrolled in the study, patients had to meet the following inclusion criteria: adults over 18 years of age, rhino-pharyngeal swab positive for SARS-CoV-2 infection, COVID-19 symptoms present for less than 10 days, patient acceptance for participation in the study. On the contrary, the study exclusion criteria were: uncooperative patients, assisted ventilation, psychiatric or neurological disorders, previous surgery or radiotherapy in the oral and nasal cavities, pre-existing self-reported smell and taste alterations, history of head trauma, allergic rhinitis, chronic rhinosinusitis.

All patients provided informed consent for participation in the study. The study protocol was approved by University Hospital of Cagliari Ethical Committee (PG/2021/5471).

Some clinical and epidemiological information was collected for all patients: age, gender and COVID-19 symptoms. All patients were followed up clinically until the nasopharyngeal swab was negative. The overall clinical severity of COVID-19 was classified according to Tian et al. [21] in mild, moderate, severe and critical.

Psychophysical olfactory evaluation was performed with the Connecticut Chemosensory Clinical Research Center test (CCCRC). The CCCRC is a validated, widely used and easy to perform psychophysical test. The methodology, the scoring system and its application in COVID-19 patients have been extensively described in previous studies [22–25]. The CCCRC includes the assessment of the olfactory threshold using solutions with increasing concentration of N-butyl acid and an identification task for common odors. The olfactory score thus

obtained allows to clinically classify the olfactory function in five categories: normal (scores 90 and 100), mild (scores 70 and 80), moderate (scores 50 and 60) or severe hyposmia (scores 20, 30 and 40) and anosmia (scores 0 and 10).

Within 24 h after the olfactory test, plasma levels of IL-6 (reference value < 5.9 pg/mL) were determined on a peripheral blood sample taken from each patient by means of a fully automated Elecsys system on a cobas e801 platform (Roche Diagnostics, Basel, Switzerland) as previously described [18].

The statistical analysis was performed with SPSS 26.0 (IBM, Armonk, NY, USA). Categorical variables are reported in numerals and percentages of the total. Descriptive statistics for quantitative variables are given as the mean \pm standard deviation (SD) or median (interquartile range–IQR). The Kruskal–Wallis Test was performed to evaluate the statistical significance of differences in olfactory scores between clinical severity groups. Post hoc analysis with Mann–Whitney *U* test was used to define the different relationships of the olfactory severity score and IL-6 blood levels with each of the COVID-19 severity subgroups. The correlation between olfactory scores and IL-6 levels was assessed with the Spearman rank correlation coefficient. The level of statistical significance was set at $p < 0.05$ with a 95% confidence interval.

Results

Seventy-four COVID-19 patients who met the inclusion and exclusion criteria were included in this study. Table 1 summarizes the epidemiological and clinical characteristics of the patients (Table 1).

COVID-19 was staged as mild in 34 patients, moderate in 26 and severe in 14 cases. At the time of psychophysical evaluation, 55 patients (74% of the cohort) presented with olfactory dysfunction including anosmia (14 cases; 18.9%), severe hyposmia (12 cases; 16.2%), moderate hyposmia (18 cases; 24.3%) and mild hyposmia (11 cases; 14.9%), the remainder having normosmia. Overall, the median olfactory score was 60 (IQR 30–87.5). There were no significant differences in olfactory scores across the different COVID-19 severity groups (Tables 2, 3).

In the patient series, the median plasma level of IL-6 was 7.7 pg/mL (IQR 3.7–18.8). The concentration of IL-6 was found to be significantly correlated with the severity of COVID-19 with a directly proportional relationship ($p < 0.001$) (Table 2). Post hoc analysis with Mann–Whitney *U* test post hoc analysis showed that IL-6 blood concentration differences were significantly different across all COVID-19 severity subgroups analyzed (Table 3). The correlation between IL-6 plasma concentrations and olfactory scores was weak ($r_s = 0.182$) and not significant ($p = 0.12$) (Fig. 1).

Table 1 General and clinical features of the study population

Gender	
Male	47 (63.5%) [95% CI 51.5–74.4%]
Female	27 (36.5%) [95% CI 25.6–48.5%]
Age (years)	63.4 ± 13.4 [95% CI 60.9–66.5]
Mean ± SD	
Days from COVID-19 symptoms onset	7.2 ± 3.2 [95% CI 6.5–7.9]
Mean ± SD	
Clinical stage	
No. of patients (%)	
Mild	34 (46%) [95% CI 34.3–57.9%]
Moderate	26 (35.1%) [95% CI 24.4–47.1%]
Severe	14 (18.9%) [95% CI 10.7–29.7%]
Critical	0 (0%) [97.5% CI 0–4.7%]
Olfactory function assessment	
No. of patients (%)	
Normal	19 (25.7%) [95% CI 16.2–37.2%]
Mild hyposmia	11 (14.9%) [95% CI 7.6–25%]
Moderate hyposmia	18 (24.3%) [95% CI 15.1–35.7%]
Severe hyposmia	12 (16.2%) [95% CI 8.7–26.6%]
Anosmia	14 (18.9%) [95% CI 10.7–29.7%]

95% CI 95% confidence interval, SD standard deviation

Discussion

To the best of our knowledge, only one article has previously evaluated the correlation between olfactory dysfunction in COVID-19 and IL-6 levels [26]. In this study, Cazzolla et al. reported a significant directly proportional correlation between IL-6 levels and the presence of self-reported olfactory dysfunction. This finding seems to be in line with what has been reported for other chronic inflammatory olfactory dysfunctions [19, 20] but, considering that high levels of IL-6 are a well-established negative prognostic factor [16–18], it would be in contrast to the fact that olfactory disorders seem to represent a favorable prognostic index in COVID-19 patients [10–13]. However, the study of Cazzolla et al. presents some major limitations which reduce the reliability of the results and that we have tried to overcome

in the present study. First, the assessment of the olfactory function is not based on psychophysical tests and it has been shown that self-reported olfactory loss alone significantly underestimates the real prevalence of smell disorders in COVID-19 patients [27, 28]. Second, the temporal distance from the onset of symptoms was not considered among the inclusion and exclusion criteria. This is crucial if we want to establish whether the presence and severity of the olfactory disturbance at the onset of COVID-19 can correlate with other established prognostic indices. In this sense, a further limitation is that only mild and moderate COVID-19 patients were included in the study by Cazzolla et al. [26].

In our study, patients were evaluated within the first 10 days of symptom onset, when olfactory dysfunction should not yet have begun to recover [29, 30]. At this early stage of COVID-19, 74.3% of patients had olfactory dysfunction. This prevalence is in line with those reported by most authors [1, 3–5, 8, 13, 22, 23, 25]. The olfactory scores did not show significant differences between the various COVID-19 severity groups. Although most authors attribute a positive prognostic value to olfactory disorders [10–13], this finding has never been detected in patients belonging to the Italian centers who participated in our prospective cohort studies during the first wave [7, 23–25]. However, plasma levels of IL-6 were shown to be a reliable negative prognostic index, demonstrating a directly proportional and significant correlation with the

Table 3 Post hoc analysis results

	Mann–Whitney <i>U</i> test
CCCRC score	
Mild versus moderate COVID-19	<i>p</i> = 0.787
Moderate versus severe COVID-19	<i>p</i> = 0.153
Mild versus severe COVID-19	<i>p</i> = 0.082
IL-6 blood level	
Mild versus moderate COVID-19	<i>p</i> = 0.008
Moderate versus severe COVID-19	<i>p</i> = 0.002
Mild versus severe COVID-19	<i>p</i> < 0.001

Table 2 Olfactory scores and IL-6 level according to clinical severity of COVID-19

COVID-19 severity [21]	Mild (<i>N</i> = 34)	Moderate (<i>N</i> = 26)	Severe (<i>N</i> = 14)
CCCRC score	60 (30–70)	50 (32.5–85)	85 (50–90)
Median (IQR)			
Kruskal–Wallis test	<i>p</i> = 0.202		
IL-6 blood level (pg/mL)	4.29 (3.5–10.6)	10.5 (6–18.2)	31.5 (17.8–44.3)
Median (IQR)			
Kruskal–Wallis test	<i>p</i> < 0.001		

CCCRC Connecticut Chemosensory Clinical Research Center test, IL-6 interleukin 6, IQR interquartile range




Fig. 1 Correlation analysis between IL-6 plasma level and CCCRC scores

severity of the clinical picture of COVID-19. The differences between the median IL-6 blood level were significant not only between patients with severe COVID-19 and those with mild and moderate forms but also between subjects with mild and moderate clinical pictures. This supports our hypothesis that serum IL-6 levels reflect the severity of major organ involvement. The correlation between olfactory scores and IL-6 levels was therefore weak and not significant.

Our findings give support to two main hypotheses. First, the prognostic value of olfactory disturbances appears to be significantly less strong than other clinical and laboratory data such as IL-6. The data presented in the literature to date are discordant, often retrospective and based on the only self-reported olfactory loss or with the need for hospitalization as the only prognostic outcome. Second, the pathogenesis of olfactory dysfunctions in COVID-19 is more likely caused by a local inflammatory [31–33] process than a systemic cytokine storm that then damages the olfactory epithelium. In this inflammatory process, it is not excluded that locally produced IL-6 may play a role in the genesis of olfactory dysfunction by acting on neurons [34] or glial cells [35] as occurs in other inflammatory and post-viral smell disorders [19, 36]. It would have been interesting to determine IL-6 levels in nasal mucus but this was not possible due to technical problems at the time of this study. However, any nasal production of IL-6 within the olfactory

epithelium is likely insufficient to significantly contribute to serum IL-6 levels.

This study has some limitations. The number of patients included in the study is not yet sufficient to draw firm conclusions. While there is a risk we have rejected significant association between olfactory scores and IL-6 levels due to type 2 statistical error, we did have sufficient participants to detect a strong and statistically significant correlation between COVID-19 severity and IL-6 which is a well-recognized negative prognostic factor. In future studies, it would be useful to evaluate the correlations between olfactory scores and both serum and mucus levels of other pro-inflammatory cytokines such as IL-10, IL-12, IL-15 and TNF- α which are secreted by cells infected with SARS-CoV-2 [37] and involved in the cytokine storm at the base of the most severe cases of COVID-19 [38]. Moreover, it could be interesting to prospectively evaluate patients by monitoring the recovery of olfactory function and correlating it with IL-6 levels in the early stages of infection. It is well known that IL-6 can be both pro and anti-inflammatory and suppresses the production of TNF- α [39]. TNF- α is increased in the olfactory mucosa following SARS-CoV-2 infection and causes damage to olfactory sensory neurons (OSNs) [40]. Perhaps elevated systemic IL-6 leads to reduced TNF- α production in olfactory epithelium and limits damage to OSNs, therefore higher IL-6 early on could be associated with faster recovery of olfactory loss. We may not have been able to see

this correlation as we tested patients in the earliest stages of the disease, during the phase of injury to the supporting cells but before the secondary loss of OSNs has occurred.

Conclusions

In COVID-19 patients, psychophysical olfactory scores did not show significant correlations with the plasma levels of a well-recognized negative prognostic factor such as IL-6. This observation casts some shadows on the positive prognostic value of olfactory dysfunctions.

Author contributions LAV, ADV and GDR have full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Conceptualization and methodology: LAV, ADV, GDR. Acquisition, analysis and interpretation of the data: CP, FG, VF, GD. Statistical analysis: AFP, GD. Critical revision of the literature: GM, SB, AGF, FB, AP. Provision of resources: SB, AGF, FB, AP. Original draft preparation: LAV, ADV, GDR. Critical revision of the manuscript for important intellectual content: CH, SS, JRL, SLB. Supervision: GDR, AFP. All the authors read and approved the final version of the manuscript.

Funding Open access funding provided by Università degli Studi di Sassari within the CRUI-CARE Agreement.

Declarations

Conflict of interest None of the authors has a financial interest in any of the products, devices or drugs mentioned in this manuscript.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Petrocelli M, Ruggiero F, Baietti AM, Pandolfi P, Salzano G, Salzano FA et al (2020) Remote psychophysical evaluation of olfactory and gustatory functions in early-stage coronavirus disease 2019 patients: the Bologna experience of 300 cases. *J Laryngol Otol* 134:571–576
- Parma V, Ohla K, Veldhuizen MG, Niv MY, Kelly CE, Bakke AJ et al (2020) More than smell—COVID-19 is associated with severe impairment of smell, taste, and chemesthesis. *Chem Senses* 45:609–622
- Vaira LA, Lechien JR, Khalife M, Petrocelli M, Hans S, Distinguin L et al (2021) Psychophysical evaluation of the olfactory function: European multi-center study on 774 COVID-19 patients. *Pathogens* 10:E62. <https://doi.org/10.3390/pathogens10010062>
- Spinato G, Fabbris C, Polesel J et al (2020) Alterations in smell or taste in mildly symptomatic outpatients with SARS-CoV-2 infection. *JAMA* 323(20):2089–2090
- Sedaghat AR, Gengler I, Speth MM (2020) Olfactory dysfunction: a highly prevalent symptom of COVID-19 with public health significance. *Otolaryngol Head Neck Surg* 163(1):12–15
- Moein ST, Hahemian SMR, Mansourafshar B, Khorrarn-Tousi A, Tabarsi P, Doty RL (2020) Smell dysfunction: a biomarker for COVID-19. *Int Forum Allergy Rhinol* 10(8):944–950
- Vaira LA, Hopkins C, Petrocelli M et al (2020) Do olfactory and gustatory psychophysical scores have prognostic value in COVID-19 patients? A prospective study of 106 patients. *J Otolaryngol Head Neck Surg* 49(1):56
- Speth MM, Singer-Cornelius T, Oberle M, Gengler I, Brockmeier SJ, Sedaghat AR (2020) Olfactory dysfunction and sinonasal symptomatology in COVID-10: prevalence, severity, timing, and associated characteristics. *Otolaryngol Head Neck Surg* 163(1):114–120
- Hopkins C, Vaira LA, De Riu G (2020) Self-reported olfactory loss in COVID-19: is it really a favorable prognostic factor? *Int Forum Allergy Rhinol* 10:926
- Lechien JR, Ducarme M, Place S et al (2020) Objective olfactory findings in hospitalized severe COVID-19 patients. *Pathogens* 9(8):E627
- Yan CH, Faraji F, Prajapati DP, Ostrander BT, DeConde AS (2020) Self-reported olfactory loss associates with outpatient clinical course in COVID-19. *Int Forum Allergy Rhinol* 10(7):821–831
- Izquierdo-Dominguez A, Rojas-Lechuga MJ, Chiesa-Estomba C et al (2020) Smell and taste dysfunctions in COVID-19 are associated with younger age in ambulatory settings—a multicenter cross-sectional study [published online June 17, 2020]. *J Investig Allergol Clin Immunol*. <https://doi.org/10.18176/jiaci.0595>
- Lechien JR, Chiesa-Estomba CM, Vaira LA, De Riu G, Cammaroto G, Idrissi YC et al (2020) Epidemiological, otolaryngological, olfactory and gustatory outcomes according to the severity of COVID-19: a study of 2579 patients. *Eur Arch Otorhinolaryngol*. <https://doi.org/10.1007/s00405-020-06548>
- Le Bon SD, Horoi M (2020) Is anosmia the price to pay in an immune-induced scorched-earth policy against COVID-19? *Med Hypotheses* 143:109881. <https://doi.org/10.1016/j.mehy.2020.109881>
- Coomes EA, Haghbayan H (2020) Interleukin-6 in COVID-19: a systematic review and meta-analysis. *Rev Med Virol* 30:1–9
- Liu F, Li L, Xu M, Wu J, Luo D, Zhu Y et al (2020) Prognostic value of interleukin-6, C-reactive protein, and procalcitonin in patients with COVID-19. *J Clin Virol* 127:104370
- Aziz M, Fatima R, Assaly R (2020) Elevated interleukin-6 and severe COVID-19: a meta-analysis. *J Med Virol* 92:2283–2285
- Herold T, Jurinovic V, Arnreich C, Lipworth BJ, Hellmuth JC, von Bergwelt-Baildon M et al (2020) Elevated levels of IL-6 and CRP predict the need for mechanical ventilation in COVID-19. *J Allergy Clin Immunol* 146:128–136
- Henkin RI, Schmidt L, Velicu I (2013) Interleukin 6 in hyposmia. *JAMA Otolaryngol Head Neck Surg* 139:728–734
- Raff AC, Lieu S, Melamed JL, Melamed ML, Quan Z, Ponda M et al (2008) Relationship of impaired olfactory function in ERS to malnutrition and retained uremic molecules. *Am J Kidney Dis* 52:102–110
- Tian S, Hu N, Lou J et al (2020) Characteristics of COVID-19 infection in Beijing. *J Infect* 80:401–406
- Vaira LA, Deiana G, Fois AG, Pirina P, Madeddu G, De Vito A et al (2020) Objective evaluation of anosmia and ageusia in

- COVID-19 patients: single-center experience on 72 cases. *Head Neck* 42:1252–1258
23. Vaira LA, Hopkins C, Salzano G, Petrocelli M, Melis A, Cucurullo M et al (2020) Olfactory and gustatory function impairment in COVID-19 patients: Italian objective multicenter-study. *Head Neck* 42:1560–1569
 24. Vaira LA, Hopkins C, Petrocelli M, Lechien JR, Cutrupi S, Salzano G et al (2020) Efficacy of corticosteroid therapy in the treatment of long-lasting olfactory disorders in COVID-19 patients. *Rhinology*. <https://doi.org/10.4193/Rhin.20.515>
 25. Vaira LA, Salzano G, Petrocelli M, Deiana G, Salzano FA, De Riu G (2020) Validation of a self-administered olfactory and gustatory test for the remotely evaluation of COVID-19 patients in home quarantine. *Head Neck* 42:1570–1576
 26. Cazzolla AP, Lovero R, Lo Muzio L, Testa NF, Schirinzi A, Palmieri G et al (2020) Taste and smell disorders in COVID-19 patients: role of interleukin-6. *ACS Chem Neurosci*. <https://doi.org/10.1021/acschemneuro.0c00447>
 27. Mazzatenta A, Neri G, Dardes D, De Luca C, Marinari S, Porreca E et al (2020) Smell and taste in severe COVID-19: self-reported vs. testing. *Front Med Lausanne* 7:589409
 28. Hannum ME, Ramirez VA, Lipson SJ, Herriman RD, Toskala AK, Lin C et al (2020) Objective sensory testing methods reveal a higher prevalence of olfactory loss in COVID-19-positive patients compared to subjective methods: a systematic review and meta-analysis. *Chem Senses* 45:865–874
 29. Vaira LA, Hopkins C, Petrocelli M, Lechin JR, Chiesa-Estomba CM, Salzano G et al (2020) Smell and taste recovery in coronavirus disease 2019 patients: a 60-day objective and prospective study. *J Laryngol Otol* 134:703–709
 30. Boscolo-Rizzo P, Menegaldo A, Fabbris C, Spinato G, Borsetto D, Vaira LA et al (2021) Six-month psychophysical evaluation of olfactory dysfunction in patients with COVID-19. *Chem Senses*. <https://doi.org/10.1093/chemse/bjab006>
 31. Kirschenbaum D, Imbach LL, Ulrich S et al (2020) Inflammatory olfactory neuropathy in two patients with COVID-19. *Lancet*. [https://doi.org/10.1016/S0140-6736\(20\)31525-7](https://doi.org/10.1016/S0140-6736(20)31525-7)
 32. Vaira LA, Hopkins C, Sandison A et al (2020) Olfactory epithelium histopathological findings in long-term COVID-19 related anosmia. *J Laryngol Otol* 134:1123–1127
 33. Vaira LA, Salzano G, Fois AF, Piombino P, De Riu G (2020) Potential pathogenesis of ageusia and anosmia in COVID-19 patients. *Int Forum Allergy Rhinol* 10:1103–1104
 34. Juttler E, Tarabin V, Schwaninger M (2002) Interleukin-6 (IL-6): a possible neuromodulator induced by neuronal activity. *Neuroscientist* 8:268–275
 35. Galiano M, Liu ZQ, Kalla R et al (2001) Interleukin-6 (IL6) and cellular response to facial nerve injury: effects on lymphocyte recruitment, early microglial activation and axonal outgrowth in IL6-deficient mice. *Eur J Neurosci* 14:327–341
 36. Rosleler S, Holtappels G, Wagenmann M, Bachert C (1995) Elevated levels of interleukins IL-1 beta, IL-6 and IL-8 in naturally acquired viral rhinitis. *Eur Arch Otorhinolaryngol* 252:561–563
 37. Bilinska K, Butowt R (2020) Anosmia in COVID-19: a bumpy road to establishing a cellular mechanism. *ACS Chem Neurosci* 11:2152–2155
 38. Ragab D, Salah Eldin H, Taimah M, Khattab R, Salem R (2020) The COVID-19 cytokine storm; what we know so far. *Nat Rev Neurol* 11:1446. <https://doi.org/10.3389/fimmu.2020.01446>
 39. Yimin KM (2006) A regulatory effect of the balance between TNF-alpha and IL-6 in the granulomatous and inflammatory response to *Rhodococcus aurantiacus* infection in mice. *J Immunol* 177:642–650
 40. Torabi A, Mohammadbagheri E, Akbari Dilmaghani N, Bayat AH, Fathi M, Vakili K et al (2020) Proinflammatory cytokines in the olfactory mucosa result in COVID-19 induced anosmia. *ACS Chem Neurosci* 11:1909–1913

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Luigi Angelo Vaira^{1,2}  · Andrea De Vito³ · Giovanna Deiana^{2,4} · Chiara Pes^{5,6} · Federica Giovanditto^{1,7} · Vito Fiore³ · Jerome R. Lechien^{8,9} · Serge-Daniel Le Bon¹⁰ · Sven Saussez^{8,9} · Giordano Madeddu³ · Sergio Babudieri³ · Antonio Pazzola⁷ · Franco Bandiera^{5,6} · Alessandro Giuseppe Fois¹¹ · Andrea Fausto Piana^{2,4} · Claire Hopkins¹² · Giacomo De Riu¹

¹ Maxillofacial Surgery Operative Unit, Department of Medical, Surgical and Experimental Sciences, University of Sassari, Viale San Pietro 43/B, 07100 Sassari, Italy

² Biomedical Science PhD School, Biomedical Science Department, University of Sassari, Sassari, Italy

³ Infectious and Tropical Diseases Unit, Department of Medical, Surgical and Experimental Sciences, University of Sassari, Sassari, Italy

⁴ Clinical Epidemiology and Medical Statistics Unit, Department of Medical, Surgical and Experimental Sciences, University of Sassari, Sassari, Italy

⁵ Internal Medicine Department, University Hospital of Sassari, Sassari, Italy

⁶ Neuro-COVID Department, University Hospital of Sassari, Sassari, Italy

⁷ Onco-COVID Department, University Hospital of Sassari, Sassari, Italy

⁸ COVID-19 Task Force of the Young-Otolaryngologists of the International Federation of Oto-Rhino-Laryngological Societies (YO-IFOS), Paris, France

⁹ Department of Human and Experimental Oncology, Faculty of Medicine UMONS Research Institute for Health Sciences and Technology, University of Mons (UMons), Mons, Belgium

¹⁰ Department of Otorhinolaryngology, CHU Saint-Pierre, Brussels, Belgium

¹¹ Respiratory Diseases Operative Unit, University Hospital of Sassari, Sassari, Italy

¹² King's College, London, UK