



Correction to: Tinnitus, sound intolerance, and mental health: the role of long-term occupational noise exposure

Zahra Jafari¹ · Thomas Coppins² · Glenn Hole² · Femi Nyatepe-Coo² · Bryan E. Kolb¹ · Majid H. Mohajerani¹

Published online: 9 May 2022


© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2022

**Correction to: European Archives of
Oto-Rhino-Laryngology**
<https://doi.org/10.1007/s00405-022-07362-2>

The original article has been corrected.

In the original publication of the article, two typos were identified in the labels of Figs. 1, 2. The correct figures are provided in this correction.

Fig. 1 Average of hearing thresholds in **A** Tin and Tin-CNE groups and **B** cases with SNHL or a normal audiogram. CNE, chronic noise exposure; SNHL, sensory-neural hearing loss



The original article can be found online at <https://doi.org/10.1007/s00405-022-07362-2>.

✉ Zahra Jafari
Zahra.jafari@uleth.ca

¹ Department of Neuroscience, Canadian Centre for Behavioural Neuroscience (CCBN), University of Lethbridge, Lethbridge, AB T1K 3M4, Canada

² Audiology First, Lethbridge, AB T1J 4B5, Canada




Fig. 2 Percentage of participants in different grades of THI, anxiety (HADS-A), depression (HADS-D), and insomnia (ISI) questionnaires. HADS-A/D, hospital anxiety and depression scale-anxiety/depression; ISI, insomnia severity index; THI, tinnitus handicap inventory

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.