



Correction to: Depressive residual symptoms are associated with illness course characteristics in a sample of outpatients with bipolar disorder

Gianluca Serafini¹ · Gustavo H. Vazquez^{2,3,4} · Xenia Gonda^{5,6,7} · Maurizio Pompili⁸ · Zoltan Rihmer⁵ · Mario Amore¹

Published online: 6 March 2018

© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to:

European Archives of Psychiatry and Clinical Neuroscience

<https://doi.org/10.1007/s00406-018-0875-5>

Please, 6–7 lines before the bottom (abstract section), delete the following statement:

“Subjects with residual symptoms were less likely to report substance abuse ($\chi^2_{(2)} = 11.937, p \leq 0.005$) and lifetime psychotic symptoms ($\chi^2_{(2)} = 10.577, p = 0.005$)”

and insert:

“Subjects with residual symptoms were less likely to report substance abuse ($\chi^2_{(2)} = 11.937, p \leq 0.005$), are more likely to report lifetime psychotic symptoms ($\chi^2_{(2)} = 10.577, p = 0.005$)”

Please, 3–4 lines before the bottom (abstract section), delete the following statement:

“After multivariate analyses, a significant positive contribution to residual symptoms was given only by duration

of current illness episode ($\beta = 0.003; p \leq 0.05$), and lifetime psychotic symptoms ($\beta = 1.094; p \leq 0.005$)”.

and insert:

“After multivariate analyses, a significant positive contribution to residual symptoms was given only by duration of current illness episode ($\beta = 0.003; p \leq 0.005$), and significant distressing life events in the last 6 months ($\beta = 1.094; p \leq 0.005$)”.

Page 4, column 2, 10 lines from the top (Results section), delete the following statement:

“likely to report substance abuse (30.3 vs. 69.7%)”

and insert:

“likely to report substance abuse (46.7 vs. 53.3%)”

Finally, throughout the Discussion section, page 10, column 1, 14 lines before the bottom, delete the following statement:

“the functional outcome in BD appear to reflect”.

and insert:

“the functional outcome in BD appears to reflect”.

The original article can be found online at <https://doi.org/10.1007/s00406-018-0875-5>.

✉ Gianluca Serafini
gianluca.serafini@unige.it

¹ Department of Neuroscience, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health (DINO GMI), Section of Psychiatry, University of Genoa, IRCCS Ospedale Policlinico San Martino, Largo Rosanna Benzi 10, 16132 Genoa, Italy

² International Consortium for Bipolar and Psychotic Disorder Research, McLean Hospital, Belmont, MA, USA

³ Department of Neuroscience, Palermo University, Buenos Aires, Argentina

⁴ Department of Psychiatry, Queens University, Kingston, Ontario, Canada

⁵ Department of Psychiatry and Psychotherapy, Kutvolgyi Clinical Center, Semmelweis University, Budapest, Hungary

⁶ MTA-SE Neuropsychopharmacology Research Group, Hungarian Academy of Sciences and Semmelweis University, Budapest, Hungary

⁷ NAP-A-SE New Antidepressant Target Research Group, Semmelweis University, Budapest, Hungary

⁸ Department of Neurosciences, Suicide Prevention Center, Sant'Andrea Hospital, University of Rome, Rome, Italy