

L. Skuterud · Ø. Pedersen · H. Staaland · K. H. Røed ·
B. Salbu · A. Liken · K. Hove


Absorption, retention and tissue distribution of radiocaesium in reindeer: effects of diet and radiocaesium source

Published online: 25 November 2004
© Springer 2004

Radiat Environ Biophys (2004) DOI 10.1007/s00411-004-0257-4

The scale of the vertical axis in Fig. 5 is too low by a factor of 100. The correct scale is given here.

Fig. 5 Activity concentrations of ^{134}Cs in reindeer tissues at the end of experiment 2 normalised to the ^{134}Cs administration rate (mean \pm SD of all 8 animals). RBC, plasma and red bone marrow concentrations are given on the right axis. Neck muscle samples were taken dorsal to the spine at the level of the vertebra 3–5



The online version of the original article can be found at <http://dx.doi.org/10.1007/s00411-004-0257-4>

L. Skuterud (✉)
Norwegian Radiation Protection Authority,
P.O.Box 55, 1332 Østerås, Norway
e-mail: Lavrans.Skuterud@nrpa.no
Tel.: +47-72-485174
Fax: +47-72-486572

Ø. Pedersen · A. Liken · K. Hove
Department of Animal Science,
Agricultural University of Norway,
1432 Ås, Norway

H. Staaland
Department of Biology and Nature Conservation,
Agricultural University of Norway,
1432 Ås, Norway

K. H. Røed
Department of Basic Sciences and Aquatic Medicine,
Norwegian College of Veterinary Medicine,
0033 Oslo, Norway

B. Salbu
Laboratory for Analytical Chemistry,
Agricultural University of Norway,
1432 Ås, Norway