



Parasitic and predatory behavior of *Alysia manducator* (Hymenoptera: Braconidae) on blow fly larvae feeding on an adult pig carcass in the Western Cape Province of South Africa: preliminary observations and forensic implications

Adeyemi Daniel Adetimehin¹ · Calvin Gerald Mole¹ · Devin Alexander Finaughty^{2,3} · Marise Heyns^{1,4}

Received: 31 October 2022 / Accepted: 11 April 2023 / Published online: 21 April 2023
© The Author(s) 2023

Abstract

Wasps are part of the entomofauna associated with vertebrate carrion. They are known to parasitize and prey on specific life stages of insect hosts such as eggs, larvae, pupae, and/or adults associated with vertebrate carrion. However, reports of parasitic behavior of wasps on carrion-associated insect life stages and their possible forensic implications are non-existent in the Western Cape Province of South Africa. This study is part of ongoing research investigating the entomofauna and their pattern of succession on an adult pig carcass in Cape Town, South Africa. During this study, the parasitic wasp *Alysia manducator* was noted parasitizing and preying on blow fly larvae associated with the decomposing carcass. The arrival of *A. manducator* coincides with the occurrence of blow fly eggs and/or larvae on the carcass. These wasps were seen in close association with the eggs and larvae of blow flies on various parts of the carcass and some wasps were seen dragging fly larvae attached to their ovipositors away from one part of the carcass to another. Some *A. manducator* were also observed walking over several larvae on the carcass while exhibiting a stabbing behavior presumably in search of a host for oviposition. We suggest that the observations recorded in this study are of considerable forensic importance as the dragging effect and predatory and stabbing behavior exhibited by *A. manducator* could potentially disrupt the feeding and development of the fly larvae on the carcass. This could subsequently alter the process of carcass decomposition and/or affect minimum post-mortem interval estimations.

Keywords Wasps · *Alysia manducator* · Parasitism · Decomposition · Post-mortem interval · Forensic entomology

Introduction

In the forensic entomology literature, attention has mainly been given to insects belonging to the orders Diptera (flies) and Coleoptera (beetles) [1]. This is ostensibly due to their inherent abilities to locate and utilize vertebrate carrion as suitable substrates for feeding and breeding [1–3]. Due to their necrophagic and reproductive tendencies on a vertebrate carrion, researchers have been able to extract valuable information from these insects such as the minimum time since death, time since human/animal neglect, change in cadaver location, and drugs/poisons presence which could further be applied in medico-legal investigations [1].

Other insect orders have, historically, been overlooked. The forensic importance of the order Hymenoptera has recently begun to attract the attention of forensic entomologists. This order comprises insects such as ants which have

✉ Adeyemi Daniel Adetimehin
yemiadetimehin@gmail.com

¹ Department of Pathology, Division of Forensic Medicine and Toxicology, University of Cape Town, Cape Town, South Africa

² Department of Human Biology, Division of Clinical Anatomy and Biological Anthropology, University of Cape Town, Cape Town, South Africa

³ School of Chemistry and Forensic Science, University of Kent, Canterbury, UK

⁴ School of Medicine, Faculty of Life and Health Sciences, Ulster University, Derry/Londonderry, UK

been documented to not only prey on immature and adult stages of other carrion-associated insects, but also inflict injuries, cause hemorrhage, and prevent fly landing and oviposition on vertebrate remains in various parts of the world [4, 5]. Another interesting insect group within this order is that of the wasps. These insects are frequent responders to decomposing vertebrate remains. Among them is a special group known as parasitoids. These parasitic wasps are necrophilous in nature; however, they do not feed on vertebrate carrion [6]. Instead, they make use of the immature (e.g., eggs, larvae, prepupae, and pupae) stages of blow flies (Diptera: Calliphoridae), flesh flies (Diptera: Sarcophagidae), house flies (Diptera: Muscidae), scuttle flies (Diptera: Phoridae), cheese skippers (Diptera: Piophilidae), and rove beetles (Coleoptera: Staphylinidae) as suitable hosts for the breeding and development of their offspring [6–11].

The parasitic and predatory behavior of wasps have been documented to have substantial implications in biological control programs involving their importation, mass breeding, and deployment in the control and management of invasive and native insect pests and plants in agricultural settings, aquatic environments, urban areas, and animal rearing facilities [12–17]. However, their context in the perspective of forensic entomology has not received the same level of attention. The parasitic and predatory behavior of wasps has been documented to alter the development, morphological appearance, and behavior of the host insect, prevent fly oviposition, prey on adult and immature insect stages, and/or deter the activities of vertebrate scavengers on or around vertebrate carrion [12, 18–23]. This, in turn, may interrupt insect colonization and therefore possibly, the decomposition of the vertebrate carrion. In addition, it can jeopardize the attempt to raise collected immature insect specimens (serving as their hosts) to adults for accurate species identification and/or their application in minimum time since death estimation during forensic investigations [6]. Equally important, wasps can cause post-mortem lacerations and injuries on vertebrate carrion which can potentially create entry sites for other insects and possibly mislead forensic investigators during cause-of-death investigations [21, 24].

Several wasp families have been reported to contain parasitic and predatory species. This includes the families Braconidae, Ichneumonidae, Pteromalidae, Figitidae, Vespidae, Eulophidae, Chalcididae, Diapriidae, Encyrtidae, and Proctotrupidae [11, 20, 24–27]. Depending on the species, parasitic wasps possess the ability to forage and parasitize specific life stages of selected insect hosts and, thus, can serve as alternative forensic indicator species after the consideration of fly evidence in the estimation of the minimum time since death, verification of cadaver relocation, and location of hidden vertebrate remains [6, 19, 28]. The limited availability of published reports on the activities of parasitoid wasps on vertebrate remains alongside their application

in forensic entomological investigations might be linked to them being neglected or unseen during forensic entomological investigations due to their small size, tendency to arrive during the mid- to late stages of decomposition, and the paucity of published data on their biology and ecology [6, 26]. Nevertheless, there exist reports on the potential value of parasitic wasps in forensic investigations and minimum time since death estimations [9, 10, 28–30].

In South Africa, parasitoid wasps such as *Nasonia vitripennis* (Hymenoptera: Pteromalidae) and *Trichopria lewisii* (Hymenoptera: Diapriidae) have been found to parasitize the pupae of the forensically important fly, *Chrysomya albiceps* (Diptera: Calliphoridae) in the Kruger National Park of the Limpopo/Mpumalanga Provinces [8]. However, we are not aware of any record of the activities of parasitoid wasps on a vertebrate carrion and carrion-associated immature insects in the Western Cape Province—a locale that is markedly different from a biogeographic perspective to that of the Limpopo and Mpumalanga Provinces across which the Kruger National Park spans. Thus, we report preliminary observations and possible forensic implications of the parasitic and predatory behavior of *A. manducator* (Hymenoptera: Braconidae) on the immature stages (i.e., eggs and larvae) of blow flies associated with a decomposing adult pig carcass in the winter season of the Western Cape Province.

Materials and methods

Study site

This study is part of an ongoing winter carrion decomposition and insect successional study within the area of Table Mountain National Park, adjacent to the University of Cape Town Upper Campus, Rondebosch, South Africa (S33°57.682'; E018°27.301'). The Table Mountain National Park is characterized by a rocky terrain situated on the Cape Peninsula, a mountainous region located at the south-western tip of the African continent [5, 31]. The Park is dominated by Mediterranean-like shrubland called fynbos—part of the Cape Floristic Kingdom—which is prone to fire [5, 32]. The park harbors 2285 plant species of which 90 are endemic [5, 32]. Pockets of invasive alien vegetation exist, dominated by *Acacia*, *Pinus*, and *Hakea* species [5, 32].

Decomposition study and animal model

As part of this study, a 60-kg adult pig (*Sus scrofa domestica* L.) carcass has been used as an animal model. Pigs have been regarded as the most suitable analogues to a human body where the establishment of baseline data is concerned due to the similarities in the integumentary, circulatory, and digestive systems [33]. The adult pig was purchased from the

University of Stellenbosch, Faculty of AgriSciences Piggery Unit after which it was humanely terminated by a single shot to the base of the brain with a 0.22 caliber long rifle by the farm manager on the day of the winter solstice (21 June 2022) of the Southern Hemisphere [34]. Immediately following termination, the adult pig carcass was rinsed and sealed within a body bag for transport to the research site. At the research site, the carcass was deployed in a steel cage covered with chicken wire (Fig. 1). Thereafter, decompositional changes and insect activities on the carcass were recorded daily.

Documentation of observations, collection, and identification of wasps

The location(s) of the wasps on, in, and around the carcass during data collection were documented using detailed field notes and manual photography. The characteristic activities of the wasps on, in, and around the carcass and their interaction with other insect life stages (e.g., fly eggs and larvae) present on and around the carcass were documented with field notes, manual photography, and video recordings. The wasps were collected from the carcass with the aid of soft tweezers after undisturbed video recordings and photographs were taken. After collection, the wasps were transferred immediately into a killing jar containing paper towels dampened with ethyl acetate [5]. Following termination, the wasps were transferred into screwcap containers after which they were taken to the Forensic Entomology Laboratory, Faculty of Health Sciences, University of Cape Town, placed in 70% ethanol, and then stored in a refrigerator until identification.

Results and discussion

The wasp specimens were identified to be *Alysia manducator* Panzer (Hymenoptera: Braconidae) by Dr. Simon van Noort (Iziko Museums of South Africa). Images of this species and further information is available on WaspWeb [35]. We observed in this study the predatory and parasitic behavior of *A. manducator* on blow fly larvae feeding on animal carrion. The arrival of *A. manducator* individuals coincides with the occurrence of blow fly eggs and larvae on the carcass. We found several individuals of *A. manducator* in close association with the eggs and larvae of blow flies on various parts of the carcass. This observation is consistent with the findings of Horenstein and Salvo [36] in central Argentina and that of our previous studies using neonate pigs in all the seasons of the Western Cape (Ade-timehin et al., unpublished data). According to Reznik et al. [19], meat samples infested with *Calliphora vicina* (Diptera: Calliphoridae) larvae were significantly attractive to *A. manducator* when compared to meat samples in similar decomposition stage but without *Ca. vicina* larvae. Similarly, *A. manducator* has been reported to parasitize the larvae of *Lucilia sericata* (Diptera: Calliphoridae) feeding on vertebrate carrion in England, UK [37].

The ability of parasitic wasps (including *A. manducator*) to locate the immature stages of suitable hosts and hosts' source of nutrition (e.g., vertebrate carrion) might be linked to their sensitivity to the chemical and visual cues of the host insect and its food source [7, 12, 19]. In fact, Heo et al. [38] reported that the parasitoid wasps namely *Exoristobia philippinensis* (Hymenoptera:

Fig. 1 Close-up view of the adult pig carcass within the steel cage



Encyrtidae) and *Dirhinus himalayanus* (Hymenoptera: Chalcididae) were able to locate and parasitize the pupae of *Liopygia ruficornis* (Diptera: Sarcophagidae) in a baited trap positioned on the rooftop of a high-rise building approximately 101 m above the ground. The forensic application of this is that the foraging behavior of wasps and their ability to detect the chemical cues of their host insect and its food source can serve as useful alternative pointers to the presence of decomposing vertebrate remains and other carrion-associated immature insect life stages (e.g., larvae) in concealed environments (e.g., in the bushveld) and multistorey buildings. Advances in biosensor research have demonstrated the use of wasp species as biological sensors in the identification or detection of specific visual, chemical, and odor cues. Particularly, this has led to the development of the “wasp hound”, a device originally used to detect volatile fungal chemicals [39]. Further research has indicated the ability of conditioning wasp species to detect certain odors or chemical cues associated with animal food products [40], explosives [41, 42], fungal pathogens [43], illicit drugs [44], and vertebrate carrion [45]. Based on the current observations, we suggest that further studies within the region should utilize the “wasp hound” in the training and conditioning of adult *A. manducator* individuals specifically in the detection of the volatile organic chemicals (VOCs) emanating from vertebrate carrion, alongside carrion-associated insects, and the vertebrate carrion itself in different depositional environments (e.g., above ground, hanged, buried, vehicular confinement, trashcan, suitcase, and drum).

Parasitic behavior: stabbing of fly larvae

We observed some individuals of *A. manducator* walking over several larvae on the carcass and then exhibiting a stabbing behavior (Online Resources 1 and 2) presumably in search of a location or host for the purpose of egg laying [7, 11, 19, 46, 47]. This was subsequently confirmed by the presence of a fly larva attached to the ovipositor of one of the collected wasp individuals (Fig. 2). We attribute the commencement of the stabbing behavior by *A. manducator* to the concentration of the decay odor emanating from the carcass and the movement of the fly larvae underneath its legs [7, 19]. Parasitic wasps including *A. manducator* have been reported to exhibit similar behavior on vertebrate carrion [7, 11, 19]. After the stabbing of the host insect, the cuticular teeth on the ovipositor of the parasitoid wasp pierces and ruptures the host insect’s skin [46, 47]. Furthermore, the stabbing behavior of parasitic wasps (including *A. manducator*) brings about the paralysis of the host insect larvae [19, 47]. Owing to this, we speculate that the observed



Fig. 2 A blow fly larva attached to the ovipositor (yellow arrow) of a collected *Alysia manducator* individual

stabbing of the blow fly larvae by *A. manducator* could potentially cause injuries/damages to the soft tissues of the blow fly larvae.

Furthermore, the stabbing behavior of parasitic wasps is a mechanism associated with oviposition [19], and after oviposition/egg eclosion, the likely consumption of the tissues of the immature host insect (e.g., pupa) by the larval stage of the parasitoid wasp could lead to the death of the insect host [18]. In addition, the subsequent development and emergence of the adult parasitoids from their host insects could compromise the morphology and rearing process of collected immature insect specimens (e.g., larvae/pupae) to adults for species confirmation during crime scene investigations. A notable study in this regard was that of Heo et al. [20]. The authors reported the presence of two exit holes on the pupal casing of *Chrysomya rufifacies* (Diptera: Calliphoridae) following the emergence of the adults of the parasitoid wasp, *E. philippinensis*. Equally important, the stabbing behavior by wasps could alter the physiology (e.g., diapause) and development of the immature insect hosts, and manipulate their biology (e.g., feeding) and behavior (e.g., movement) [12, 18, 48]. For example, Cammack et al. [12] reported that the presence of *N. vitripennis* females led to an increase in the developmental rate of *L. sericata* pupae. Similarly, parasitism by *A. manducator* triggered premature pupation in *L. sericata* [48]. Furthermore, Cammack et al. [12] revealed that *L. sericata* larvae burrowed into the soil to escape parasitism by *N. vitripennis*. The resulting impacts of the stabbing behavior by *A. manducator* are substantial and from the forensic entomological point of view, it can potentially influence the estimation of the age of immature fly specimens and their utilization in minimum time since death estimations. Thus, we suggest that forensic entomologists consider the presence/absence of wasps in, on, or around vertebrate remains prior to and after the collection of immature entomological evidence before their utilization in minimum time since death estimations.

Predation and dragging of fly larvae

We found one *A. manducator* individual close to the oral cavity preying on a blow fly larva (Online Resource 3). Some individuals of *A. manducator* were also seen dragging away blow fly larvae attached to their ovipositors (Fig. 3; Online Resources 4 and 5). The ability of *A. manducator* to drag or pull away the immature stages of flies from or toward the carcass or from one body part to another can potentially alter the process of decomposition. For instance, the dragging of a fly larva that has successfully established an entry point or close to establishing one on the skin of a carcass to another region of the body that has not been impacted, can potentially increase the feeding time the specific larva has to go through to gain access to the soft tissues and internal organs of the body.

The persistence of the ovipositor and the continuous dragging of the blow fly larvae by *A. manducator* could potentially overdose the host insect with more toxins and may bring about the premature death of the host larvae [47]. Also, it should be pointed out that the persistence of the wasp's ovipositor in the blow fly larva and the resultant dragging could be as a result of the level of experience of the female parasitoid wasp in egg laying [47]. In the literature, some studies have documented how parasitoid wasps drag or pull away their host insects with the aid of their ovipositors and mandibles [49, 50]. While the structure of the ovipositors in parasitic Hymenoptera are similar, parasitoid wasps such *Leptopinila heterotoma* (Diptera: Hymenoptera) have been reported to possess a structure called the “ovipositor clip” in their ovipositors which play a key role in holding their hosts [47].

Present but exhibiting no observable parasitic and/or predatory behavior

Interestingly, for some *A. manducator* individuals, we did not observe them exhibiting any stabbing and/or predatory



Fig. 3 An *Alysia manducator* individual on the soil surface dragging a blow fly larva attached to its ovipositor (yellow arrow)

behavior on or toward the blow fly larvae. Although *A. manducator* individuals are known to be parasitoids of the larvae of flies associated with vertebrate remains [7], their behavioral variation as observed in this study might be due to the age and size of the fly larvae, maturity status and oviposition experience of the female wasp, parasitism status of the fly larvae, and/or odor trail left behind on/around the site of oviposition by the host parent [7, 19, 47, 51]. According to Reznik et al. [19], the increase in probing/contact ratio of *A. manducator* on *Ca. vicina* larvae, for example, may be attributed to the increase in size of the larvae within the 5-day developmental period. However, the possible increase in strength of the larvae due to the size increase may have contributed to the reduction in the oviposition and attack of *A. manducator* on *Ca. vicina* larvae (Reznik et al. [19]). Furthermore, the authors revealed that the decrease in the probing behavior of *A. manducator* on mature *Ca. vicina* larvae could be as a result of the reduction in the smell of the larvae as evidenced by their empty intestinal contents.

Future research

The current study presents observations on the predatory and parasitic behavior of the wasp *A. manducator*, highlighting the need for further research to ascertain the degree of effect such species may have on the estimation of the minimum time since death. While previous research has demonstrated developmental, morphological, physiological, and behavioral changes in host species of parasitoid wasps [12, 18–20, 48], further laboratory-based research is needed to fully understand the medico-legal implications of such behavior. The use of wasp species in the detection of human remains is a further avenue of interest, as advances in the literature have demonstrated the effectiveness of a device called the “wasp hound” in the training and conditioning of wasp species as biological sensors in the identification or detection of specific visual, chemical, and odor cues associated with food products, explosives, illicit drugs, and vertebrate carrion [40–45].

Conclusion

The current study presents observations on the predatory and parasitic behavior of the wasp *A. manducator* associated with a decomposing adult pig carcass. While much of the observations have previously been reported in ecological and entomological literature, the current study highlights the importance of these observations for forensic entomology.

We observed *A. manducator* stabbing the larvae of blow flies on multiple occasions. Also, we noted the dragging of some larvae away and/or toward the carcass. We speculate

that the (1) stabbing behavior of *A. manducator* can cause injuries to the soft tissues of the larvae which can alter the development and behavior of blow fly larvae and their utilization in minimum time since death estimations; (2) dragging of blow fly larvae away from the carcass or from one part of the carcass to another can influence the rate of decomposition of a vertebrate carrion particularly in the early stages where minimal larvae are present; (3) likely oviposition of the wasps and the development of their offspring on/in the larvae can potentially inhibit the development of the collected immature insects to adults for accurate species identification; and (4) ability of wasps to locate their insect host and its associated food source can provide useful indications as to the location of cadavers and cadaver-associated immature insects in concealed environments.

These observations highlight the need for further research to ascertain the degree of effect such species may have on the estimation of the minimum time since death. Particularly, there is a need for more laboratory and field studies on the biology, ecology, and foraging behavior of *A. manducator* on food resources including vertebrate carcasses in the presence or absence of the immature stages (e.g., eggs, larvae, and pupae) of similar/multiple forensically important fly species. This may help clarify if *A. manducator* exhibit any host specificity in relation to its parasitic behavior toward larvae of similar/multiple fly species and how such behavior affects the larvae's feeding and development. We also suggest the need for laboratory and field studies focused on examining the effectiveness of trained/conditioned adult *A. manducator* individuals in the detection of vertebrate carrion and carrion-associated chemical cues in our region. Lastly, as with other similarly cited articles from different parts of the world, we urge forensic entomologists to take cognizance of the presence or absence of wasps in/around the crime/death scene during entomological evidence collection as this may have an influence on the development of the insect evidence prior to or after collection and their utilization in minimum post-mortem interval estimations.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00414-023-03001-5>.

Acknowledgements The authors thank the Authorities of the South African National Parks and Table Mountain National Park for hosting the research project within their premises, Mr. John Morris (Farm Manager: Mariendahl Experimental Farm, Stellenbosch University) for providing the adult pig used for this research, and Dr. Simon van Noort of the Iziko Museums of South Africa for the identification of the collected wasp specimens. The authors also thank the South African National Research Foundation and University of Cape Town for providing A.D.A. with the Grantholder Student-Linked Bursary, International Students' Scholarship, JW Jagger Centenary Gift Scholarship, and Postgraduate Research Training Grant for his academic program. The financial assistance of the National Research Foundation towards this research is hereby acknowledged. Opinions expressed and conclusions arrived at are those of the authors and are not necessarily to be attributed to the National Research Foundation.

Author contribution A.D.A.: conceptualization, methodology, writing—original draft, writing—review and editing, visualization, and investigation. M.H.: conceptualization, methodology, resources and funding acquisition, writing—review and editing, supervision, project administration. D.A.F.: conceptualization, methodology, writing—review and editing, and supervision. C.G.M.: writing—review and editing, supervision, and project administration.

Funding Open access funding provided by University of Cape Town. This study was funded by the South African National Research Foundation (NRF) through a Research Grant awarded to M.H. (grant number: CSUR116299).

Data availability All data generated or analyzed in relation to this study are included in this published article and its supplementary information files.

Declarations

Ethics approval Ethical clearance for the use of the adult pig and the entire study was obtained from the University of Cape Town, Faculty of Health Sciences Animal Ethics Committee (UCT FHS AEC Reference Number: 021_021).

Conflict of interest The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Byrd JH, Castner JL (2010) Forensic entomology the utility of arthropods in legal investigations, 2nd edn. CRC Press, Boca Raton, London, New York
2. Greenberg B (1991) Flies as forensic investigators. *J Med Entomol* 28(5):565–577
3. Matuszewski S, Szafałowicz M (2013) Temperature-dependent appearance of forensically useful beetles on carcasses. *Forensic Sci Int* 229:92–99
4. Eubanks MD, Lin C, Tarone AM (2019) The role of ants in vertebrate carrion decomposition. *Food Webs* 18:e00109
5. Adetimehin AD, Mole CG, Finaughty DA, Heyns M (2022) Caught in the act: impact of *Crematogaster* cf. *liengmei* (Hymenoptera: Formicidae) necrophagous behavior on neonate pigs (*Sus scrofa domestica* L.) in the Western Cape Province of South Africa. *Int J Legal Med*. <https://doi.org/10.1007/s00414-022-02835-9>
6. Rivers DB (2016) Parasitic Hymenoptera as forensic indicator species. In: Shetty B, Padubidri J (eds) *Forensic analysis – from death to justice*. IntechOpen, London, pp 67–83
7. Laing J (1937) Host-finding by insect parasites. 1. Observations on the finding of host by *Alysia manducator*, *Mormoniella vitripennis* and *Trichogramma evanescens*. *J Anim Ecol* 6(2):298–317

8. Braack LEO (1987) Community dynamics of carrion-attendant arthropods in Tropical African Woodland. *Oecologia* 72(3):402–409
9. Grassberger M, Frank C (2003) Temperature-related development of the parasitoid wasp *Nasonia vitripennis* as forensic indicator. *Med Vet Entomol* 17:257–262
10. Disney RHL, Munk T (2004) Potential use of Braconidae (Hymenoptera) in forensic cases. *Med Vet Entomol* 18:442–444
11. Frederickx C, Dekeirsschieter J, Verheggen FJ, Haubruge E (2013) The community of Hymenoptera parasitizing necrophagous Diptera in an urban biotope. *J Insect Sci* 13:32
12. Cammack JA, Adler PH, Tomberlin JK, Arai Y, Bridges JWC (2010) Influence of parasitism and soil compaction on pupation of the green bottle fly, *Lucilia sericata*. *Entomol Exp Appl* 136:134–141
13. Machtinger ET, Geden CJ, Kaufman PE, House AM (2015) Use of pupal parasitoids as biological control agents of filth flies on equine facilities. *J Integr Pest Manag* 6(1):16
14. Kraus EC, Coetzee J, van Noort S, Olmi M (2019) First record of an indigenous South African parasitoid wasp on an imported biological control agent, the water hyacinth hopper. *Biocontrol Sci Tech* 29(12):1234–1241
15. Southon RJ, Fernandes OA, Nascimento FS, Sumner S (2019) Social wasps are effective biocontrol agents of key lepidopteran crop pests. *Proc R Soc B* 286:20191676
16. Zhi-zhi W, Yin-quan L, Min S, Jian-hua H, Xue-xin C (2019) Parasitoid wasps as effective biological control agents. *J Integr Agric* 18(4):705–715
17. van Noort S, Smith R, Coetzee JA (2021) Identity of parasitoid wasps (Hymenoptera, Braconidae and Eulophidae) reared from aquatic leaf-mining flies (Diptera, Ephydriidae) on invasive Brazilian waterweed *Egeria densa* in South Africa. *Afr Invertebr* 62(1):287–314
18. Vinogradova EB, Zinovjeva KB (1972) Experimental investigation of the seasonal aspect of the relationship between blowflies and their parasites. *J Insect Physiol* 18(9):1629–1638
19. Reznik SY, Chernoguz DG, Zinovjeva KB (1992) Host searching, oviposition preferences and optimal synchronization in *Alysia manducator* (Hymenoptera: Braconidae), a parasitoid of the blowfly. *Calliphora vicina* Oikos 65(1):81–88
20. Heo CC, Nazni WA, Lee HL, John J, Baharudin O, Chen CD, Lau KW, Sofian-Azirun M (2009) Predation on pupa of *Chrysomya rufifacies* (Marquart) (Diptera: Calliphoridae) by parasitoid, *Exoristobia philippinensis* Ashmead (Hymenoptera: Encyrtidae) and *Ophyra spinigera* larva (Diptera: Muscidae). *Trop Biomed* 26(3):369–372
21. Moretti TDC, Giannotti E, Thyssen PJ, Solis DR, Godoy WAC (2011) Bait and habitat preferences, and temporal variability of social wasps (Hymenoptera: Vespidae) attracted to vertebrate carrion. *J Med Entomol* 48(5):1069–1075
22. Spencer EE, Barton PS, Ripple WJ, Newsome TM (2020) Invasive European wasps alter scavenging dynamics around carrion. *Food Webs* 24:e00144
23. Cairncross RJ, Barton PS, Bonat S, Crowther MS, Dickman CR, Vandersteen J, Newsome TM (2022) The predatory impacts of invasive European wasps on flies are facilitated by carcasses with open wounds. *Food Webs* 31:e00227
24. Somavilla A, Linard V, Rafael JA (2019) Social wasps (Vespidae: Polistinae) on carcasses of *Rattus norvegicus* (Mammalia: Muridae) in the Central Amazonia, Brazil: possible forensic implications. *Revista Brasileira de Entomologia* 63:18–21
25. Grassberger M, Frank C (2004) Initial study of arthropod succession on pig carrion in a central European urban habitat. *J Med Entomol* 41(3):511–523
26. Voss SC, Spafford H, Dadour IR (2009) Hymenopteran parasitoids of forensic importance: host associations, seasonality, and prevalence of parasitoids of carrion flies in Western Australia. *J Med Entomol* 46(5):1210–1219
27. Kolyada V, Perkovsky E (2011) A new species of the genus *Disogmus* Förster (Hymenoptera, Proctotrupoidea, Proctotrupidae) from the Eocene Rovno amber. *Zookeys* 130:455–459
28. Amendt J, Krettek R, Niess C, Zehner R, Bratzke H (2000) Forensic entomology in Germany. *Forensic Sci Int* 113:309–314
29. Huchet JB, Greenberg B (2010) Flies, Mochicas and burial practices: a case study from Huaca de la Luna, Peru. *J Archaeol Sci* 37:2846–2856
30. Reigada C, Gião JZ, Galindo LA, Godoy WAC (2011) Survival of submerged blowfly species and their parasitoids: implications for post-mortem submersion interval. *Forensic Sci Int* 212:126–129
31. Poulsen ZC, Hoffman MT (2015) Changes in the distribution of indigenous forest in Table Mountain National Park during the 20th Century. *S Afr J Bot* 101:49–56
32. van Wilgen BW, Forsyth GG, Prins P (2012) The management of fire-adapted ecosystems in an urban setting: The case of Table Mountain National Park South Africa. *Ecol Soc* 17(1):8
33. Matuszewski S, Hall MJR, Moreau G, Schoenly KG, Tarone AM, Villet MH (2020) Pigs vs people: the use of pigs as analogues for humans in forensic entomology and taphonomy research. *Int J Legal Med* 134:793–810
34. Kelly JA (2006) The influence of clothing, wrapping and physical trauma on carcass decomposition and arthropod succession in central South Africa. Dissertation, University of the Free State
35. van Noort S (2022) WaspWeb: Hymenoptera of the Afrotropical region. URL: www.waspweb.org. Accessed 19 October 2022
36. Horenstein MB, Salvo A (2012) Community dynamics of carrion flies and their parasitoids in experimental carcasses in central Argentina. *J Insect Sci* 12:8
37. Smith KE, Wall R (1997) The use of carrion as breeding sites by the blowfly *Lucilia sericata* and other Calliphoridae. *Med Vet Entomol* 11(1):38–44
38. Heo CC, Latif B, Kurahashi H, Tan SH, Chew WK, Nazni WA, Omar B (2017) Preliminary observations of necrophagous dipteran activities and a new host record of parasitoids on a high-rise building. *Pol J Entomol* 86:165–180
39. Rains GC, Utley SL, Lewis WJ (2006) Behavioral monitoring of trained insects for chemical detection. *Biotechnol Prog* 22(1):2–8
40. Wäckers F, Olson D, Rains G, Lundby F, Haugen J-E (2011) Boar taint detection using parasitoid biosensors. *J Food Sci* 76(1):41–47
41. Tomberlin JK, Tertuliano M, Rains G, Lewis WJ (2005) Conditioned *Microplitis croceipes* Cresson (Hymenoptera: Braconidae) detect and respond to 2,4-DNT: development of a biological sensor. *J Forensic Sci* 50(5):1187–1190
42. Tomberlin JK, Rains GC, Sanford MR (2008) Development of *Microplitis croceipes* as a biological sensor. *Entomol Exp Appl* 128:249–257
43. Rains GC, Tomberlin JK, D'Alessandro M, Lewis WJ (2004) Limits of volatile chemical detection of a parasitoid wasp, *Microplitis croceipes*, and an electronic nose: a comparative study. *Trans ASAE* 47(6):2145
44. Olsen D, Rains G (2014) Use of a parasitic wasp as a biosensor. *Biosensors* 4:150–160
45. Frederickx C, Verheggen FJ, Brostaux Y, Haubruge E (2014) Associative learning of *Nasonia vitripennis* Walker (Hymenoptera: Pteromalidae) to methyldisulfanylmethane. *J Forensic Sci* 59(2):413–416
46. Swain RB, Green W, Portman R (1938) Notes on oviposition and sex ratio in *Hyposoter pilosulus* Prov. (Hym.: Ichneumonidae). *J Kansas Entomol Soc* 11(1):7–9

47. van Lenteren JC, Isidoro N, Bin F (1998) Functional anatomy of the ovipositor clip in the parasitoid *Leptopilina heterotoma* (Thompson) (Hymenoptera: Eucoilidae) a structure to grip escaping host larvae. *Int J Insect Morphol Embryol* 27(3):263–268
48. Holdaway FG, Evans AC (1930) Parasitism a stimulus to pupation: *Alysia manducator* in relation to the host *Lucilia sericata*. *Nature* 125:598–599
49. Evans HE (1962) Evolution of prey-carrying mechanisms in wasps. *Evolution* 16(4):468–483
50. Rubink WL, Evans HE (1979) Notes on the nesting behavior of the Bethyloid wasp, *Epyris eriogoni* Kieffer, in southern Texas. *Psyche* 86(4):313–319
51. Mukai H, Kitajima H (2021) Laboratory evaluation of *Orthocentrus brachycerus* (Hymenoptera: Ichneumonidae), as a potential biological control agent in mushroom cultivation. *J Appl Entomol* 145:348–357

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.