

## The value of skin biopsy with recording of intraepidermal nerve fiber density and quantitative sensory testing in the assessment of small fiber involvement in patients with different causes of polyneuropathy

Maria Nebuchennykh · Sissel Løseth ·  
Sigurd Lindal · Svein Ivar Mellgren

Published online: 6 June 2009  
© Springer-Verlag 2009

**Erratum to: J Neurol**  
**DOI 10.1007/s00415-009-5065-y**

Unfortunately, one paragraph under the heading “Reference material and normal values” was published with errors. The correct formula is given here.

The Z-score was calculated according to the formula:  
[LnIENF (patient value) – (2.733 – 0.004 \* age – 0.178 \* gender (1 men, 0 women))]/0.343.

---

The online version of the original article can be found under  
doi:[10.1007/s00415-009-5065-y](https://doi.org/10.1007/s00415-009-5065-y).

M. Nebuchennykh (✉) · S. Løseth · S. I. Mellgren  
Department of Neurology, University Hospital of North Norway,  
9038 Tromsø, Norway  
e-mail: maria.nebuchennykh@unn.no

M. Nebuchennykh · S. Løseth · S. I. Mellgren  
Institute of Clinical Medicine, University of Tromsø,  
9038 Tromsø, Norway

S. Lindal  
Department of Pathology, University Hospital of North Norway,  
Tromsø, Norway