Erratum to: Botulinum toxin type A in post-stroke lower limb spasticity: a multicenter, double-blind, placebo-controlled trial

Ryuji Kaji · Yuka Osako · Kazuaki Suyama ·
Toshio Maeda · Yasuyuki Uechi · Masaru Iwasaki

Published online: 8 June 2010
© Springer-Verlag 2010

Erratum to: J Neurol DOI 10.1007/s00415-010-5526-3

There was an error in the eligibility criteria in “Methods” section. Instead of:

“Patients

Male or female patients aged 20–80 years and weighing at least 50 kg were eligible if they had had a stroke at least 6 months prior to treatment and had equinus deformity (plantar flexion of the ankle) as demonstrated by a score

of **>3** for ankle flexors on the Modified Ashworth Scale (MAS).”

It should read:

“Patients

Male or female patients aged 20–80 years and weighing at least 50 kg were eligible if they had had a stroke at least 6 months prior to treatment and had equinus deformity (plantar flexion of the ankle) as demonstrated by a score of **3 or 4** for ankle flexors on the Modified Ashworth Scale (MAS).”

On behalf of GSK1358820 Spasticity Study Group.

The online version of the original article can be found under doi:[10.1007/s00415-010-5526-3](https://doi.org/10.1007/s00415-010-5526-3).

R. Kaji (✉)
Department of Clinical Neuroscience, Institute of Health
Biosciences, Tokushima University Graduate School,
2-50-1 Kuramotocho, Tokushima, Tokushima 770-8503, Japan
e-mail: rkaji@clin.med.tokushima-u.ac.jp

Y. Osako · K. Suyama · T. Maeda · Y. Uechi · M. Iwasaki
Japan Development and Medical Affairs Division,
GlaxoSmithKline K.K., 4-6-15 Sendagaya, Shibuya-ku,
Tokyo 151-8566, Japan