Erratum to: Effects of acute relapses on neuropsychological status in multiple sclerosis patients

S. A. Morrow · S. Jurgensen · F. Forrestal ·
Frederick E. Munchauer · R. H. B. Benedict

Published online: 2 April 2011
© Springer-Verlag 2011

Erratum to: J Neurol DOI 10.1007/s00415-011-5975-3

Table 1 was published with errors. Correct table is given here:

Table 1 Pre-relapse demographics and disability measures

	Cases	Controls	<i>p</i> value
Age (years)			
Mean ± SD	44.0 ± 7.1	42.3 ± 7.2	0.387*
Gender			
<i>N</i> (%) female	35 (66.0%)	75 (65.2%)	0.917 ⁺
EDSS prior to relapse			
Median (range)	3.0 (1.0–7.5)	2.5 (0.0–7.0)	0.717 ⁺
Cerebral FS prior to relapse			
Median (range)	0.0 (0.0–2.0)	0.0 (0.0–2.0)	0.777 ⁺
SDMT 2 visits prior to relapse			
Mean ± SD	52.3 ± 11.6	52.5 ± 11.3	0.937*
SDMT 1 visit prior to relapse			
Mean ± SD	53.6 ± 10.9	53.4 ± 11.6	0.658*
MSNQ 1 visit prior to relapse			
Mean ± SD	15.0 ± 11.2	13.0 ± 10.8	0.112*
MSNQ 2 visits prior to relapse			
Mean ± SD	14.1 ± 10.6	13.3 ± 11.0	0.666*
Corticosteroid treatment			
<i>N</i> (%)	37 (56.9%)	N/A	

SD standard deviation; *EDSS* expanded disability status scale; *FS* function scale; *SDMT* Symbol Digit Modalities Test; *MSNQ* Multiple Sclerosis Neuropsychological Screening Questionnaire

* ANOVA, ⁺ Chi-square

The online version of the original article can be found under doi:[10.1007/s00415-011-5975-3](https://doi.org/10.1007/s00415-011-5975-3).

S. A. Morrow · R. H. B. Benedict
Jacobs Neurological Institute,
State University of New York at Buffalo,
Buffalo, NY, USA

S. Jurgensen · F. Forrestal · F. E. Munchauer
Biogen Idec, Inc., Cambridge, MA, USA

R. H. B. Benedict (✉)
Buffalo General Hospital, 100 High Street,
Buffalo, NY 14203, USA
e-mail: benedict@buffalo.edu