



Correction to: Effectiveness of Rotigotine plus intensive and goal-based rehabilitation versus Rotigotine alone in “de-novo” Parkinsonian subjects: a randomized controlled trial with 18-month follow-up

D. Ferrazzoli¹ · P. Orтели¹ · G. Riboldazzi^{2,3} · R. Maestri⁴ · G. Frazzitta¹

Published online: 17 August 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to: Journal of Neurology (2018) 265:906–916
<https://doi.org/10.1007/s00415-018-8792-0>

The original version of this article unfortunately contained a mistake. The funding information was incorrect. The corrected funding information is given below.

This study was funded by UCB Biopharma—Investigator-Initiated Study (UCB IIS-2014-101555).

The original article can be found online at <https://doi.org/10.1007/s00415-018-8792-0>.

✉ D. Ferrazzoli
davideferrazzoli@gmail.com

¹ Department of Parkinson’s Disease, Movement Disorders and Brain Injury Rehabilitation, “Moriggia-Pelascini” Hospital-Via Pelascini 3, Gravedona ed Uniti (CO), 22015 Como, Italy

² Parkinson’s Disease and Movement Disorders Center, ASST dei Sette Laghi, Varese, Italy

³ Parkinson’s Disease Rehabilitation Center, Fondazione Borghi, Brebbia, Varese, Italy

⁴ Department of Biomedical Engineering, Istituti Clinici Scientifici Maugeri Spa Società Benefit, IRCCS, Via per Montescano 3, Montescano, 27040 Pavia, Italy