

Kaan Gündüz
Carol L. Shields
Ilhan Günalp
Esra Erden
Jerry A. Shields

Orbital schwannoma: correlation of magnetic resonance imaging and pathologic findings

Published online: 10 February 2004
© Springer-Verlag 2004

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s00417-003-0681-1>

K. Gündüz (✉)
G.M.K. Bulvarı 116/3,
06570 Maltepe, Turkey
e-mail: eyemd@ada.net.tr
Fax: +90-312-2291812

K. Gündüz · I. Günalp
Oncology Service,
Department of Ophthalmology,
Ankara University Faculty of Medicine,
Ankara, Turkey

C. L. Shields · J. A. Shields
Oncology Service, Wills Eye Hospital,
Thomas Jefferson University,
Philadelphia, Pennsylvania, USA

E. Erden
Department of Pathology,
Ankara University Faculty of Medicine,
Ankara, Turkey

Graefe's Arch Clin Exp Ophthalmol (2003)
241:593–597

Materials and Methods section, 2nd paragraph, 1th line:

The first sentence should read “After preoperative MRI was obtained, the patients underwent orbitotomy for tumor removal” instead of “After preoperative MRI was obtained, the patients underwent orbitotomy and the tumors were totally excised.”