

CORRECTION



Correction to: Quantitative analysis of choroidal blood flow parameters in optical coherence tomography and angiography in central serous chorioretinopathy

Bo-Een Hwang^{1,2} · Jae-Hyuck Kwak^{1,2} · Joo-Young Kim^{1,2} · Rae-Young Kim^{1,2} · Mirinae Kim^{1,2} · Young-Geun Park^{1,2} · Young-Hoon Park^{1,2}

Published online: 21 March 2022
© Springer-Verlag GmbH Germany, part of Springer Nature 2022

Correction to: Graefe's Archive for Clinical and Experimental Ophthalmology
<https://doi.org/10.1007/s00417-022-05588-8>

In the published version of this article, Table 3 contained errors.

Table 3 should be presented as below.

This is being corrected in this publication.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00417-022-05588-8>.

✉ Young-Hoon Park
parkyh@catholic.ac.kr

¹ Department of Ophthalmology and Visual Science, Seoul St. Mary's Hospital, The Catholic University of Korea School of Medicine, 222 Banpo-daero, Seocho-gu, Seoul 06591, Republic of Korea

² Catholic Institute for Visual Science, The Catholic University of Korea School of Medicine, Seoul, Republic of Korea

Table 3 Relationships between choriocapillaris(CC) flow voids and choroidal OCT parameters (CVI and CT)

Phansalkar3							
Acute CSC eye							
Fovea	Standardize β	R ²	P value	Extrafovea	Standardize β	R ²	P value
CT	0.570	0.325	0.009	CT	0.474	0.225	0.035
CVI	-0.189	0.036	0.424	CVI	-0.098	0.010	0.682
Recovered eye							
Fovea	Standardize β	R ²	P value	Extrafovea	Standardize β	R ²	P value
CT	0.589	0.346	0.006	CT	0.156	0.024	0.512
CVI	-0.220	0.048	0.352	CVI	-0.327	0.107	0.159
Fellow eye							
Fovea	Standardize β	R ²	P value	Extrafovea	Standardize β	R ²	P value
CT	-0.281	0.079	0.230	CT	-0.358	0.128	0.122
CVI	0.293	0.086	0.210	CVI	0.028	0.001	0.907
Control eye							
Fovea	Standardize β	R ²	P value	Extrafovea	Standardize β	R ²	P value
CT	0.004	<0.001	0.986	CT	-0.189	0.036	0.424
CVI	0.068	0.005	0.777	CVI	0.101	0.010	0.672
Phansalkar15							
Acute CSC eye							
Fovea	Standardize β	R ²	P value	Extrafovea	Standardize β	R ²	P value
CT	0.562	0.316	0.010	CT	0.462	0.214	0.040
CVI	-0.175	0.031	0.461	CVI	-0.094	0.009	0.694
Recovered eye							
Fovea	Standardize β	R ²	P value	Extrafovea	Standardize β	R ²	P value
CT	0.575	0.330	0.008	CT	0.125	0.016	0.599
CVI	-0.208	0.043	0.379	CVI	-0.312	0.098	0.180
Fellow eye							
Fovea	Standardize β	R ²	P value	Extrafovea	Standardize β	R ²	P value
CT	-0.233	0.054	0.323	CT	-0.347	0.121	0.134
CVI	0.206	0.042	0.384	CVI	0.206	0.043	0.383
Control eye							
Fovea	Standardize β	R ²	P value	Extrafovea	Standardize β	R ²	P value
CT	0.040	0.002	0.868	CT	-0.238	0.057	0.313
CVI	0.142	0.020	0.550	CVI	0.105	0.011	0.659

β , regression coefficient. p-values that were statistically significant are highlighted in bold

“Phansalkar 3 or 15” means using window radius 3, 15 pixels when Phansalkar threshold was applied to images

“Fovea” means the circular area of which radius is 50 pixels (= 703 μm)

“Extrafovea” means the circular area of which radius is 100 pixels (= 1406 μm)

CT: choroidal thickness; CVI: choroidal vascularity index