



HK-2 cell response to TGF- β highly depends on cell culture medium formulations

Gantsetseg Garmaa¹ · Anna Manžéger^{1,2} · Samaneh Haghighi¹ · Gábor Kökény^{1,2}

Accepted: 31 August 2023 / Published online: 26 September 2023
© The Author(s) 2023

Abstract

The immortalized human renal proximal tubular epithelial cell line HK-2 is most commonly used to study renal cell physiology and human kidney diseases with tubulointerstitial fibrosis such as diabetic nephropathy, obstructive uropathy or allograft fibrosis. Epithelial-to-mesenchymal transition (EMT) is the main pathological process of tubulointerstitial fibrosis in vitro. Transforming growth factor-beta (TGF- β) is a key inducer of EMT. Several pro-fibrotic gene expression differences have been observed in a TGF- β -induced EMT model of HK-2 cells. However, growth conditions and medium formulations might greatly impact these differences. We investigated gene and protein expression of HK-2 cells cultured in six medium formulations. TGF- β 1 increased the expression of *ACTA2*, *TGFB1*, *COL4A1*, *EGR2*, *VIM* and *CTGF* genes while reducing *PPARG* in all medium formulations. Interestingly, TGF- β 1 treatment either increased or decreased *EGR1*, *FN*, *IL6* and *C3* gene expression, depending on medium formulations. The cell morphology was slightly affected, but immunoblots revealed TGFB1 and vimentin protein overexpression in all media. However, fibronectin expression as well as the nuclear translocation of EGR1 was medium dependent. In conclusion, our study demonstrates that, using the HK-2 in vitro model of EMT, the meticulous selection of appropriate cell culture medium formulation is essential to achieve reliable scientific results.

Keywords Kidney fibrosis · TGF-beta · EMT · Cell culture · PTEC · Transcription factor

Abbreviations

ACTA2	Actin alpha 2, smooth muscle	CTGF	Connective tissue growth factor
BPE	Bovine pituitary extract	DMEM 2%	Dulbecco's modified Eagle's medium supplemented with 2% fetal bovine serum
ATCC	American type cell collection	DAPI	4',6-Diamidino-2-phenylindole
BCA	Bicinchoninic acid	DMEM 5%	Dulbecco's modified Eagle's medium supplemented with 2% fetal bovine serum
ciPTEC	Human conditionally immortalized proximal tubular epithelial cell line	DMEM 10%	Dulbecco's modified Eagle's medium supplemented with 2% fetal bovine serum
C3	Complement C3	DMEM/F12 10%	Dulbecco's Modified Eagle Medium/Nutrient Mixture F-12 supplemented with 10% fetal bovine serum
COL4A1	Collagen type IV alpha 1 chain	ECL	Enhanced chemiluminescence
		EGR1	Early growth response factor 1
		EGR2	Early growth response factor 2
		EMT	Epithelial-to-mesenchymal transition
		FN	Fibronectin
		GAPDH	Glyceraldehyde-3-phosphate dehydrogenase
		HK-2	Immortalized proximal tubule epithelial cell line

✉ Gábor Kökény
kokeny.gabor@med.semmelweis-univ.hu

Gantsetseg Garmaa
gantsetseg.garmaa@gmail.com

Anna Manžéger
anna.manzeger@gmail.com

Samaneh Haghighi
haghighi_samane@yahoo.com

¹ Institute of Translational Medicine, Semmelweis University, Nagyvárad tér 4, Budapest 1089, Hungary

² International Nephrology Research and Training Center, Semmelweis University, Nagyvárad tér 4, Budapest 1089, Hungary

HPV16	Human papillomavirus
IL6	Interleukin 6
KFSM	Keratinocyte serum-free medium
PTEC	Proximal tubular epithelial cells
PTEC 2%	Hormonally defined medium supplemented with 2% fetal bovine serum
PPARG	Peroxisome proliferator activated receptor gamma
PBS	Phosphate buffer saline
rEGF	Recombinant epidermal growth factor
RPTEC/TERT1	HTERT-immortalized epithelial cell line
SA7K	Pseudo-immortalized human renal proximal tubule epithelial cell line
SD	Standard deviation
TGF- β	Transforming growth factor beta cytokine
TGFB1	Transforming growth factor beta 1
VIM	Vimentin
ZO-1	Zonula occludens-1

Introduction

The kidney is a complex organ composed of at least 16 functionally distinct epithelial cell types (Balzer et al. 2022). The proximal tubular epithelial cells (PTEC), along with the tubular vasculature, make up > 80% of the renal cortex and are among the most abundant epithelial cell types (Balzer et al. 2022). They play essential roles in fluid, amino acid and sodium reabsorption and contribute significantly to

pathological changes within the cortical tubulointerstitium (Sato et al. 2020).

To study the function of PTEC cells, specialized isolation and culture techniques were developed that eliminate interference from other renal cell types, hemodynamics and neural activity. Several human cell lines, such as ciPTEC (Wilmer et al. 2010), RPTEC/TERT1 (Wieser et al. 2008) and SA7K (Li et al. 2017), have been newly established over the last decade. Still, the immortalized proximal tubule epithelial cell line HK-2 is most commonly used (Valdés et al. 2021) to study cell physiology, pharmacology (Jenkinson et al. 2012) and toxicology (Li et al. 2017) as well as various kidney diseases. HK-2 was derived from normal adult human renal cortex and immortalized through transduction with human papillomavirus (HPV16) E6/E7 genes, maintained for up to 30 passages (Ryan et al. 1994).

Regardless of the lack of some transporters, HK-2 is the mainly used in vitro model to study renal epithelial-to-mesenchymal transition (EMT) (Wang et al. 2018). EMT is recognized as an essential component of tissue fibrogenesis following kidney injury (Wang et al. 2018). Several studies show that tubular epithelial cells undergo phenotypic conversion in vitro after being incubated with fibrogenic transforming growth factor beta-1 (TGF- β 1) (Tian and Phillips 2003). During EMT, the cells lose their epithelial proteins, such as E-cadherin and zonula occludens-1 (ZO-1), and start to produce the mesenchymal markers vimentin, α -SMA, type I collagen, fibronectin (Liu 2004) and connective tissue growth factor (CTGF) (Cheng et al. 2006).

Although HK-2 cell providers recommended the use of serum-free medium supplemented with bovine pituitary extract (BPE) and human recombinant epidermal growth

Table 1 Cell culture media and their formulation

Medium abbreviations	Formulation
DMEM 2%	DMEM with 1000 mg/l glucose, L-glutamine, sodium bicarbonate (Invitrogen, Carlsbad, CA), and 2% fetal bovine serum (Invitrogen), 50 U/ml penicillin and 50 μ g/ml streptomycin (Invitrogen)
DMEM 5%	DMEM with 1000 mg/l glucose, L-glutamine, sodium bicarbonate (Invitrogen, Carlsbad, CA) and 5% fetal bovine serum (Invitrogen), 50 U/ml penicillin and 50 μ g/ml streptomycin (Invitrogen)
DMEM 10%	DMEM with 1000 mg/l glucose, L-glutamine, sodium bicarbonate (Invitrogen, Carlsbad, CA) and 10% fetal bovine serum (Invitrogen), 50 U/ml penicillin and 50 μ g/ml streptomycin (Invitrogen)
DMEM/F12 10%	DMEM/F12 medium contains a 1:1 mixture of DMEM medium and Ham's F12 medium, with L-glutamine, 15 mM HEPES, sodium bicarbonate (Invitrogen, Carlsbad, CA) and 10% fetal bovine serum (Invitrogen), 50 U/ml penicillin and 50 μ g/ml streptomycin (Invitrogen)
PTEC (hormonally defined)	DMEM/F12 (Invitrogen, Carlsbad, CA) supplemented with 2% fetal bovine serum (Invitrogen), 5 μ g/ml insulin, 5 μ g/ml transferrin, 5 ng/ml selenium, 40 ng/ml hydrocortisone, 5 pg/ml triiodo-L-thyronine, 50 U/ml penicillin and 50 μ g/ml streptomycin (Invitrogen)
KFSM	Keratinocyte serum-free growth medium (Invitrogen, Carlsbad, CA) supplemented with 0.05 mg/ml bovine pituitary extract (BPE), 5 ng/ml human recombinant epidermal growth factor (rEGF) (Invitrogen, Carlsbad, CA), 50 U/ml penicillin and 50 μ g/ml streptomycin (Invitrogen)

BPE Bovine pituitary extract, *rEGF* human recombinant epidermal growth factor, *DMEM 2%* DMEM with 2% FBS supplemented medium, *DMEM 5%* DMEM with 5% FBS supplemented medium, *DMEM/F12 10%* DMEM with 10% FBS supplemented medium, *PTEC* hormonally defined medium, *KFSM* keratinocyte serum-free growth medium

factor (rEGF) (Ryan et al. 1994), most of the biomedical studies documented the use of DMEM or DMEM/F12 media supplemented with fetal bovine serum (FBS) or hormones (Bozic et al. 2011, 2020; Kang et al. 2019). Thus, we aimed to establish and compare the TGF- β 1-induced EMT model in HK-2 cells in various culture medium formulations commonly used for epithelial cells. We further aimed to investigate how different culture media affect the pro-fibrotic gene expression pattern.

Here, we report different cell behavior (growth rate, morphology, differential gene and protein expression) upon changing the medium culture formulations. Our results underline that appropriate cell culture media should be carefully selected for specific scientific questions to avoid potential misinterpretation of experimental results obtained from the widely used HK-2 in vitro model.

Materials and methods

Cell culture

HK-2 cells were purchased from the American Type Cell Collection (ATCC, #CRL-2190). Cells were cultured in T75 flasks with six different growth media (Table 1) at 37 °C in a humid atmosphere of 95% air and 5% CO₂. The medium was refreshed every 48 to 72 h until the desired confluency. After 3 to 5 days, when cells reached 70–80% confluency, HK-2

cells were trypsinized and passaged at a 1:3 ratio or seeded on 6 or 24-well plates at the desired density for immediate experimental use. HK-2 cells were seeded $\sim 10^5$ cells or 3×10^4 per well in 6- or 24-well plates, respectively, and cultured in different growth media overnight, then serum starved for 24 h. Recombinant human TGF- β 1 (10 ng/ml, Sigma-Aldrich) was next added to the appropriate wells for 24 h. Microphotographs of treated and untreated cells in the culture plate wells were taken with Nikon Coolpix 4500 digital camera attached to a Nikon Eclipse TS100 microscope at 200 \times magnification. Cells were then harvested in

Table 2 Primer sequences (5'–3') used for quantitative PCR

<i>hACTA2</i> , forward	5'-AGATCAAGATCATTGCCC-3'
<i>hACTA2</i> , reverse	5'-TTCATCGTATTCCTGTTTC-3'
<i>hCOLAA1</i> , forward	5'-AAAGGGGAGATCAAGGGATAG-3'
<i>hCOLAA1</i> , reverse	5'-TCACCTTTTTCTCCAGGTAG-3'
<i>hC3</i> , forward	5'-CGGATCTTCACCGTTCAACCA-3'
<i>hC3</i> , reverse	5'-GATGCCTTCCGGTTCTCAA-3'
<i>hCTGF</i> , forward	5'-TTAAGAAGGGGCAAAAAGTGC-3'
<i>hCTGF</i> , reverse	5'-CATACTCCACAGAATTAGCTC-3'
<i>hEGRI</i> , forward	5'-ACCTGACCGCAGAGTCTTTT-3'
<i>hEGRI</i> , reverse	5'-GAGTGGTTTGGCTGGGGTAA-3'
<i>hFN</i> , forward	5'-CCATAGCTGAGAAGTGTTTTG-3'
<i>hFN</i> , reverse	5'-CAAGTACAATCTACCATCATCC-3'
<i>hGAPDH</i> , forward	5'-CATGAGAAGTATGACAACAGCCT-3'
<i>hGAPDH</i> , reverse	5'-AGTCCTTCCACGATACCAAAGT-3'
<i>hIL6</i> , forward	5'-GCAGAAAAAGGCAAAGAATC-3'
<i>hIL6</i> , reverse	5'-CTACATTTGCCGAAGAGC-3'
<i>hTGFB1</i> , forward	5'-GGA AAT TGA GGG CTT TCG CC-3'
<i>hTGFB1</i> , reverse	5'-CCG GTA GTG AACCCG TTG AT-3'
<i>hVIM</i> , forward	5'-GGAAACTAATCTGGATCACTC-3'
<i>hVIM</i> , reverse	5'-CATCTCTAGTTTCAACCGTC-3'
<i>hPPARG</i> , forward	5'-AAAGAAGCCAACACTAAACC-3'
<i>hPPARG</i> , reverse	5'-TGGTCATTTTCGTTAAAGGC-3'




Fig. 1 Effect of different medium formulations on HK-2 cell morphology in control (CTL) and TGF- β 1-treated (TGFb) groups. Light microscopy, 200 \times magnification; scale bar represents 100 μ m. 2% DMEM DMEM with 2% FBS supplemented medium, 5% DMEM DMEM with 5% FBS supplemented medium, 10% DMEM DMEM with 10% FBS supplemented medium, F12 10% DMEM/F12 with 10% FBS supplemented medium, PTEC hormonally defined medium, KFSM keratinocyte serum-free growth medium




Fig. 2 TGF-β1 mRNA (*TGFB1*) and protein (TGFB1) expression of HK-2 cells in different medium formulations. **a** Representative qPCR results of TGF-β1-induced effects on HK-2 cells cultured in different medium formulations for 24 h. The gene expression of *TGFB1* was normalized to *GAPDH* and indicated as fold expression relative to the respective controls (mean ± SD). Significant differences between the control vs. TGFβ ($n=7-16/\text{group}$) group shown as a red line and TGFβ vs. TGFβ groups between culture media shown as blue line

are indicated as $p < 0.05^*$, $p < 0.01^{**}$ and $p < 0.001^{***}$. Independent samples Mann-Whitney *U* test and Kruskal-Wallis test. **b** Representative immunoblot of TGFB protein expression is shown for each group; GAPDH was used as a loading control. *TGFB1*: transforming growth factor mRNA, *TGFB* transforming growth factor beta protein, *CTL* control group, *TGFβ* transforming growth factor beta-1 (10 ng/ml)-treated group

Trizol (Invitrogen, Thermo, USA) for total RNA isolation according to the manufacturer's protocol or in ice-cold RIPA buffer for protein extraction, as mentioned below.

RNA isolation and qPCR

Total RNA from Trizol samples was extracted according to the manufacturer's protocol. RNA concentration and purity were verified on a Nanodrop 2000 (Thermo, USA) and then reverse transcribed using the High-Capacity cDNA kit (Applied Biosystems/life Technologies, Carlsbad, CA, USA), as stated previously (Kökény et al. 2022). Each PCR reaction with specific primers (Table 2) was performed on a Bio-Rad CFX thermal cycler (Bio-Rad Hungary, Budapest, Hungary) in duplicates using the SensiFast SYBR Green PCR Master Mix (Bioline, Germany). The specificity and effectivity of PCR reactions were verified by melting curve analysis. Target gene expression was normalized to glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) mRNA expression using the $2^{-\Delta\Delta C_t}$ formula and expressed as fold expression relative to a control sample. Each gene expression is presented as mean ± standard deviation (SD).

Immunoblot

HK-2 cells were lysed in ice-cold RIPA buffer containing "Complete Mini" protease inhibitor cocktail (Roche, Mannheim, Germany). Protein concentration was determined using BCA assay (Thermo Scientific, Waltham, MA, USA). Equal amounts (20 μg) of protein were loaded in Laemmli buffer on 8% or 12% SDS-polyacrylamide gels. Separated

samples were transferred to nitrocellulose membranes, blocked with 5% skim milk and incubated with primary antibodies (1:2000, see Supplemental Table 1) overnight at 4 °C and then with the appropriate HRP-conjugated secondary antibodies as described previously (Kökény et al. 2022). Blots were visualized with enhanced chemiluminescence (ECL) detection kit (Thermo).

Immunofluorescence

The protein expressions of EGR1 and TGFB1 were assessed using immunofluorescence. HK-2 cells (10,000/well) were seeded on glass coverslips in a 24-well plate. After 24-h treatment with 10 ng/ml TGF-β1 in serum-free medium, cells were washed and fixed in methanol, then permeabilized with 0.25% Triton-X. Non-specific binding of secondary antibody was blocked using 2% donkey serum for 30 min. EGR1 (1:200) and TGF-β (1:200) antibodies (see Supplemental Table 1) or PBS for negative controls (omitting primary antibody) was applied overnight, and then the cells were incubated with donkey anti-rabbit IgG-A594 (1:200) (Jackson ImmunoResearch) at room temperature in the dark. Coverslips were mounted with Vectashield containing DAPI (Vector Laboratories, Burlingame, CA, USA). Cells were visualized and photographed under UV light using a Leica DMR-HC microscope at 400× magnification. Primary antibody specificity was evaluated on immunoblots.

Statistical analysis

Experimental data are presented as mean ± SD. Statistical analysis was performed using SPSS 28.0.0 for Windows




Fig. 3 EMT-related gene expression pattern of HK-2 cells. Gene expression of control (CTL) and TGF-β1-treated (TGFβ) cells in each culture medium is expressed as fold expression relative to the corresponding controls. Expression of each gene was normalized to GAPDH, and mean expression levels with standard deviation (\pm SD) are shown; **a** *FN* ($n=7-8$ /group), **b** *VIM* ($n=7-8$ /group), **c** *ACTA2* ($n=7-24$ /group). Significant differences within the groups are indicated as red lines with $p<0.05^*$, $p<0.01^{**}$ and $p<0.001^{***}$. Inter-group differences are blue lines with $p<0.05^*$, $p<0.01^{**}$ and $p<0.001^{***}$. Independent samples Mann-Whitney *U* test and Kruskal-Wallis test

(SPSS Inc). Shapiro-Wilk test was performed to analyze normal distribution of the data. Pairwise comparisons were conducted using the independent samples Mann-Whitney *U* test or Kruskal-Wallis test, followed by Bonferroni correction, as indicated, to analyze normalized expression levels. Spearman and Pearson's correlation was used for the relationship between continuous and ordinal variables. The significance level was $p<0.05$ at the 95% confidence level.

Results

Cell morphology

We examined the effect of six culture medium formulations (see Table 1) on cell morphology. Cells without TGF-β1 treatment depicted epithelial morphology, whereas TGF-β1 treatment (10 ng/ml) for 24-h induced a slightly elongated morphology in all medium formulations (Fig. 1). However, HK-2 cultured in KFSM appeared smaller and did not replicate as fast as cells cultured in DMEM or DMEM F12-based formulations on day 5 after the passage. At sub-confluent density, HK-2 in DMEM 5% FBS most likely appeared round-shaped and polygonal (Fig. 1). HK-2 cells reached 80% confluence at culture day 7 in KFSM and day 5 in DMEM and DMEM F12-based media. On day 7, space was still visible in a T-75 flask with KFSM and DMEM 2% FBS medium. In addition, cells in DMEM with 10% FBS appeared larger (Fig. 1).

Gene and protein expression variations related to EMT and pro-fibrotic markers

TGF-β1 promoted phenotype changes of HK-2 cells toward EMT, as confirmed by the slightly elongated morphology (Fig. 1), the significantly increased *TGFβ1* gene (Fig. 2a) and protein expression (Fig. 2b), and highly expressed mesenchymal markers by qPCR (Fig. 3a–c). EMT-related changes at the mRNA level were observed regardless of the culture medium formulations (Fig. 3). The mRNA expression of vimentin (*VIM*) was induced in all formulations (Fig. 3b); however, immunoblot showed its overexpression in all media except DMEM 5% (Supplementary Fig. S1a, c). TGF-β1 partially induced the mRNA expression of *FN* in PTEC 2%, DMEM F12 10% and KFSM (Fig. 3a) and the protein expression in almost all media except DMEM/F12 10% and KFSM (Supplementary Fig. S1b, c).

CTGF, *COL4A1* and *EGR2* expressions were significantly increased upon TGF-β1 treatment in all investigated media (Fig. 4a, b, Fig. 5b). Interestingly, medium formulations had a substantial impact on pro-fibrotic *EGR1* expression. TGF-β1 treatment increased *EGR1* mRNA expression significantly only in DMEM 5% and DMEM/F12 10% media, but in PTEC 2% and DMEM 10%, *EGR1* decreased (Fig. 5a, Table 3).

PPARG expression decreased significantly in all culture media upon TGF-β1 treatment (Fig. 5c), although there were differences between culture media; the most potent effect was observed using PTEC formulation (0.39 ± 0.08 in PTEC 2% vs. 0.68 ± 0.10 in KFSM and 0.76 ± 0.09 in DMEM F12 10%).

Among the inflammatory markers, *IL6* mRNA was not induced in TGF-β1-treated HK-2 cells cultured in PTEC 2%




Fig. 4 Pro-fibrotic gene expression pattern depends on culture medium formulation. Gene expression of control (CTL) and TGF-β1-treated (TGFβ) HK-2 cells in each culture medium are expressed as fold expression relative to the corresponding controls as follows: **a** CTGF ($n=7-8$ /group), **b** COL4A1 ($n=5-6$ /group). Expression of

medium in addition to the DMEM 2% (Fig. 6a, Table 3). The expression of the complement C3 protein-coding gene, C3, decreased in KFSM and increased in DMEM 5% but did not change in any other media upon TGFβ1 treatment (Fig. 6b).

Effect of serum concentration and additive hormones on the EMT process

To clarify the effect of serum concentration in a culture medium, we cultured HK-2 cells in DMEM supplemented with 2%, 5% or 10% FBS. By Spearman's correlation, there was no significant correlation between the percentage of FBS and the expression of mRNAs listed in Figs. 3, 4, 5 and 6 in the abovementioned three media.

To further clarify the effect of different formulations besides FBS, we cultured HK-2 cells on a hormonally defined medium (PTEC, Table 2). Notably, immunoblot depicted significantly (1.7-fold) overexpressed fibronectin in PTEC medium (Supplemental Fig. S1b).

We also compared DMEM vs. DMEM F12-based medium supplemented with 10% FBS. The mean mRNA expression of *FN*, *ACTA2*, *EGR1* and *EGR2* was differentially expressed in TGF-β1-induced HK-2 cells cultured in these two media ($p < 0.05$) (Figs. 3a, c, 5a, b). Interestingly, FN protein expression was reduced by 0.73-fold in DMEM F12 10% (Supplementary Fig. S1b).

Finally, these data indicated that TGF-β1 stimulates partial EMT and pro-fibrotic program of HK-2 cells differently depending on cell culture medium formulations.

Different cellular localization of EGR1 protein by immunocytochemistry

Based on the differential expression of mRNAs in culture medium formulations, we investigated the expression

each gene was normalized to GAPDH and shown as mean \pm SD. Significant differences within the groups are indicated as red lines with $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$. Inter-group differences are blue lines with $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$. Independent samples Mann-Whitney *U* test and Kruskal-Wallis test

pattern of EGR1 in TGF-β-induced HK-2 cells by immunofluorescence. TGF-β induced EGR1 translocation to the nucleus in DMEM F12 10% and KFSM media indicating transcriptional activation (Fig. 7a). Cells grown in other media (DMEM 2%, 5%, 10% and PTEC 2%) depicted only cytoplasmic overexpression of EGR1 (Fig. 7a). Still, according to both gene expression and immunoblot analysis, we observed TGFB1 protein expression induced by TGF-β1 treatment independent of the used culture medium (Fig. 7b).

Discussion

HK-2 is a widely used in vitro model to study EMT in kidney tubules. However, published studies use different cell culture media (Bozic et al. 2011, 2020; Li et al. 2005), which may lead to inhomogeneous or even controversial experimental results. Thus, we investigated how the most commonly used culture medium formulations affect cell behavior upon TGF-β1 administration, depicted as cell morphology, gene and protein expression pattern. To the best of our knowledge, this is the first comparative study on HK-2 cell morphology and expression pattern in different culture medium formulations.

Cell morphology

In general, TGF-β1 induced EMT in all used culture media. Our results indicate that HK-2 cells cultured in serum-free and DMEM 2% medium showed a slow growth rate and smaller cell size than those cultured in other media. Based on our study, we found that cells grown in a higher proportion of FBS (10%) appeared enlarged. However, no evidence suggests a connection between PTEC cell size and FBS concentration. Interestingly, one study reported an increase in




Fig. 5 Gene expression pattern of transcription factors upon culture medium formulations. Gene expression of control (CTL) and TGF- β 1-treated (TGFb) HK-2 cells in each culture medium are expressed as fold expression relative to the corresponding controls as follows: **a** *EGR1* ($n=5-12$ /group), **b** *EGR2* ($n=5-14$ /group), **c** *PPARG* ($n=5-13$ /group). Expression of each gene was normalized to GAPDH and shown as mean \pm SD. Significant differences within the groups are indicated as red lines with * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$. Inter-group differences are blue lines with * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$. Independent samples Mann-Whitney *U* test and Kruskal-Wallis test

size, similar to human pancreatic cancer spheres (Sasaki et al. 2019), when FBS was present. Further research is necessary to fully understand the effects of FBS on cell growth and size. In contrast, cells grown in a hormonally defined medium looked more flattened and elongated. Notably, the slower growth rate of cells in the DMEM 2% medium was predictable as FBS is the main component of culture media supporting cell growth (Yao and Asayama 2016; Puck et al. 1958).

Gene and protein expression

Despite the morphological differences, gene expression of the classic EMT markers *ACTA2*, *TGF β* , *VIM* and *CTGF* was significantly upregulated in TGF- β 1-induced HK-2 cells regardless of the culture medium, indicating that any of these medium formulations is feasible for investigation of the expression changes of these genes in a TGF- β 1-induced system. This corroborates previously published data (Brennan et al. 2012). Our EMT model partially induced the expression of mesenchymal (α -SMA, vimentin, fibronectin) markers. Notably, protein expression differed somewhat from the corresponding mRNA expression results. Khundmiri and colleagues noted that protein expression in HK-2 cells could differ from mRNA expression seen in human kidney and primary PTEC because of the low percentage (26%) of proximal tubule-specific transcripts detected in the HK-2 cell line, as obtained from RNA-Seq (Khundmiri et al. 2021).

In addition, the observed differential gene and protein expression could be due to a lack of serum in the serum-free medium or interactions with other additives in the culture medium. Regarding specific additives, EGF is considered a mitogen and supports cell proliferation, but its combination with TGF- β has been reported to cause primary PTEC cell hypertrophy and excessive accumulation of extracellular matrix proteins (Franch et al. 1995). This may explain the higher expression of *COL4A1* in the KFSM medium supplemented with EGF compared to other media in our study.

Peroxisome proliferator receptor-gamma (PPAR- γ) is a nuclear receptor superfamily member and has been demonstrated to ameliorate renal fibrogenesis (Zhao et al. 2016; Németh et al. 2019). *PPARG* is downregulated in fibrosis, and PPAR- γ agonists are promising therapeutic agents in fibrotic diseases (Kökény et al. 2020; Ghallab and Seddek 2020; Németh et al. 2019). Our results corroborate these studies depicting a strong TGF- β 1-related reduction of *PPARG* expression in all investigated medium formulations (Fig. 5c).

Early growth response factor-1 and -2 (*EGR1*, *EGR2*) have been implicated as important pro-fibrotic transcription factors in kidney diseases (Bhattacharyya et al. 2013, 2011; Ho et al. 2016; Sun et al. 2014; Vollmann et al. 2017). We have previously demonstrated a *PPARG*-*EGR1* axis in TGF- β 1-induced renal fibrosis (Németh et al. 2019). Interestingly *EGR1* expression upon TGF- β 1 stimulation fluctuated from medium to medium in our present study. Despite increased TGF- β mRNA and protein expression in parallel with the reduced *PPARG* in all medium formulations, we observed nuclear translocation of *EGR1* in only two media. This suggests a culture medium-dependent transcriptional activity for *EGR1*. In contrast, *EGR2* expression was significantly and similarly induced by TGF- β 1 in all investigated media.

Table 3 TGF- β -induced pro-fibrotic EGR1 and inflammatory IL-6 gene expression responses based on cell culture media

Cell culture medium and treatment	EGR1 gene expression			IL-6 gene expression		
	Mean \pm SD	Effect of TGF- β treatment (TGFb vs CTL)	Effect of serum content (TGFb vs TGFb)	Mean \pm SD	Effect of TGF- β treatment (TGFb vs CTL)	Effect of serum content (TGFb vs TGFb)
DMEM 2% CTL DMEM 2% TGFb	0,91 \pm 0,22 0,70 \pm 0,15	p=0.22 (ns)		1,05 \pm 0,20 1,14 \pm 0,18	p=0.42 (ns)	p=0.046 vs F12 10%
DMEM 5% CTL DMEM 5% TGFb	1,00 \pm 0,08 1,54 \pm 0,41	p=0.013	p=0.033 vs DMEM 10% p=0.02 vs PTEC	1,00 \pm 0,09 1,55 \pm 0,51	p=0.01	
DMEM 10% CTL DMEM 10% TGFb	1,00 \pm 0,11 0,71 \pm 0,18	p<0.001	p=0.033 vs DMEM 5% p=0.006 vs F12 10%	1,00 \pm 0,14 1,56 \pm 0,21	p=0.008	
F12 10% CTL F12 10% TGFb	1,00 \pm 0,09 1,91 \pm 0,41	p<0.001	p=0.006 vs DMEM 10% p=0.003 vs PTEC	1,00 \pm 0,07 1,68 \pm 0,34	p=0.03	p=0.046 vs DMEM 2% p=0.003 vs PTEC
PTEC CTL PTEC TGFb	1,05 \pm 0,11 0,67 \pm 0,23	p<0.001	p=0.02 vs DMEM 5% p=0.003 vs F12 10%	0,96 \pm 0,16 0,95 \pm 0,14	p=0.931 (ns)	p=0.003 vs F12 10% p=0.039 vs KSFM
KSFM CTL KSFM TGFb	1,00 \pm 0,08 0,72 \pm 0,26	p=0.065 (ns)		1,00 \pm 0,08 1,58 \pm 0,23	p=0.002	p=0.039 vs PTEC
Statistical method:		Mann-Whitney	Kruskal-Wallis		Mann-Whitney	Kruskal-Wallis

Legend: The effect of TGF- β treatment (TGFb) in each cell culture medium was assessed pairwise using the Mann-Whitney U-test and are indicated as p values (TGFb vs CTL). The effect of different cell culture media formulations on TGF- β -induced cellular responses was further analyzed with Kruskal-Wallis test for multiple comparisons and are indicated as p values with the corresponding group (ns=not significant).

Interestingly, the expression pattern of inflammatory mediators IL6 and C3 was also dependent on cell culture media. In the kidney, proximal tubular cells are exposed to various cytokines from glomerular filtration, and they also synthesize complement proteins, including C3 and factors B and H (Gerritsma et al. 1997; Peake et al. 1999). In our study, C3 pro-

hormones and EGF (Gerritsma et al. 1998), supporting our observation as C3 was downregulated by 50% in the serum-free KSFM. The differences in FBS-supplemented media need to be investigated.

Study limitations





Fig. 6 Cell culture medium-dependent expression of inflammatory genes. Gene expression of control (CTL) and TGF- β 1-treated (TGFb) HK-2 cells in each culture medium are expressed as fold expression relative to the corresponding controls as follows: **a** IL6 ($n=5-7$), **b** C3 ($n=7-11$ /group). Gene expression was normalized to GAPDH

and shown as mean \pm SD. Significant differences within the groups are indicated as red lines with * $p \leq 0.05$, ** $p \leq 0.01$ and *** $p \leq 0.001$. Inter-group differences are blue lines with * $p \leq 0.05$, ** $p \leq 0.01$ and *** $p \leq 0.001$. Independent samples Mann-Whitney U test and Kruskal-Wallis test

duction seemed to depend on the medium serum content as in DMEM and DMEM/F12-based formulations, and TGF- β -induced C3 overexpression only in DMEM 5% FBS medium. C3 has been demonstrated to decrease upon TGF- β 1 in primary PTEC cells in a serum-free medium supplemented with

First, it is necessary to consider the shortcomings of cultured HK-2 cells as a model of the proximal tubule. HK-2 cells and many of the available proximal tubule model cell lines fail to replicate the differential expression of apical/basal membrane transporters and metabolizing enzymes. These are one of the

Fig. 7 EGR1 and TGFB1 immunocytochemistry of TGF- β 1 induced HK-2 cells. EGR1 (a) and TGFB1 (b) protein expression of control and TGF- β 1-treated (10 ng/ml for 24 h) cells in different medium formulations. Immunofluorescence, 400 \times magnification. Neg. contr.: representative photograph of the negative control sample (omitting the primary antibodies). Scale bar represents 50 μ m. Red stain: EGR1 or TGFB1; blue stain: nuclear stain (DAPI). Magenta: double stain



characteristics of renal proximal tubular cells in vivo (Sanchez-Romero et al. 2016) with additional display of metabolic zonation with different enzyme machinery and transporters along proximal tubule segments S1, S2 and S3 (Lee et al. 2015), and HK-2 cells might be representative of only one of them. Nevertheless, HK-2 cell line is the most widely used experimental model in translational research (Valdés et al. 2021).

Investigating the long-term effect of TGF- β 1-induced gene and protein expression and regulation might provide a more detailed understanding of the effect of different medium formulations in HK-2 cells. We also did not investigate medium additives individually; further studies probably need to

investigate the effect on single additives, hormones, growth factors and insulin, e.g., on the TGF- β -induced EMT model in HK-2 cells.

Conclusions

In conclusion, our study demonstrates how cell culture medium formulation (an essential component for cell growth and morphology) and its modification can affect the HK-2 cell behavior during TGF- β 1 induced in vitro EMT. TGF- β 1 stimulates the HK-2 EMT process in any investigated

culture medium formulations but with different EGR1 activation and inflammatory response. Choosing cell culture medium formulations and explaining the expression of EMT and pro-fibrotic markers in TGF- β 1-induced HK-2 cells is essential and should not be overlooked.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00418-023-02237-x>.

Acknowledgements The invaluable technical assistance of Rita Bencs and Krisztina Fazekas is greatly acknowledged. Financial support by the Stipendium Hungaricum Scholarship (Tempus Public Foundation, Hungary) and SE 250+ Excellence PhD Scholarship (Simmelweis University) to GG as well as by the Bolyai Scholarship of the Hungarian Academy of Sciences (BO/00304/20/5 to GK) and the ÚNKP Bolyai+ Scholarship from the Hungarian Ministry of Innovation and Technology and the National Research, Development and Innovation Office (to GK) is greatly acknowledged. Institutional internal funding has been attributed to GK.

Author contributions G.G. performed cell culture and qPCR experiments, statistical evaluation and drafted the manuscript. S.H. performed cell culture experiments. A.M. performed immunoblots and immunocytochemistry and supervised statistical evaluation. G.K. conceptualized the study, prepared the figures and wrote the final manuscript. All authors reviewed the manuscript.

Funding Open access funding provided by Semmelweis University. Tempus Közalapítvány, Stipendium Hungaricum, Stipendium Hungaricum, Semmelweis Egyetem, Institutional internal funding, Institutional internal funding, Magyar Tudományos Akadémia, Bolyai Scholarship BO/00304/20/5, Innovációs és Technológiai Minisztérium, ÚNKP Bolyai + Scholarship.

Data availability Experimental data are available upon reasonable request from the corresponding author.

Declarations

Conflict of interest None of the authors have any competing interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Balzer MS, Rohacs T, Susztak K (2022) How many cell types are in the kidney and what do they do? *Annual Rev Physiol* 84:507–531. <https://doi.org/10.1146/annurev-physiol-052521-121841>
- Bhattacharyya S, Wu M, Fang F, Tourtellotte W, Feghali-Bostwick C, Varga J (2011) Early growth response transcription factors: key mediators of fibrosis and novel targets for anti-fibrotic therapy. *Matrix Biol* 30(4):235–242. <https://doi.org/10.1016/j.matbio.2011.03.005>
- Bhattacharyya S, Fang F, Tourtellotte W, Varga J (2013) Egr-1: new conductor for the tissue repair orchestra directs harmony (regeneration) or cacophony (fibrosis). *J Pathol* 229(2):286–297. <https://doi.org/10.1002/path.4131>
- Bozic M, de Rooij J, Parisi E, Ortega MR, Fernandez E, Valdivielso JM (2011) Glutamatergic signaling maintains the epithelial phenotype of proximal tubular cells. *J Am Soc Nephrol* 22(6):1099–1111. <https://doi.org/10.1681/ASN.2010070701>
- Bozic M, Caus MA, Rodrigues-Diez RA, Pedraza NA, Ruiz-Ortega M, Garí E, Galle P, Panadés MJ, Martínez A, Fernández E, Valdivielso JM (2020) Protective role of renal proximal tubular alpha-synuclein in the pathogenesis of kidney fibrosis. *Nat Commun* 11(1):1943. <https://doi.org/10.1038/s41467-020-15732-9>
- Brennan EP, Morine MJ, Walsh DW, Roxburgh SA, Lindenmeyer MT, Brazil DP, Gaora PO, Roche HM, Sadlier DM, Cohen CD, Consortium G, Godson C, Martin F (2012) Next-generation sequencing identifies TGF-beta1-associated gene expression profiles in renal epithelial cells reiterated in human diabetic nephropathy. *Biochim Biophys Acta* 4:589–599. <https://doi.org/10.1016/j.bbadis.2012.01.008>
- Cheng O, Thuillier R, Sampson E, Schultz G, Ruiz P, Zhang X, Yuen PST, Mannon RB (2006) Connective tissue growth factor is a biomarker and mediator of kidney allograft fibrosis. *Am J Transplant* 6(10):2292–2306. <https://doi.org/10.1111/j.1600-6143.2006.01493.x>
- Franch HA, Shay JW, Alpern RJ, Preisig PA (1995) Involvement of pRB family in TGF beta-dependent epithelial cell hypertrophy. *J Cell Biol* 129(1):245–254. <https://doi.org/10.1083/jcb.129.1.245>
- Gerritsma JS, Gerritsen AF, De Ley M, van Es LA, Daha MR (1997) Interferon-gamma induces biosynthesis of complement components C2, C4 and factor H by human proximal tubular epithelial cells. *Cytokine* 9(4):276–283. <https://doi.org/10.1006/cyto.1996.0164>
- Gerritsma JSJ, van Kooten C, Gerritsen AF, van Es LA, Daha MR (1998) Transforming growth factor- β 1 regulates chemokine and complement production by human proximal tubular epithelial cells. *Kidney Int* 53(3):609–616. <https://doi.org/10.1046/j.1523-1755.1998.00799.x>
- Ghallab A, Seddek A (2020) PPAR γ as therapeutic target for anti-fibrotic therapy. *EXCLI J* 19:227–229. <https://doi.org/10.17179/excli2020-1136>
- Ho LC, Sung JM, Shen YT, Jheng HF, Chen SH, Tsai PJ, Tsai YS (2016) Egr-1 deficiency protects from renal inflammation and fibrosis. *J Mol Med (berl)* 94(8):933–942. <https://doi.org/10.1007/s00109-016-1403-6>
- Jenkinson SE, Chung GW, van Loon E, Bakar NS, Dalzell AM, Brown CDA (2012) The limitations of renal epithelial cell line HK-2 as a model of drug transporter expression and function in the proximal tubule. *Pflgers Arch* 464(6):601–611. <https://doi.org/10.1007/s00424-012-1163-2>
- Kang HM, Lim JH, Noh KH, Park D, Cho HS, Susztak K, Jung CR (2019) Effective reconstruction of functional organotypic kidney spheroid for in vitro nephrotoxicity studies. *Sci Rep* 9(1):17610. <https://doi.org/10.1038/s41598-019-53855-2>
- Khundmiri SA, Chen LA, Lederer ED, Yang CR, Knepper MA (2021) Transcriptomes of major proximal tubule cell culture models. *J Am Soc Nephrol* 32(1):86–97. <https://doi.org/10.1681/ASN.2020010009>
- Kökény G, Calvier L, Legchenko E, Chouvarine P, Mózes MM, Hansmann G (2020) PPAR γ is a gatekeeper for extracellular matrix and vascular cell homeostasis: beneficial role in pulmonary hypertension and renal/cardiac/pulmonary fibrosis. *Curr Opin Nephrol*

- Hypertens 29(2):171–179. <https://doi.org/10.1097/MNH.0000000000000580>
- Kökény G, Németh Á, Kopp JB, Chen W, Oler AJ, Manzégér A, Rosivall L, Mózes MM (2022) Susceptibility to kidney fibrosis in mice is associated with early growth response-2 protein and tissue inhibitor of metalloproteinase-1 expression. *Kidney Int* 102(2):337–354. <https://doi.org/10.1016/j.kint.2022.03.029>
- Lee JW, Chou CL, Knepper MA (2015) Deep sequencing in microdissected renal tubules identifies nephron segment-specific transcriptomes. *J Am Soc Nephrol* 26(11):2669–2677. <https://doi.org/10.1681/ASN.2014111067>
- Li H, Ruan XZ, Powis SH, Fernando R, Mon WY, Wheeler DC, Moorhead JF, Varghese Z (2005) EPA and DHA reduce LPS-induced inflammation responses in HK-2 cells: evidence for a PPAR-gamma-dependent mechanism. *Kidney Int* 67(3):867–874. <https://doi.org/10.1111/j.1523-1755.2005.00151.x>
- Li S, Zhao J, Huang R, Steiner T, Bourner M, Mitchell M, Thompson DC, Zhao B, Xia M (2017) Development and application of human renal proximal tubule epithelial cells for assessment of compound toxicity. *Curr Chem Genom Transl Med* 11:19–30. <https://doi.org/10.2174/2213988501711010019>
- Liu Y (2004) Epithelial to mesenchymal transition in renal fibrogenesis: pathologic significance, molecular mechanism, and therapeutic intervention. *J Am Soc Nephrol* 15(1):1–12. <https://doi.org/10.1097/01.asn.0000106015.29070.e7>
- Németh Á, Mózes MM, Calvier L, Hansmann G, Kökény G (2019) The PPAR γ agonist pioglitazone prevents TGF- β induced renal fibrosis by repressing EGR-1 and STAT3. *BMC Nephrol* 20(1):245. <https://doi.org/10.1186/s12882-019-1431-x>
- Peake PW, O'Grady S, Pussell BA, Charlesworth JA (1999) C3a is made by proximal tubular HK-2 cells and activates them via the C3a receptor. *Kidney Int* 56(5):1729–1736. <https://doi.org/10.1046/j.1523-1755.1999.00722.x>
- Puck TT, Cieciera SJ, Robinson A (1958) Genetics of somatic mammalian cells. III. Long-term cultivation of euploid cells from human and animal subjects. *J Exp Med* 108(6):945–956. <https://doi.org/10.1084/jem.108.6.945>
- Ryan MJ, Johnson G, Kirk J, Fuerstenberg SM, Zager RA, Torok-Storb B (1994) HK-2: an immortalized proximal tubule epithelial cell line from normal adult human kidney. *Kidney Int* 45(1):48–57. <https://doi.org/10.1038/ki.1994.6>
- Sanchez-Romero N, Schophuizen CM, Gimenez I, Masereeuw R (2016) In vitro systems to study nephrotoxicology: 2D versus 3D models. *Eur J Pharmacol* 790:36–45. <https://doi.org/10.1016/j.ejphar.2016.07.010>
- Sasaki N, Toyoda M, Hasegawa F, Fujiwara M, Gomi F, Ishiwata T (2019) Fetal bovine serum enlarges the size of human pancreatic cancer spheres accompanied by an increase in the expression of cancer stem cell markers. *Biochem Biophys Res Commun* 514(1):112–117. <https://doi.org/10.1016/j.bbrc.2019.04.117>
- Sato Y, Takahashi M, Yanagita M (2020) Pathophysiology of AKI to CKD progression. *Semin Nephrol* 40(2):206–215. <https://doi.org/10.1016/j.semnephrol.2020.01.011>
- Sun S, Ning X, Zhai Y, Du R, Lu Y, He L, Li R, Wu W, Sun W, Wang H (2014) Egr-1 mediates chronic hypoxia-induced renal interstitial fibrosis via the PKC/ERK pathway. *Am J Nephrol* 39(5):436–448. <https://doi.org/10.1159/000362249>
- Tian YC, Phillips AO (2003) TGF-beta1-mediated inhibition of HK-2 cell migration. *J Am Soc Nephrol* 14(3):631–640. <https://doi.org/10.1097/01.asn.0000053418.56286.5e>
- Valdés A, Lucio-Cazaña FJ, Castro-Puyana M, García-Pastor C, Fiehn O, Marina ML (2021) Comprehensive metabolomic study of the response of HK-2 cells to hyperglycemic hypoxic diabetic-like milieu. *Sci Rep* 11(1):5058. <https://doi.org/10.1038/s41598-021-84590-2>
- Vollmann EH, Cao L, Amatucci A, Reynolds T, Hamann S, Dalkilic-Liddle I, Cameron TO, Hossbach M, Kauffman KJ, Mir FF, Anderson DG, Novobrantseva T, Koteliansky V, Kisseleva T, Brenner D, Duffield J, Burkly LC (2017) Identification of novel fibrosis modifiers by in vivo siRNA silencing. *Mol Ther Nucleic Acids* 7:314–323. <https://doi.org/10.1016/j.omtn.2017.04.014>
- Wang Z, Divanyan A, Jourde'heuil FL, Goldman RD, Ridge KM, Jourde'heuil D, Lopez-Soler RI (2018) Vimentin expression is required for the development of EMT-related renal fibrosis following unilateral ureteral obstruction in mice. *Am J Physiol Renal Physiol* 315(4):F769–F780. <https://doi.org/10.1152/ajprenal.00340.2017>
- Wieser M, Stadler G, Jennings P, Streubel B, Pfaller W, Ambros P, Riedl C, Katinger H, Grillari J, Grillari-Voglauer R (2008) hTERT alone immortalizes epithelial cells of renal proximal tubules without changing their functional characteristics. *Am J Physiol Renal Physiol* 295(5):F1365–F1375. <https://doi.org/10.1152/ajprenal.90405.2008>
- Wilmer MJ, Saleem MA, Masereeuw R, Ni L, van der Velden TJ, Russel FG, Mathieson PW, Monnens LA, van den Heuvel LP, Levchenko EN (2010) Novel conditionally immortalized human proximal tubule cell line expressing functional influx and efflux transporters. *Cell Tissue Res* 339(2):449–457. <https://doi.org/10.1007/s00441-009-0882-y>
- Yao T, Asayama Y (2016) Human preimplantation embryo culture media: past, present, and future. *J Mamm Ova Res* 33(1):17–34. <https://doi.org/10.1274/jmor.33.17>
- Zhao M, Chen Y, Ding G, Xu Y, Bai M, Zhang Y, Jia Z, Huang S, Zhang A (2016) Renal tubular epithelium-targeted peroxisome proliferator-activated receptor- γ maintains the epithelial phenotype and antagonizes renal fibrogenesis. *Oncotarget* 7(40):64690–64701. <https://doi.org/10.18632/oncotarget.11811>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.