Correction to: High-but not moderate-intensity exercise acutely attenuates hypercapnia-induced vasodilation of the internal carotid artery in young men

Rintaro Sakamoto¹ · Masaki Katayose² · Yutaka Yamada¹ · Toru Neki² · Tatsuki Kamoda² · Katsuyuki Tamai² · Kotomi Yamazaki² · Erika Iwamoto²

Published online: 24 August 2021
© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Correction to: European Journal of Applied Physiology
<https://doi.org/10.1007/s00421-021-04721-5>

The original version of this article unfortunately contained a mistake. The formula used for SR_{AUC} calculation under the section “Hypercapnia-induced vasodilation and ICA variables” of the Method section was incorrect.

The correct formula should read

$\Sigma [1/2 (x_{i+1} - x_i) (y_{i+1} - y_i) + (x_{i+1} - x_i) y_i]$, where x is time and y is SR .

This correction did not have any major impact on the main results or their interpretation.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00421-021-04721-5>.

✉ Erika Iwamoto
e.iwamoto@sapmed.ac.jp

¹ Department of Physical Therapy, Graduate School of Health Sciences, Sapporo Medical University, Sapporo, Japan

² School of Health Science, Sapporo Medical University, Sapporo, Japan