Correction to: Brown adipose tissue fat-fraction is associated with skeletal muscle adiposity

Madoka Ogawa^{1,2,12} · Kalle Koskensalo^{3,4,5,6} · Sanna Laurila^{3,4,6,7} · Milja Holstila^{3,4,8} · Minna Laheesmaa^{3,4} · Kirsi A. Virtanen^{4,9} · Hidehiro Iida^{3,4} · Hiroshi Akima^{1,10} · Pirjo Nuutila^{3,4,11}

Published online: 23 October 2021
© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Correction to: European Journal of Applied Physiology
<https://doi.org/10.1007/s00421-021-04816-z>

The original version of this article unfortunately contained a mistake. In Table 2, Multifidus lumborum column was mistakenly included.

The corrected Table 2 is given below.

The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00421-021-04816-z>.

✉ Madoka Ogawa
madoka.ogawa20@gmail.com

- ¹ Graduate School of Education and Human Development, Nagoya University, Aichi, Japan
- ² Nippon Sport Science University, Tokyo, Japan
- ³ Turku PET centre, University of Turku, Turku, Finland
- ⁴ Turku PET centre, Turku University Hospital, Turku, Finland
- ⁵ Department of Medical Physics, Turku University Hospital, Turku, Finland
- ⁶ Heart Center, Turku University Hospital, Turku, Finland

- ⁷ Satakunta Central Hospital, Pori, Finland
- ⁸ Medical Imaging Centre of Southwest Finland, Turku University Hospital, Turku, Finland
- ⁹ Institute of Public Health and Clinical Nutrition, University of Eastern Finland, Kuopio, Finland
- ¹⁰ Research Center of Health, Physical Fitness and Sports, Nagoya University, Aichi, Japan
- ¹¹ Department of Endocrinology, Turku University Hospital, Turku, Finland
- ¹² Kyoto Sangyo University, Kyoto, Japan

Table 2 Relationship between cold stimulated brown adipose tissue (BAT) glucose uptake rate and fat-fraction and age, body mass index (BMI), body fat percentage and waist circumference

	Glucose uptake of BAT ($\mu\text{mol}/100\text{ g}/\text{min}$)		Fat-fraction of BAT (%)	
	r_s	n	r_s	n
Age (years)	−0.307	18	0.158	19
BMI (kg/m^2)	0.004	18	0.450	19
Body fat (%)	−0.042	18	0.480*	19
Waist circumference (cm)	−0.128	18	0.562*	19

* $P < 0.05$