



## Correction to: Prevalence of Triad–RED–S symptoms in high-level Kenyan male and female distance runners and corresponding control groups

Lauri Önnik<sup>1</sup> · Martin Mooses<sup>1</sup> · Silva Suvi<sup>1</sup> · Diresibachew W. Haile<sup>2,3</sup> · Robert Ojiambo<sup>2,4</sup> · Amy R. Lane<sup>5</sup> · A. C. Hackney<sup>5</sup>

Published online: 24 October 2021  
© Springer-Verlag GmbH Germany, part of Springer Nature 2021

**Correction to:** *European Journal of Applied Physiology*  
<https://doi.org/10.1007/s00421-021-04827-w>

The original version of this article unfortunately contained a mistake. Author name Diresibachew W. Haile was incorrectly written as Diresibashew W. Haile.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The original article can be found online at <https://doi.org/10.1007/s00421-021-04827-w>.

---

✉ Lauri Önnik  
lauri.onnik@ut.ee

<sup>1</sup> University of Tartu, Ujula 4, 51008 Tartu, Estonia

<sup>2</sup> Moi University, Eldoret, Kenya

<sup>3</sup> Addis Ababa University, Addis Ababa, Ethiopia

<sup>4</sup> University of Global Health Equity, Butaro, Rwanda

<sup>5</sup> University of North Carolina, Chapel Hill, NC, USA