



Effect of repeated post-resistance exercise cold or hot water immersion on in-season inflammatory responses in academy rugby players: a randomised controlled cross-over design

Barry G. Horgan^{1,2,3} · Nicholas P. West⁴ · Nicolin Tee^{1,5} · Shona L. Halson^{1,6} · Eric J. Drinkwater^{2,7} · Dale W. Chapman^{1,2,8} · G. Gregory Haff^{2,9}

Received: 12 May 2023 / Accepted: 25 January 2024
© The Author(s) 2024

Abstract

Purpose Uncertainty exists if post-resistance exercise hydrotherapy attenuates chronic inflammatory and hormone responses. The effects of repeated post-resistance exercise water immersion on inflammatory and hormone responses in athletes were investigated.

Methods Male, academy Super Rugby players ($n = 18$, 19.9 ± 1.5 y, 1.85 ± 0.06 m, 98.3 ± 10.7 kg) participated in a 12-week programme divided into 3×4 -week blocks of post-resistance exercise water immersion (either, no immersion control [CON]; cold [CWI]; or hot [HWI] water immersion), utilising a randomised cross-over pre-post design. Fasted, morning blood measures were collected prior to commencement of first intervention block, and every fourth week thereafter. Linear mixed-effects models were used to analyse main (treatment, time) and interaction effects.

Results Repeated CWI ($p = 0.025$, $g = 0.05$) and HWI ($p < 0.001$, $g = 0.62$) reduced creatine kinase (CK), compared to CON. HWI decreased ($p = 0.013$, $g = 0.59$) interleukin (IL)-1ra, compared to CON. HWI increased ($p < 0.001$ – 0.026 , $g = 0.06$ – 0.17) growth factors (PDGF-BB, IGF-1), compared to CON and CWI. CWI increased ($p = 0.004$, $g = 0.46$) heat shock protein-72 (HSP-72), compared to HWI.

Conclusion Post-resistance exercise CWI or HWI resulted in trivial and moderate reductions in CK, respectively, which may be partly due to hydrostatic effects of water immersion. Post-resistance exercise HWI moderately decreased IL-1ra, which may be associated with post-resistance exercise skeletal muscle inflammation influencing chronic resistance exercise adaptive responses. Following post-resistance exercise water immersion, CWI increased HSP-72 suggesting a thermoregulatory response indicating improved adaptive inflammatory responses to temperature changes, while HWI increased growth factors (PDGF-BB, IGF-1) indicating different systematic signalling pathway activation. Our data supports the continued use of post-resistance exercise water immersion recovery strategies of any temperature during in-season competition phases for improved inflammatory adaptive responses in athletes.

Keywords Strength training · Team sports · Hydrotherapy · Recovery · Inflammation

Communicated by Fabio fischetti.

✉ Barry G. Horgan
barry.horgan2@gmail.com

¹ Australian Institute of Sport (AIS), Australian Sports Commission, Bruce, ACT 2617, Australia

² School of Medical and Health Sciences, Edith Cowan University (ECU), Joondalup, WA, Australia

³ Brumbies Rugby, Bruce, ACT, Australia

⁴ School of Medical Science, Menzies Health Institute QLD, Griffith University, Gold coast, Queensland, Australia

⁵ Mary MacKillop Institute of Health Research, Australian Catholic University, Melbourne, VIC, Australia

⁶ School of Behavioural and Health Sciences, Australian Catholic University, Banyo, Queensland, Australia

⁷ Centre for Sport Research, School of Exercise & Nutrition Sciences, Deakin University, Geelong, VIC, Australia

⁸ Curtin University, Bentley, WA 6102, Australia

⁹ Directorate of Psychology and Sport, University of Salford, Greater Manchester, Salford, UK

Abbreviations

| | |
|------|--------------------------------|
| CK | Creatine kinase |
| CON | Control |
| CWI | Cold water immersion |
| HSP | Heat shock protein |
| HWI | Hot water immersion |
| IGF | Insulin-like growth factor |
| IL | Interleukin |
| PDGF | Platelet-derived growth factor |
| TNF | Tumour necrosis factor |

Introduction

Athletes engage in resistance exercise across a season, to maintain and develop muscular hypertrophy, strength and power. Each individual resistance exercise session induces a systemic signalling response, which accrues over time following multiple repeated exposures. Athletes also often complete post-resistance exercise recovery strategies, e.g., water immersion, to promote recovery between training sessions. The hydrostatic pressure and temperature effects of water immersion alter the acute post-exercise physiological responses (Petersen and Fyfe 2021), by reducing acute muscle damage and swelling in athlete (Horgan et al. 2022a, b, c). Over time, these acute recovery responses may accumulate to accrue (promote) or mediate (diminish) the chronic systemic inflammatory signalling responses to resistance exercise, (Cornish et al. 2020).

Cold water immersion (CWI) involves the immersion of the body, i.e., single limb, partial- or whole-body (to neck, excluding head), in water temperatures ranging between 8 and 20 °C for a duration range of 5 to 30 min (Moore et al. 2022). The use of CWI throughout a taper period during competition preparation resulted in unclear (15 °C × 15 min, whole-body) (Halson et al. 2014) or positive (15 °C × 15 min, whole-body; 10 °C × 10 min, partial-body) (Halson et al. 2014; Tavares et al. 2020) performance responses, which may be associated with attenuated increases in post-exercise inflammation. If the practice of applying post-resistance exercise CWI (8 °C × 20 min, single-limb; 10 °C × 10 min, partial-body) occurs repeatedly over time, this may result in the accumulation of multiple acute reductions in myofibrillar protein synthesis (Fuchs et al. 2020a, b) and the down-regulation of satellite cell activity and hypertrophy signalling pathways (Roberts et al. 2015). These findings have contributed to scepticism regarding repeated CWI application following resistance exercise in athletes.

Optimising anabolic hormonal status, i.e., testosterone, has also been shown to be favourable for increases in lean muscle mass and strength in healthy males (Bhasin et al. 2001). A single post-resistance exercise hot water immersion (HWI) session (39 °C × 15 min, whole-body) has been

shown to result in small acute increases in testosterone concentration in athletes, as compared to a contrast water immersion strategy (Horgan et al. 2022b(in press)). Acute responses to post-resistance exercise HWI application also resulted in increased perceived sleep quality, while reducing next day perceptions of fatigue in athletes (Horgan et al. 2022b(in press)). As such, HWI may offer practitioners an alternative post-resistance exercise strategy to promote positive adaptive responses in athletes. Physiologically, HWI (~ 44 °C × 15–20 min, partial-body) increases peripheral muscular blood flow, as well as skin, muscle and core temperature (Bonde-Petersen et al. 1992). As highly trained athletes are reported to immerse themselves in hot water temperatures of ~ 38–42 °C (Rodrigues et al. 2020; Méline et al. 2021), it remains unclear if this strategy will stimulate positive adaptation (Fuchs et al. 2020a, b; Stevens et al. 2020) when applied repeatedly over time. Hence, the cumulative effects of applying post-resistance exercise CWI versus HWI across multiple sessions on the medium- to long-term hormone responses in athletes warrants further investigation.

Compared to recreationally active subjects, highly trained athletes demonstrate different inflammatory responses to exercise (Damas et al. 2018). Increased training age and status is generally associated with lower exercise-induced muscle damage, e.g., creatine kinase (CK), myoglobin (Damas et al. 2018); adapted increases, e.g., interleukin (IL)-4, IL-6, IL-7, IL-8, IL-15, insulin-like growth factor (IGF)-1, heat shock protein (HSP)-72; or decreases, e.g., IL-1ra, IL-1b, (Dennis et al. 2004) tumour necrosis factor (TNF)-α in inflammatory cytokines and myokines (Cornish et al. 2020), growth factors (Kraemer et al. 2017), and temperature-dependent, muscle regulating HSP-72 (Fyfe et al. 2019). Research investigating repeated post-resistance exercise water immersion in athletes is ambiguous, with few studies highlighting the discrepancies between responses in trained versus untrained subjects (Fyfe et al. 2019). Many studies utilise ‘recreationally active’ subjects, from which results cannot be directly translated to athletes. In studies which utilised highly trained athletes, CWI application occurred following sport-specific training and not following resistance exercise (Ihsan et al. 2021).

To date researchers have not investigated the cumulative effects of repeated post-resistance exercise water immersion on the adaptive inflammatory responses in a highly trained athlete cohort (Roberts et al. 2015). Studies utilising athletes in high performance training settings which investigate the underlying inflammatory adaptive responses to repeated post-resistance exercise water immersion are essential to advance the understanding of the role of post-resistance exercise water immersion, and how it can be periodised across training and competition. This study aimed to determine the cumulative effects following a repeated

post-resistance exercise (i.e., 4 weeks @ 2/week) CWI (15 °C × 15 min, whole-body) versus HWI (39 °C × 15 min, whole-body) strategy, compared to CON, on the systemic (blood) inflammatory and hormone responses following a 4-week intervention. We hypothesised that following repeated bouts of post-resistance exercise water immersion that: chronic hormone (testosterone, and testosterone to cortisol ratio) and growth factor (IGF-1, IGFBP-3 and PDGF-BB) responses would increase following all water immersion strategies (i.e., CWI or HWI); post-resistance exercise CWI would attenuate, and HWI would promote increases in blood markers of muscle damage (CK, myoglobin), pro-inflammatory cytokines (IL-1b, IL-6, IL-8, IL-17a and TNF- α) and heat-shock protein (HSP-72); and post-resistance exercise CWI would promote increases in anti-inflammatory cytokines (IL-1ra, IL-4 and IL-10), compared to HWI.

Methods

Subjects

Using a repeated measure study design, with conservative inputs of an effect size $f = 0.15$, $\alpha = 0.1$, 3 groups (CWI, HWI, CON) with an estimated total sample of 24 participants with 3 measures (time) and a correlation among repeated measures of 0.85, the achieved beta (power) is 0.823 (G*Power 3.1.9.2 software). This estimated sample accounts for additional participants to improve the confidence of correlations and to account for the risk of participant drop-out. Thirty-one sub-elite, male, academy Super Rugby players participated. Upon study commencement, subjects were participating in their club adult rugby in-season competition schedule, while also volunteering their participation in this study. In addition to Super Rugby academy and club rugby training, subjects were required to have completed a minimum twice weekly resistance training for > 12 months prior to study commencement. Study approval was obtained from Edith Cowan University and the Australian Institute of Sport (AIS) Human Research Ethics Committees (Approval#:17049). Clinical trial registration was obtained (ANZCTR#:12617000366358), with procedures adhering to the Declaration of Helsinki (2013). Subjects were screened for health and cardiovascular risk using a pre-exercise questionnaire (Adult Pre-exercise Screening System, ESSA, Australia). Subjects were also surveyed regarding any medication or nutritional supplements they were consuming by a qualified sports dietician, prior to confirming suitability to participate in the study. Subjects were informed of study risks and requirements prior to volunteering written informed consent. Prior to study commencement, subjects

were familiarised with exercise procedures, post-exercise treatment interventions and measurement test protocols. Subjects were required to complete a minimum of 87.5% (i.e., 7 of 8) of resistance exercise sessions and post-exercise intervention strategies to meet the inclusion criteria for each 4-week block. Subjects were excluded due to injuries sustained during their rugby training or games. Data for eighteen subjects ($n = 18$, age: 19.9 ± 1.5 y, height: 1.85 ± 0.06 m, weight: 98.3 ± 10.7 kg, body-fat: $18.7 \pm 4.1\%$, body mass index: 28.4 ± 2.5 kg·m², squat jump: 43.1 ± 7.5 cm, counter-movement jump: 45.4 ± 8.4 cm; Yo-Yo Intermittent Recovery Test Level 1: 1560 ± 440 m) were included in the analysis, distributed across CON ($n = 17$), CWI ($n = 17$) and HWI ($n = 15$) interventions.

Study design

A randomised controlled cross-over pre-post design was used, in which subjects undertook a 12-week (4-week × 3-intervention) programme (refer Figs. 1, 2, (Horgan et al. 2022)). Participants undertook twice weekly resistance exercise sessions, immediately followed by the post-resistance exercise treatment intervention (i.e., CON, CWI, HWI) sessions were performed across each 4-week block (refer Tables 1, 2, (Horgan, et al. 2022a)), for a total of eight resistance exercise and post-resistance exercise treatment intervention sessions each per 4-week block. Resistance exercise was individualised using percentage of body-mass, exercise load-movement velocity, or one repetition maximum. Sessions commenced at the same time of day (i.e., evening), were supervised and performed at room temperature (23–25 °C), with treatment interventions commencing 15-min post-exercise. Training load was monitored using session rating of perceived exertion (sRPE) (McGuigan and Foster 2004). Subjects consumed whey protein isolate (BodyScience, Australia) post-exercise, providing 25 g of protein, 3.2 g of leucine and 5.8 g of branched-chain amino acids. Subjects remained in the same post-resistance exercise intervention strategy, i.e., 15-min CON, CWI (15.1 ± 0.1 °C) or HWI (39.3 ± 0.6 °C) for each 4-week block and crossed-over thereafter. The CON condition performed static stretching at room temperature (23.0 ± 0.6 °C). Subjects in the water immersion strategies sat in the pool, immersed to the neck. Pool water was circulated, with temperatures monitored using a thermometer (Testo AG, Germany). Subjects were advised to avoid showering or bathing for > 2-h after all treatment conditions. Measurement testing occurred following a rest day, immediately prior to commencing the intervention (pre-test) and at the end of each treatment block (i.e., at the end of week 4, 8 and 12), in the morning prior to the subjects' weekly competitive game. At each testing time-point, fasting, rested morning blood samples were collected,

Fig. 1 Study design overview





Fig. 2 Training intervention schedule overview. Resistance exercise day 1 (Table 1), and day 2 (Table 2), was performed each Monday, and Wednesday respectively, prior to the post-exercise treatment intervention strategy (i.e., CON, CWI or HWI)



which occurred 36-h following the completion of the last exercise session.

Blood measures

Blood samples were collected via forearm venepuncture to assess systemic blood-based biomarkers (i.e., CK,

myoglobin, IL-1ra, IL-1b, IL-4, IL-6, IL-8, IL-10, IL-17a, TNF-α, IGF-1, IGFBP-3, PDGF-BB, HSP-72, testosterone, cortisol) in response to the exercise and intervention. Samples were prepared for cryo-storage and duplicate samples were analysed in the laboratory using methods as previously described (Horgan et al. 2022a, b, c), except for CK. In-season pre-game CK values of 832 ± 812 AU have been reported

Table 1 Description of resistance exercise session 1 completed prior to each post-exercise intervention strategy

| Resistance exercise day 1 | | Week 1 | Week 2 | Week 3 | Week 4 | Rest |
|---|-------|-------------|----------------|----------------|----------------|---------------------|
| A. Warm up. LB: SL Squat, OH Lateral leg raise, SL Hip extension; UB: YTW's, External shoulder raises, SA Push up | Reps | 2×10 | 2×10 | 2×10 | 2×10 | n/a |
| | % 1RM | n/a | n/a | n/a | n/a | |
| B1. BB Back squat | Reps | 4×8 | 8, 8, 6, 5 | 6, 6, 5, 3 | 5, 5, 3, 2 | On 4 min turnaround |
| | % 1RM | 66 | 66, 66, 72, 77 | 72, 72, 77, 82 | 77, 77, 82, 87 | |
| B2. Roman chair prone SL Isometric hold | Reps | 3×30 s | 3×30 s | 3×30 s | 3×30 s | |
| | % 1RM | n/a | n/a | n/a | n/a | |
| C1. BB rack pull | Reps | 3×6 | 3×6 | 3×5 | 3×4 | On 4 min turnaround |
| | % 1RM | 65 | 70 | 75 | 80 | |
| C2. Nordic curl | Reps | 3×4 | 3×4 | 3×4 | 3×4 | |
| | % 1RM | n/a | n/a | n/a | n/a | |
| D1. BB Bench press | Reps | 4×8 | 8, 8, 6, 5 | 6, 6, 5, 3 | 5, 5, 3, 2 | On 4 min turnaround |
| | % 1RM | 66 | 66, 66, 72, 77 | 72, 72, 77, 82 | 77, 77, 82, 87 | |
| D2. Pull up (prone grip) | Reps | 4×8 | 8, 8, 6, 5 | 6, 6, 5, 3 | 5, 5, 3, 2 | |
| | % 1RM | 66 | 66, 66, 72, 77 | 72, 72, 77, 82 | 77, 77, 82, 87 | |
| E1. DB SA Shoulder press | Reps | 3×10 | 3×8 | 3×6 | 3×5 | On 4 min turnaround |
| | % 1RM | 65 | 70 | 75 | 80 | |
| E2. SA DB Row | Reps | 3×10 | 3×8 | 3×6 | 3×5 | |
| | % 1RM | 65 | 70 | 75 | 80 | |
| F. Core. 60 s Front plank, 30 s es Side bridge, 60 s Prone bridge | Reps | 3 sets each | 3 sets each | 3 sets each | 3 sets each | 15 s between sets |
| | % 1RM | n/a | n/a | n/a | n/a | |

LB Lower-body, UB Upper-body, OH Overhead, SA Single-arm, SL Single-leg, Reps Repetitions, %1RM Percent of 1 repetition maximum, BB Barbell, DB Dumbbell, s Seconds, es Each side, min Minute, n/a Non-applicable, 2×10: Indicates 2 sets×10 repetitions were completed; 8,8,6,5: Indicates that a set of 8, 8, 6 and 5 repetitions were performed i.e. a total of 4 sets

in high-performance university rugby players and were used as an upper limit (i.e., mean + SD) reference (Mashiko et al. 2004). Inter- and intra-assay CV% were calculated for CK (1.1%, N/A), myoglobin (3.4 ± 3.6 , $6.6 \pm 6.7\%$), HSP-72 (5.2 ± 5.1 , $8.3 \pm 7.1\%$), IL-1ra (4.0 ± 3.8 , $9.9 \pm 10.5\%$), IL-4 (5.1 ± 4.4 , $9.3 \pm 10.1\%$), IL-10 (4.6 ± 4.2 , $8.9 \pm 11.3\%$), IL-1b (5.8 ± 5.1 , $8.2 \pm 8.4\%$), IL-6 (4.3 ± 3.6 , $8.5 \pm 12.2\%$), IL-8 (3.4 ± 3.6 , $6.6 \pm 6.7\%$), IL-17a (6.2 ± 4.4 , $8.5 \pm 10.7\%$), TNF- α (6.5 , 2.7%), IGF-1 (5.3%, N/A), IGFBP-3 (4.8 ± 4.0 , $11.7 \pm 19.6\%$), PDGF-BB (8.0 ± 4.7 , $12.2 \pm 12.1\%$), testosterone (14.0, 12.0%) and cortisol (9.0, 4.0%). The dynamic range for cortisol and testosterone and is 1000–3,600,000 pg/mL and 0.7–875 nM, respectively.

Load monitoring

sRPE was surveyed for all team training, competition, and resistance exercise sessions (refer Table 3, (Horgan et al. 2022)). Five-day food diaries were collected. Due to challenges with the accuracy of reporting food diary data (Burke 2015), dietician and statistician advice received advised against analysing the diaries due to missing data-points and risk of bias associated with this data collection method. Thermal sensation (TS) (refer Fig. 3, (Horgan et al. 2022)) was assessed for each post-exercise intervention strategy

using a 0–8 Likert scale (0=“unbearably cold”; 8=“unbearably hot”) immediately prior (0-min), at one minute (1-min), mid-way (7.5-min) and on completion (15-min) of each strategy.

Statistical analyses

Descriptive statistics were calculated prior to analysis using software (Stata/IC 15.1, StataCorp LP, USA). Missing data did not show any obvious pattern and were deemed to be at random and no imputation was made. Data were assessed visually via Q-Q plots, as well as using Shapiro–Wilk’s normality test. Outliers were assessed and identified as ‘severe’ if they were ± 3 SD below or above the 25th or 75th quartiles and the authors searched for any obvious reason to exclude these data. Data were analysed using raw values. Linear mixed-effects models (LMM) were conducted to analyse main (treatment, time) and interaction (treatment×time) effects. Subject were set as a random intercept and the model was fitted via maximum likelihood estimation using a 1×1 covariance matrix structure. The model analysed treatment intervention (i.e., CON, CWI, HWI) and time (group order, i.e., block A, B, C) as fixed effects on the dependent variable,

Table 2 Description of resistance exercise session 2 completed prior to each post-exercise intervention strategy

| Resistance exercise day 2 | | Week 1 | Week 2 | Week 3 | Week 4 | Rest |
|---|-------|-------------|-------------|----------------|----------------|---------------------|
| A. Warm up LB: SL Squat, OH Lateral leg raise, SL Hip extension; UB: YTW's, External shoulder raise and prone press | Reps | 2×10 | 2×10 | 2×10 | 2×10 | n/a |
| | % 1RM | n/a | n/a | n/a | n/a | |
| B. 15-min run drilling and accelerations (15-min). 60 s Skipping, A-march, A-skip, A-drill, Accelerations | Reps | 3×20 m | 3×20 m | 3×20 m | 3×20 m | On 4 min turnaround |
| | % 1RM | n/a | n/a | n/a | n/a | |
| C1. Jump monitoring (SJ, and CMJ) | Reps | 2×5 | 2×5 | 2×5 | 2×5 | |
| | % 1RM | n/a | n/a | n/a | n/a | |
| C2. BB Jump Squat | Reps | 4×5 | 4×5 | 5,5,4,3 | 4,4,3,2 | On 4 min turnaround |
| | % 1RM | 65 | 70 | 70, 70, 75, 80 | 75, 75, 80, 85 | |
| D1. BB Power shrug (floor) | Reps | 4×5 | 4×5 | 5,5,4,3 | 4,4,3,2 | |
| | % 1RM | 65 | 70 | 70, 70, 75, 80 | 75, 75, 80, 85 | |
| D2. BB 2-step Step up to Hip-lock | Reps | 4×4 es | 4×4 es | 4×4 es | 4×4 es | On 4 min turnaround |
| | % 1RM | 30 kg | 30 kg | 30 kg | 30 kg | |
| E1. 3-way DB Shoulder raise | Reps | 3×6 | 3×5 | 3×4 | 3×3 | |
| | % 1RM | 65 | 70 | 75 | 80 | |
| E2. DB Alternating bench Press | Reps | 3×10 | 3×8 | 3×6 | 3×5 | On 4 min turnaround |
| | % 1RM | 65 | 70 | 75 | 80 | |
| E3. Seated row | Reps | 3×10 | 3×8 | 3×6 | 3×5 | |
| | % 1RM | 65 | 70 | 75 | 80 | |
| F. Core. 12×Hanging leg raise, 20×BB Roll-out, 10 es x Lateral pallof press | Reps | 3 sets each | 3 sets each | 3 sets each | 3 sets each | 15 s between sets |
| | % 1RM | n/a | n/a | n/a | n/a | |

LB Lower-body, UB Upper-body, OH Overhead, SA Single-arm, SL: Single-leg, Reps Repetitions, % 1RM Percent of 1 repetition maximum, BB Barbell, DB: Dumbbell, es Each side, s Seconds, min Minute, N/A Non-applicable, 2×10 Indicates 2 sets×10 repetitions were completed; 5, 5, 4, 3: Indicates that a set of 5, 5, 4 and 3 repetitions were performed i.e. a total of 4 sets

Table 3 Descriptive statistics for measures describing the resistance exercise sessions, total weekly training load and post-exercise intervention strategies

| Training load variable | CON (N=17) | CWI (N=17) | HWI (N=15) |
|------------------------------------|--------------|--------------|--------------|
| Resistance exercise session 1 sRPE | 316.1±79.5 | 321.5±101.2 | 345.5±95.9 |
| Resistance exercise session 2 sRPE | 327.5±86.5 | 324.8±107.6 | 314.9±82.4 |
| Total weekly sRPE | 1574.4±523.7 | 1591.5±607.9 | 1789.8±528.9 |

sRPE Session rating of perceived exertion

resulting in nine combinations of treatment and time, with two subjects in each such combination. Significance was set at $p \leq 0.05$. Following LMM, main and interaction effects were evaluated using paired samples t-test and corrected using Scheffé's method of multiple comparison to control for Type 1 error. Intra-class correlations were computed following LMM. Model fit assumptions were assessed via the calculation of residuals. As body composition influences the magnitude of muscle temperature change (Stephens et al. 2017), subject percent of body-fat (BF%) was used as a fixed effect covariate for response variables. Where significant differences were observed between interventions, practical differences in means were interpreted using Hedges g (1981) effect size (ES) statistic,

where $g < 0.2$ = "trivial", $0.2 \leq g < 0.5$ = "small", $0.50 \leq g < 0.8$ = "medium" and $g \geq 0.8$ = "large".

Results

The treatment groups were balanced for total weekly in-season training load across all rugby and resistance exercise sessions (refer Table 3, (Horgan et al. 2022)). The post-exercise water immersion resulted in significant (time, treatment, and time×treatment interaction effects) changes in TS (refer Fig. 3, (Horgan et al. 2022)), compared to CON.




Fig. 3 Mean \pm SD thermal sensation response over time (min post-immersion) pre- (0 min) and per-immersion (1-, 7.5- and 15-min post-immersion) intervention strategy following whole body resistance exercise with either CON (line with unfilled circle $n=17$), CWI (hyphen with filled square $n=17$), or HWI (hyphen with filled circle $n=15$) intervention. Significant main effects were observed for treatment ($p < 0.001$), time ($p < 0.001$), and treatment \times time interaction effect ($p < 0.001$) with Tukey-adjusted p -values reporting differences for individual treatment \times time pair-wise comparisons at 1-, 7.5- and 15-min following CWI, and HWI, compared to CON condition; ***: $p < 0.001$ versus CON at pre-immersion baseline; ###: $p < 0.001$ versus CON within time-point

Blood measures

Muscle damage

Changes in serum CK concentration are shown in Fig. 4 (Panels A, B). Significant reductions in Δ CK (Fig. 4B) occurred following CWI ($p = 0.025$, $g = 0.05$) and HWI ($p < 0.001$, $g = 0.62$), compared to CON. Myoglobin concentration did not differ significantly between interventions (data not shown).

Heat shock protein

The effect of post-exercise intervention strategies on serum HSP-72 concentration is shown in Fig. 4 (panels C, D). CWI resulted in significant ($p = 0.004$, $g = 0.46$) increases in HSP-72, compared to HWI.

Anti-inflammatory cytokines

Figure 4 (panels E, F) shows the effect of post-exercise intervention strategies on the concentration of serum IL-1ra. CON was associated with significant ($p = 0.013$, $g = 0.59$) increases in IL-1ra, compared to HWI. IL-4 and IL-10 concentrations did not differ significantly between interventions (data not shown).

Pro-inflammatory cytokines

Following resistance exercise, the post-exercise intervention strategies did not influence ($p > 0.05$) the concentration of serum pro-inflammatory cytokine concentrations (i.e., IL-1b, IL-6, IL-8, IL-17a, TNF- α ; data not shown). Significant positive (IL-8: $p = 0.001$, $r = 0.22$; IL-17a: $p = 0.029$, $r = 0.26$) and negative (TNF- α : $p < 0.001$, $r = 0.37$) associations were observed with the %BF covariate.

Growth factors

Figure 5 shows the effect of post-exercise intervention strategies on serum growth factor concentrations. IGF-1 (Fig. 5, panels A, B) was significantly lower following CON ($p = 0.001$, $g = 0.06$) and CWI ($p < 0.001$, $g = 0.11$), compared to HWI. PDGF-BB (Fig. 5, panels C, D) was significantly lower following CON ($p = 0.026$, $g = 0.17$), compared to HWI. IGFBP-3 concentration did not differ significantly between interventions (data not shown).

Hormones

Figure 6 shows the effect of post-exercise intervention strategies on serum hormone concentrations.


Testosterone, cortisol, and testosterone to cortisol ratio (TC ratio) concentrations did not differ significantly between interventions (Fig. 6).

Discussion

During an in-season phase, this study investigated the effects of repeated post-resistance exercise CWI, HWI or CON on inflammatory responses in highly trained athletes. We found that repeated post-resistance exercise HWI resulted in moderate reductions in anti-inflammatory cytokine IL-1ra ($p = 0.013$), and muscle damage, i.e., CK ($p < 0.001$); while, HWI also resulted in trivial increases ($p < 0.001$ – 0.026) in growth factors PDGF-BB and IGF-1, as compared to CON and CWI. Conversely, we found that repeated post-resistance exercise CWI resulted in trivial reductions ($p = 0.025$) in muscle damage, i.e., CK, and small increases ($p = 0.004$) in heat shock protein HSP-72, as compared to HWI. This is one of the first studies to investigate the effects of repeated post-resistance exercise water immersion on an extensive array of inflammatory measures in highly trained athletes.

This study's findings for muscle damage (i.e., CK) following repeated post-resistance exercise water immersion are consistent with previous research investigating acute responses following a single immersion (Horgan et al. 2022a, b, c), where irrespective of temperature (i.e., cold vs. hot) post-exercise water immersion resulted in

Fig. 4 Mean ($\pm 95\%$ C.I.) muscle damage, heat shock protein and anti-inflammatory response pre- (white bar) versus post-intervention (grey bar), and mean ($\pm 95\%$ C.I.) pre-minus post-intervention delta (Δ) (black bar, and individual data points) for CK (panels A, B), HSP-72 (panels C, D) and IL-1ra (panels E, F) following whole body resistance exercise with either control (CON, $n=17$), cold (CWI, $n=17$) or hot (HWI, $n=15$) water immersion as treatment. Significant main effects for treatment were observed in CK (where CON > CWI, $p=0.025$, $g=0.05$; and CON > HWI, $p<0.001$, $g=0.62$) and HSP-72 (where CWI > HWI, $p=0.004$, $g=0.46$); Significant main effects for treatment were observed in IL-1ra (where CON > HWI, $p=0.013$, $g=0.59$); *: $p\leq 0.05$, **: $p\leq 0.01$, ***: $p\leq 0.001$



trivial-to-moderate reductions in CK, compared to CON. Research to date has predominantly focused on acute effects of post-exercise water immersion. As such, this study is one of the first to investigate the effects of repeated post-resistance exercise water immersion on exercise-induced muscle damage using an all-athlete cohort (Tavares et al. 2018; Méline et al. 2021). The contribution from muscle damage

to long-term hypertrophic adaptive responses to resistance exercise is believed to reduce in proportion with increased training status (Damas et al. 2018). Therefore, muscle damage in trained athletes can be considered as non-desirable symptomatology associated with resistance exercise. These findings are favourable in terms of the proposed hydrostatic pressure and temperature effects of post-exercise water




Fig. 5 Mean ($\pm 95\%$ C.I.) growth factor responses pre- (white bar) versus post-intervention (grey bar), and mean ($\pm 95\%$ C.I.) pre- minus post-intervention delta (Δ) (black bar, and individual data points) for IGF-1 (panels **A**, **B**) and PDGF-BB (panels **C**, **D**) following whole body resistance exercise with either control (CON, n=17),


cold (CWI, n=17) or hot (HWI, n=15) water immersion as treatment. Significant main effects for treatment were observed in IGF-1 (where HWI>CON, $p=0.001$, $g=0.06$; and HWI>CWI, $p<0.001$, $g=0.11$) and PDGF-BB (where HWI>CON, $p=0.026$, $g=0.17$); *: $p\leq 0.05$, ***: $p\leq 0.001$

immersion, which reduces muscle damage biomarkers following resistance exercise. This study observed a large variation in CK, which may be partly explained by other non-modifiable factors, e.g., genetics, ethnicity, gender or age (Scalco et al. 2016). A limitation of the study is the single time-point for blood measures at the end of each 4-week block, given differences in plasma kinetics between biomarkers (Scalco et al. 2016). Future studies may consider

examining the longer-term (i.e., > 4-week) pre- versus post-resistance exercise and intervention response of these markers.

Chronic, resting, serum IL-1ra anti-inflammatory cytokine levels were moderately lower ($p=0.013$) following post-exercise HWI, compared to CON. The chronic effects of resistance exercise on serum cytokines in highly trained athletes remains uncertain. Anti-inflammatory cytokines

Fig. 6 Mean and individual (open circles) hormone responses pre- (white bar) versus post-intervention (grey bar), and mean (\pm 95% C.I.) pre- minus post-intervention delta (black bar) for testosterone (panel A, B), cortisol (panel C, D) and TC ratio (panel E, F) following whole body resistance exercise with either control (CON, $n=17$), cold (CWI, $n=17$) or hot (HWI, $n=15$) water immersion as treatment. Significant main effects for treatment were observed in testosterone (where $\text{HWI} > \text{CON}$, $p=0.012$); *, $p \leq 0.05$, **, $p \leq 0.01$, ***, $p \leq 0.001$, ****, $p \leq 0.0001$



are thought to play a key role in the inflammatory cascade initiated by damaged muscle from resistance exercise (Methenitis et al. 2021). In our study, the CON group displayed significantly higher muscle damage (i.e., CK), which may be a reason for the elevated IL-1ra when blood testing occurred \sim 36-h after the last session. Similarly, elevated core temperature (\sim 39.5 °C) in mice has previously been associated with increased circulating IL-1 levels,

providing accelerated recovery and adaptive responses (Dinarelli 2012). It should be noted that while there may be some similarities between mice and humans in terms of thermoregulation, there equally are as many challenges in directly comparing both due to the unique thermoregulatory systems of both species (Gordon 2017). For example, humans can take up to 115 min for the rectal temperature to reach 38.8 ± 0.14 °C while sitting passively in HWI at 42 °C

(Rodrigues et al. 2020). Future research utilising athletes should investigate the systemic effects that post-exercise water immersion has on acute and chronic responses of genetic and intracellular RNA signalling using closed tube technology, which isolates intracellular RNA, as well as on immune cell subsets using mass cytometry.

Pro-inflammatory cytokines were either positively (i.e., IL-8, IL-17a) or negatively (i.e., TNF- α) associated with percent of body-fat, which was utilised as a covariate in the statistical analysis. This is consistent with findings by Aguiar et al. (2020), where adiposity was positively correlated with pro-inflammatory cytokines in master athletes (Aguiar et al. 2020), with the unexpected exception of TNF- α . A limitation of this study was that we did not provide complete dietary standardisation across the study, with dietary supplementation shown to reduce eccentric muscle damage (Córdova-Martínez et al. 2021) or inflammation markers (e.g., TNF- α) in athletes (Willis et al. 2012). However, our randomised cross-over study design may reduce the effect of this limitation. Another consideration related to nutrition as a covariate for adaptive responses is whether the temperature effects of post-exercise water immersion may influence appetite. Future research should investigate the effects the temperature of post-resistance exercise water immersion may have on markers of appetite regulation.

Post-resistance exercise HWI resulted in trivial increases in IGF-1, compared to CWI or CON strategies, while post-resistance exercise HWI resulted in trivial increases in PDGF-BB, compared to CON. The authors are not aware of any research investigating post-resistance exercise CWI versus HWI in athletes on chronic systemic growth factor responses. Synergistic effects of IGF-1 and PDGF-BB have been reported to provide therapeutic anti-inflammatory osteoarthritic benefits (Montaseri et al. 2011). Collectively, the above results suggest CWI may lower, albeit with trivial effect ($g = 0.06\text{--}0.11$), systemic circulating anabolic growth factors, i.e., IGF-1. Our finding is consistent with observations that post-resistance exercise CWI may attenuate the anabolic response in resistance trained males (Earp et al. 2019). In summary, synergistic effects in the systemic circulation of anabolic growth factors (i.e., IGF-1, PDGF-BB) accrue following repeated post-resistance exercise HWI application in athletes, and may contribute towards promoting an anabolic response in athletes, similar to previous studies utilising various forms of heating in combination with exercise stress to enhance adaptation and attenuate atrophy in human and rodent subjects (McGorm et al. 2018; McGorm 2019).

The immersion protocols in the current study reduced ($\sim 47\%$) or increased ($\sim 57\%$) thermal sensation following CWI or HWI, respectively. Our novel study utilising an all athlete participant cohort is the first to demonstrate small increases occurred in thermo-sensitive serum HSP-72

following repeated CWI, compared to HWI. Our findings are in contrast to evidence demonstrating no change in intramuscular HSP-72 following four weeks of repeated post-endurance exercise CWI (7-week @ 3-day/week, 15 min \times 10 °C) following high-intensity interval cycling exercise in physically active participants (Aguiar et al. 2016), as well as evidence demonstrating that repeated post-resistance exercise CWI (4-week @ 3-day/week, 15 min \times 10 °C) attenuated total intramuscular HSP-72 in recreationally active participants (Fyfe et al. 2019). This unexpected chronic increase in serum HSP-72 following CWI may be explained via a thermal shock-like response due to relative or absolute temperature changes following cooling and upon the homeostatic return to normothermia, i.e., re-warming following a cold shock (Neutelings et al. 2013). Similarly, HSP function and roles differ depending on the tissue of origin (i.e., intracellular versus extracellular), with environmental factors, and exercise mode, duration and intensity all influencing availability (Yamada et al. 2008). It is plausible that our findings are specific to extracellular serum HSP availability following post-resistance exercise CWI in athletes. Given the importance of HSPs in regulating exercise-induced muscular adaptation, this outcome may also partially explain our finding that post-resistance exercise CWI did not attenuate lean muscle mass in athletes. Similarly, reduced resting levels of serum HSP-72 have also been suggested as evidence of an improved inflammatory profile (Hoekstra et al. 2018), as well as evidence of improved heat adaptation via reduced HSP production in response to thermal stressors, e.g., HWI (Gibson et al. 2017).

Practical applications

Post-resistance exercise water immersion strategies, such as CWI, HWI, etc., may be used to provide trivial-to-moderate reductions in muscle damage (i.e., CK) following a 4-week in-season schedule in highly trained athletes. Water immersion strategies as utilised in this study, may be prescribed for highly trained athletes during an in-season competition phase, posing little risk of attenuating inflammatory adaptive responses. These results support the notion that water immersion may be periodised, i.e., the use of hot increases systemic growth factor circulation (i.e., PDGF-BB, IGF-1), while the use of cold increases thermo-sensitive systemic heat shock protein (i.e., HSP-72) circulation. Practitioners working with athletes should consider individual responses, e.g., as evident by mean \pm 95% confidence interval ranges, to post-resistance exercise recovery strategies to identify each athlete's 'adaptive sweet spot'. Future study designs should consider the differences in inflammatory responses

to exercise and hydrotherapy strategies for trained athletes versus recreationally active, non-athletes.

Conclusions

This investigation provides insights into systemic inflammatory interactions and responses following post-resistance exercise water immersion. This study provides evidence to inform post-resistance exercise water immersion prescription, i.e., timing and temperature, across a 4-week in-season competition phase for highly trained athletes. Our study, which utilised an athlete cohort, supports the theory of potential differences between athletic and recreationally active participants, which may be partly explained by understanding differing muscle protein synthesis and hypertrophy signalling pathways in trained versus untrained subjects (Damas et al. 2018; Cornish et al. 2020). As athletes become accustomed to resistance exercise, less exercise-induced muscle damage is experienced suggesting that muscle damage is not the primary mechanistic driver for muscular adaptation in athletes (Rossi et al. 2018). It is likely that the synergistic effects of multiple inflammatory responses following resistance exercise trigger both muscle damage and other pro- and anti-inflammatory cytokine, hormone, growth factor, heat shock protein and haematological responses which may cumulatively mediate the downstream muscle protein synthesis and hypertrophy signalling pathways. Future studies should investigate the effect of repeated post-resistance exercise water immersion on other mechanistic adaptive responses in athletes, such as changes in muscle architecture and electromyography.

Acknowledgements The authors would like to acknowledge that this research and work was carried out on the Traditional Lands of the Ngunnawal people, and pay our respect to Elders both past, present, and future. We would like to thank the subjects, as well as their strength and conditioning coach, Benjamin Norcott, for their time, dedication, and efforts during the study. The authors wish to thank the Systems Biology and Data Science Group at Griffith University for their assistance with analysing the immunology biomarkers. The assistance and support during data collection from Felicity Bright, Dr. James Broatch, Steven Hughes, Dr. Jessica Stephens, and Alice Wallett is also greatly acknowledged. The authors would also like to thank Dr. Marijke Welvaert and Dr. James Gilmore for their statistical advice. Finally, we would like to acknowledge the University of Canberra researcher development team for their assistance and support throughout this study.

Author contributions All authors conceived and designed the research and conducted the experiments. BH and NT processed the data, which was statistically analysed by BH. All authors reviewed the statistical analysis and results. BH prepared the manuscript. All authors read and approved the manuscript.

Funding Open Access funding enabled and organized by CAUL and its Member Institutions. Funding for this study was obtained via research grants from the Australian Institute of Sport (AIS) High Performance Sport Research Funds (HPSRF) awarded to BH., GH., DC., SH. and

ED., Deakin University awarded to ED., Griffith University awarded to NW., and Edith Cowan University (ECU) awarded to GH. and DC. BH. was also in receipt of an industry engagement post-graduate research scholarship from AIS and ECU. Deakin University, Australian Institute of Sport, Griffith University, Edith Cowan University

Data availability The datasets generated and analysed during the current study are not publicly available due to sensitivity associated with high-performance athlete data, however, are available from the corresponding author on reasonable request.

Declarations

Conflict of interests The authors have no competing interests to declare that are relevant to the content of this article.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Aguiar PF, Magalhães SM, Fonseca IAT, da Costa Santos VB, de Matos MA, Peixoto MFD, Nakamura FY, Crandall C, Araújo HN, Silveira LR (2016) Post-exercise cold water immersion does not alter high intensity interval training-induced exercise performance and Hsp72 responses, but enhances mitochondrial markers. *Cell Stress Chaperones* 21(5):1–12
- Aguiar SS, Sousa CV, Deus LA, Rosa TS, Sales MM, Neves RVP, Barbosa LP, Santos PA, Campbell CS, Simões HG (2020) Oxidative stress, inflammatory cytokines and body composition of master athletes: the interplay. *Exp Gerontol* 130:110806
- Bhasin S, Woodhouse L, Casaburi R, Singh AB, Bhasin D, Berman N, Chen X, Yarasheski KE, Magliano L, Dzau C (2001) Testosterone dose-response relationships in healthy young men. *Am J Physiol Endocrinol Metab* 281(6):E1172–1181
- Bonde-Petersen F, Schultz-Pedersen L, Dragsted N (1992) Peripheral and central blood flow in man during cold, thermoneutral, and hot water immersion. *Aviat Space Environ Med* 63(5):346–350
- Burke L (2015) Dietary assessment methods for the athlete: Pros and cons of different methods. *Sports Sci Exch* 28(150):1–6
- Córdova-Martínez A, Caballero-García A, Bello HJ, Pérez-Valdecantos D, Roche E (2021) Effect of glutamine supplementation on muscular damage biomarkers in professional basketball players. *Nutrients* 13(6):2073
- Cornish SM, Bugera EM, Duhamel TA, Peeler JD, Anderson JE (2020) A focused review of myokines as a potential contributor to muscle hypertrophy from resistance-based exercise. *Eur J Appl Physiol* 120:941–959
- Damas F, Libardi CA, Ugrinowitsch C (2018) The development of skeletal muscle hypertrophy through resistance training: the role of muscle damage and muscle protein synthesis. *Eur J Appl Physiol* 118(3):485–500

- Dennis RA, Trappe TA, Simpson P, Carroll C, Emma Huang B, Nagarajan R, Bearden E, Gurley C, Duff GW, Evans WJ (2004) Interleukin-1 polymorphisms are associated with the inflammatory response in human muscle to acute resistance exercise. *J Physiol* 560(3):617–626
- Dinarello CA (2012) Keep up the heat on IL-1. *Blood* 120(13):2538–2539
- Earp JE, Hatfield DL, Sherman A, Lee EC, Kraemer WJ (2019) Cold-water immersion blunts and delays increases in circulating testosterone and cytokines post-resistance exercise. *Eur J Appl Physiol* 119(8):1901–1907
- Fuchs CJ, Kouw IW, Churchward-Venne TA, Smeets JS, Senden JM, van Marken Lichtenbelt WD, Verdijk LB, van Loon LJ (2020a) Post exercise cooling impairs muscle protein synthesis rates in recreational athletes. *J Physiol* 598(4):755–772
- Fuchs CJ, Smeets JS, Senden JM, Zorenc AH, Goessens JP, van Marken Lichtenbelt WD, Verdijk LB, van Loon LJ (2020b) Hot-water immersion does not increase postprandial muscle protein synthesis rates during recovery from resistance-type exercise in healthy, young males. *J Appl Physiol* 128(4):1012–1022
- Fyfe JJ, Broatch JR, Trewin AJ, Hanson ED, Argus CK, Garnham AP, Halson SL, Polman RC, Bishop DJ, Petersen AC (2019) Cold water immersion attenuates anabolic signalling and skeletal muscle fiber hypertrophy, but not strength gain, following whole-body resistance training. *J Appl Physiol* 127(5):1403–1418
- Gibson OR, Taylor L, Watt PW, Maxwell NS (2017) Cross-adaptation: heat and cold adaptation to improve physiological and cellular responses to hypoxia. *Sports Med* 47(9):1751–1768
- Gordon CJ (2017) The mouse thermoregulatory system: Its impact on translating biomedical data to humans. *Physiol Behav* 179:55–66
- Halson SL, Bartram J, West N, Stephens J, Argus CK, Driller MW, Sargent C, Lastella M, Hopkins WG, Martin DT (2014) Does hydrotherapy help or hinder adaptation to training in competitive cyclists? *Med Sci Sports Exerc* 46(8):1631–1639
- Hoekstra SP, Bishop NC, Faulkner SH, Bailey SJ, Leicht CA (2018) Acute and chronic effects of hot water immersion on inflammation and metabolism in sedentary, overweight adults. *J Appl Physiol* 125(6):2008–2018
- Horgan BG, Halson SL, Drinkwater EJ, West NP, Tee N, Alcock RD, Chapman DW, Haff GG (2022a) No effect of repeated post-resistance exercise cold or hot water immersion on in-season body composition and performance responses in academy rugby players: a randomised controlled cross-over design. *Eur J Appl Physiol* 123(2):353–359
- Horgan BG, Tee N, West NP, Drinkwater EJ, Halson SL, Colomer CM, Fonda CJ, Tatham J, Chapman DW, Haff GG (2022b) Acute performance, daily well-being and hormone responses to water immersion following resistance exercise in junior international and sub-elite male volleyball athletes. *J Strength Cond Res* 10:1519
- Horgan BG, West NP, Tee N, Drinkwater EJ, Halson SL, Vider J, Fonda CJ, Haff GG, Chapman DW (2022c) Acute inflammatory, anthropometric, and perceptual (muscle soreness) effects of postresistance exercise water immersion in junior international and subelite male volleyball athletes. *J Strength Cond Res* 36(12):3473–3484
- Ihsan M, Abbiss CR, Allan R (2021) Adaptations to post-exercise cold water immersion: friend, foe, or futile? *Front Sports Act Living* 3(201):714148
- Kraemer WJ, Ratamess NA, Nindl BC (2017) Recovery responses of testosterone, growth hormone, and IGF-1 after resistance exercise. *J Appl Physiol* 122(3):549–558
- Mashiko T, Umeda T, Nakaji S, Sugawara K (2004) Position related analysis of the appearance of and relationship between post-match physical and mental fatigue in university rugby football players. *Br J Sports Med* 38(5):617–621
- McGorm H (2019) The effects of hot water immersion on recovery, performance and adaptation to resistance exercise. University of Queensland, Doctor of Philosophy
- McGorm H, Roberts LA, Coombes JS, Peake JM (2018) Turning up the heat: an evaluation of the evidence for heating to promote exercise recovery, muscle rehabilitation and adaptation. *Sports Med* 48(6):1311–1328
- McGuigan MR, Foster C (2004) A new approach to monitoring resistance training. *Strength Cond J* 26(6):42
- Méline T, Solsona R, Antonietti J-P, Borrani F, Candau R, Sanchez AM (2021) Influence of post-exercise hot-water therapy on adaptations to training over 4 weeks in elite short-track speed skaters. *J Exerc Sci Fit* 19(2):134–142
- Methenitis S, Stergiou I, Antonopoulou S, Nomikos T (2021) Can exercise-induced muscle damage be a good model for the investigation of the anti-inflammatory properties of diet in humans? *Biomedicine* 9(1):36
- Montaseri A, Busch F, Mobasheri A, Buhrmann C, Aldinger C, Rad JS, Shakibaei M (2011) IGF-1 and PDGF-bb suppress IL-1 β -induced cartilage degradation through down-regulation of NF- κ B signaling: involvement of Src/PI-3K/AKT pathway. *PLoS One* 6(12):e28663
- Moore E, Fuller JT, Bellenger CR, Saunders S, Halson SL, Broatch JR, Buckley JD (2022) Effects of cold-water immersion compared with other recovery modalities on athletic performance following acute strenuous exercise in physically active participants: a systematic review, meta-analysis, and meta-regression. *Sports Med* 53(3):1–19
- Neutelings T, Lambert CA, Nussgens BV, Colige AC (2013) Effects of mild cold shock (25 °C) followed by warming up at 37 °C on the cellular stress response. *PLoS One* 8(7):e69687
- Petersen AC, Fyfe JJ (2021) Post-exercise cold water immersion effects on physiological adaptations to resistance training and the underlying mechanisms in skeletal muscle: a narrative review *Frontiers Sports and Active Living* 3:79
- Roberts LA, Raastad T, Markworth JF, Figueiredo VC, Egner IM, Shield A, Cameron-Smith D, Coombes JS, Peake JM (2015) Post-exercise cold water immersion attenuates acute anabolic signalling and long-term adaptations in muscle to strength training. *J Physiol* 593(18):4285–4301
- Rodrigues P, Trajano GS, Wharton L, Minett GM (2020) Muscle temperature kinetics and thermoregulatory responses to 42 °C hot-water immersion in healthy males and females. *Eur J Appl Physiol* 120(12):2611–2624
- Rossi FE, De Freitas MC, Zanchi NE, Lira FS, Cholewa JM (2018) The role of inflammation and immune cells in blood flow restriction training adaptation: a review. *Front Physiol* 9:1376
- Scalco RS, Snoeck M, Quinlivan R, Treves S, Laforêt P, Jungbluth H, Voermans NC (2016) Exertional rhabdomyolysis: physiological response or manifestation of an underlying myopathy? *BMJ Open Sport Exerc Med* 2(1):e000151
- Stephens JM, Halson SL, Miller J, Slater GJ, Chapman DW, Askew CD (2017) Effect of body composition on physiological responses to cold water immersion and the recovery of exercise performance. *Int J Sports Physiol Perform* 13(3):1–28
- Stevens CJ, Ross ML, Carr AJ, Vallance B, Best R, Urwin C, Périard JD, Burke L (2020) Post exercise hot-water immersion does not further enhance heat adaptation or performance in endurance athletes training in a hot environment. *Int J Sports Physiol Perform* 16(4):480–488
- Tavares F, Beaven M, Teles J, Baker D, Healey P, Smith TB, Driller M (2018) The effects of chronic cold water immersion in elite rugby players. *Int J Sports Physiol Perform* 14(2):1–23
- Tavares F, Simoes M, Matos B, Smith TB, Driller M (2020) The acute and longer-term effects of cold water immersion in highly-trained volleyball athletes during an intense training block. *Front Sports Act Living* 2:568420

- Willis KS, Smith DT, Broughton KS, Larson-Meyer DE (2012) Vitamin D status and biomarkers of inflammation in runners. *Open Access J Sports Med* 3:35
- Yamada P, Amorim F, Moseley P, Schneider S (2008) Heat shock protein 72 response to exercise in humans. *Sports Med* 38(9):715–733

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.