

Erratum to: Surgical aspects and complications of continuous intraperitoneal insulin infusion with an implantable pump

Jan Willem Haveman · Susan J. J. Logtenberg ·
Nanne Kleefstra · Klaas H. Groenier · Henk J. G. Bilo ·
Adri M. Blomme

Published online: 30 August 2012
© Springer-Verlag 2012

Erratum to: Langenbecks Arch Surg (2010) 395:65–71
DOI 10.1007/s00423-008-0437-9

The original version of this article by Haveman et al. unfortunately contained an error concerning the statistical analysis. The 95 % confidence interval (CI) reported in this publication was 26–130 months, not 13–79 months as initially reported.

The correct results are as follows:

We calculated the median operation free period between the first and second operation. This operation free period increased from 21 months (95 % CI: 10–33 months) for patients operated before 2000 to 78 months (95 % CI: 26–130 months) for patients operated in or after 2000 ($p=0.039$; log-rank test, see Fig. 2).

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00423-008-0437-9>.

J. W. Haveman (✉) · A. M. Blomme
Department of Surgery, Isala Clinics,
P.O. Box 10400, 8000 GK Zwolle, The Netherlands
e-mail: janwillemhaveman@gmail.com

S. J. J. Logtenberg · N. Kleefstra · H. J. G. Bilo
Diabetes Centre, Isala Clinics,
Zwolle, The Netherlands

N. Kleefstra
Langerhans Medical Research Group,
Zwolle, The Netherlands

K. H. Groenier
Department of General Practice,
University Medical Center Groningen,
Groningen, The Netherlands

K. H. Groenier · H. J. G. Bilo
University of Groningen,
Groningen, The Netherlands

H. J. G. Bilo
Department of Internal Medicine,
University Medical Center Groningen,
Groningen, The Netherlands