Correction to: A tailored approach in lymph node-positive perihilar cholangiocarcinoma

Christian Benzing¹ · Felix Krenzien¹ · Alexa Mieg¹ · Annika Wolfsberger¹ · Andreas Andreou¹ · Nora Nevermann¹ · Uwe Pelzer² · Uli Fehrenbach³ · Lena Marie Haiden¹ · Robert Öllinger¹ · Wenzel Schöning¹ · Moritz Schmelzle¹ · Johann Pratschke¹

Published online: 31 October 2021

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Correction to: Langenbeck's Archives of Surgery (2021) 406:1499–1509
<https://doi.org/10.1007/s00423-021-02154-4>

Figures 3 and 4 did not precisely match the text and were incorrect. Figure 3 refers to the survival according to the side of hepatic resection and not R-status. In Figure 4, the reasons for left/right hepatectomy not being feasible were reversed.

Corrected figures and figure legends are shown below:

Figure 3: Overall and disease-free survival of patients with lymph node positive perihilar cholangiocarcinoma according to side of hepatic resection

Kaplan Meier curves of patients with lymph node metastases and resected perihilar cholangiocarcinoma according to side of hepatic resection. A. Overall survival. B.

The original article can be found online at <https://doi.org/10.1007/s00423-021-02154-4>.

✉ Moritz Schmelzle
moritz.schmelzle@charite.de

¹ Department of Surgery, Campus Charité Mitte Campus Virchow-Klinikum, Experimental Surgery and Regenerative Medicine, Charité – Universitätsmedizin Berlin, Berlin, Germany

² Department of Hematology, Oncology and Tumor Immunology, Charité – Universitätsmedizin Berlin, Berlin, Germany

³ Department of Radiology, Charité - Universitätsmedizin Berlin, Berlin, Germany

Overall survival excluding 90-day mortality. C. Disease-free survival. D. Disease-free survival excluding 90-day mortality. E. Overall survival after propensity score

matching excluding 90-day mortality. F. Disease-free survival after propensity score matching excluding 90-day mortality.

Figure 4: Suggested therapy algorithm in resectable perihilar cholangiocarcinoma

Suggested tailored approach in patients diagnosed with resectable perihilar cholangiocarcinoma without distant metastases (Figure created with Biorender.com)

The original article has been corrected

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.