

Abscisic acid levels in tomato ovaries are regulated by *LeNCED1* and *SICYP707A1*

Lisette Maria Catharina Nitsch · Carla Oplaat · Richard Feron · Qian Ma · Mieke Wolters-Arts · Peter Hedden · Celestina Mariani · Wim Hendrik Vriezen

Received: 1 August 2008 / Accepted: 19 February 2009 / Published online: 26 March 2009
© The Author(s) 2009. This article is published with open access at Springerlink.com

Abstract Although the hormones, gibberellin and auxin, are known to play a role in the initiation of fruits, no such function has yet been demonstrated for abscisic acid (ABA). However, ABA signaling and ABA responses are high in tomato (*Solanum lycopersicum* L.) ovaries before pollination and decrease thereafter (Vriezen et al. in *New Phytol* 177:60–76, 2008). As a first step to understanding the role of ABA in ovary development and fruit set in tomato, we analyzed ABA content and the expression of genes involved in its metabolism in relation to pollination. We show that ABA levels are relatively high in mature ovaries and decrease directly after pollination, while an increase in the ABA metabolite dihydrophaseic acid was

measured. An important regulator of ABA biosynthesis in tomato is 9-*cis*-epoxy-carotenoid dioxygenase (*LeNCED1*), whose mRNA level in ovaries is reduced after pollination. The increased catabolism is likely caused by strong induction of one of four newly identified putative (+)ABA 8'-hydroxylase genes. This gene was named *SICYP707A1* and is expressed specifically in ovules and placenta. Transgenic plants, overexpressing *SICYP707A1*, have reduced ABA levels and exhibit ABA-deficient phenotypes suggesting that this gene encodes a functional ABA 8'-hydroxylase. Gibberellin and auxin application have different effects on the *LeNCED1* and *SICYP707A1* gene expression. The crosstalk between auxins, gibberellins and ABA during fruit set is discussed.

L. M. C. Nitsch · C. Oplaat · M. Wolters-Arts · C. Mariani
Department of Plant Cell Biology, IWWR,
Radboud University Nijmegen, Heyendaalseweg 135,
6525 AJ Nijmegen, The Netherlands

Q. Ma · P. Hedden
Crop Performance and Improvement Division,
Rothamsted Research Centre, Harpenden,
Hertfordshire AL5 2JQ, United Kingdom

Present Address:
R. Feron
KEYGENE N.V., P.O. Box 216, 6700 AE Wageningen,
The Netherlands

Present Address:
Q. Ma
National Key Laboratory of Crop Genetic Improvement,
Huazhong Agricultural University, 430070 Wuhan, China

Present Address:
W. H. Vriezen (✉)
Nunhems Netherlands B.V., P.O. Box 4005,
6080 AA Haalen, The Netherlands
e-mail: w.vriezen@science.ru.nl

Keywords ABA metabolism · Abscisic acid · (+)ABA 8'-hydroxylase · *LeNCED1* · *SICYP707A1* · Tomato

Abbreviations

ABA	Abscisic acid
GA	Gibberellin
NCED	9- <i>cis</i> -Epoxy-carotenoid dioxygenase
DPA	Dihydrophaseic acid
ABA-GE	ABA-glucose ester
PA	Phaseic acid
neo-PA	neo-Phaseic acid

Introduction

A major and nutritionally important part of the human diet is composed of fruits or fruit-derived products. In addition, fruits are important in the life cycle of plants, since they protect the seeds during their development and allow seed

dispersal. Therefore, there has been considerable research on fruit development, for which tomato has been used as a model system for physiological and molecular studies for a long time. The first step of fruit development, fruit initiation, is very important from an agro-economical point of view. It is a very delicate phase that is sensitive to abiotic factors, which often decrease fruit set and thus reduce yields. From a developmental point of view, fruit initiation, commonly referred to as fruit set, is the stage at which the ovary makes the decision to abort or to proceed with further cell division and fruit development (Gillaspy et al. 1993). Normally, fruit set is dependent on successful completion of pollination and fertilization (Gillaspy et al. 1993). However, it is not known precisely how the signals from pollination and fertilization stimulate cell division and fruit initiation.

It has long been known that application of hormones such as auxins or gibberellins can induce fruit set in the absence of fertilization. The resulting parthenocarpic fruit growth is a clear demonstration of the importance of hormones in fruit initiation and development. There are a number of parthenocarpic tomato mutants, such as *pat*, *pat2* and *pat3/4*, in which it was shown that the GA content and/or GA biosynthesis were increased significantly (Mazzucato et al. 1998; Fos et al. 2000, 2001; Olimpieri et al. 2007). Furthermore, overexpression of the *iaaM* gene, which encodes an enzyme involved in auxin biosynthesis, also induced parthenocarpic fruit development in tomato and many other species (Rotino et al. 1997; Mezzetti et al. 2004). Changes in both GA and auxin content were thus associated with parthenocarpic phenotypes. The altered hormone balance in the ovary of parthenocarpic plants seems, therefore, to substitute for pollination and fertilization (Gorguet et al. 2005).

Some of the genes functioning in the biosynthesis and signaling cascade of these hormones during fruit initiation have been identified recently. For instance, it was shown that expression of a GA 20-oxidase gene was induced by pollination in tomato (Rebers et al. 1999; Serrani et al. 2007), and the importance of *SIDELLA*, which represses GA signaling and tomato fruit set, has also been demonstrated (Marti et al. 2007). In addition, *IAA9* and *ARF8*, both transcriptional regulators of the auxin response, were shown to be part of an important signal for fruit set (Wang et al. 2005; Goetz et al. 2007). However, many other genes remain to be discovered and investigated. We have used transcript profiling to gain a better insight into the genes involved in fruit initiation, focusing on genes involved in hormonal signaling. Several groups of genes were found to have altered mRNA levels after pollination including genes involved in the cell cycle, posttranscriptional gene regulation, GA biosynthesis and auxin signaling (Vriezen et al. 2008). Notably, several genes related to abscisic acid

(ABA), such as *ABA RESPONSE ELEMENT BINDING PROTEIN 1*, *ABA INSENSITIVE1*-like genes, and dehydrin genes, were found to be highly expressed in mature ovaries while their expression decreased after pollination (Vriezen et al. 2008). Many factors related to ABA signaling and ABA responses are thus higher in mature, unpollinated ovaries and are quickly reduced after pollination. ABA is known to play a role in seed and bud dormancy, and in the regulation of abiotic stress responses (Horvath et al. 2003; Bartels and Sunkar 2005; Finkelstein et al. 2008). During tomato fruit development, its role is thought to be primarily restricted to the latter half of seed development in processes such as reserve deposition, desiccation tolerance and seed dormancy (Gillaspy et al. 1993), and a role in the first phase of fruit initiation has not been established. ABA has, however, been mentioned as a player in pea fruit set (Garcia-Martinez and Carbonell 1980; Rodrigo and Garcia-Martinez 1998). Here, we describe the changes found in the levels of ABA and the ABA metabolites dihydrophaseic acid (DPA) and phaseic acid (PA) in tomato ovaries and the regulation of ABA biosynthesis and catabolism genes during fruit initiation. Our data suggest that ABA is an additional player in the regulation of tomato fruit set together with auxin and GA.

Materials and methods

Plant material

Tomato plants (*Solanum lycopersicum* L. cv. Moneymaker from Enza Zaden, Enkhuizen, The Netherlands) were grown under greenhouse conditions from March to October under a 16/8 h day–night rhythm. Supplementary lights (600 W high-pressure sodium lights) turned on below 200 W/m² and turned off above 300 W/m². Temperature was kept above 20°C during the light period and 17°C during the dark period with the PRIVA Integro versie 724 system. Plants were watered daily and given fertilizer weekly.

Treatments

Flowers were emasculated 3 days before full bloom (dbf) and at full bloom either hand pollinated or treated with 2 µL of either 1 mM GA₃ (Duchefa Biochemie BV, Haarlem, The Netherlands) in 1% (v/v) ethanol or 1 mM 4-Cl-IAA (Sigma-Aldrich, St. Louis, MO, USA) in 2% (v/v) ethanol, or 1 mM ABA (Acros, Geel, Belgium) in 10% (v/v) methanol. 4-Cl-IAA is a more stable form of auxin than IAA and was used because it appeared to induce tomato fruit set more reliably than IAA. Control flowers were treated with 2 µL of the solvent (10% methanol or 1% ethanol). Hormone treatments were repeated after 2 days. Flowers were

harvested 3 days after first treatment unless otherwise noted. Plant material was sampled between 1100 and 1300 hours and directly frozen in liquid nitrogen. When necessary, plant material was dissected using binoculars and frozen thereafter in liquid nitrogen. The different tissues were dissected from young flowers that were about 3 dbf. Roots and hypocotyls were isolated from 10-day-old seedlings.

ABA analysis

Abscisic acid was quantified in triplicates by GC–MS as described for GA analysis (Griffiths et al. 2006), but with modifications. Freeze-dried samples were homogenized in 50 mL 80% (v/v) methanol/water in 100 mL flasks after which 300 ng 3-methyl- $^{2}\text{H}_3$ ABA was added as internal standard. Samples were purified as described previously except that the pooled ethyl acetate phases after elution from the Varian Bond Elut NH_2 cartridge (100 mg; Kinesis, St. Neots, UK) were evaporated to dryness in vacuo, and dissolved in ethyl acetate (20 μL) of which 2 μL was injected into a TR-1 capillary column (30 m \times 0.25 mm \times 0.25 mm film thickness; Thermo Fisher Scientific, Madison, WI, USA) at 50°C. The split valve (50:1) was opened after 2 min and the temperature increased at 20°C/min to 180°C and then at 4°C/min to 300°C. The instrument was operated in selective ion monitoring mode, monitoring the ions at m/z 190 and 162 for ABA and 193 and 166 for $^{2}\text{H}_3$ ABA. The amount of ABA was determined from the peak areas for the ions m/z 190 and 193 by reference to a calibration curve.

ABA catabolite analysis

The ABA catabolite analysis was performed at the Plant Biotechnology Institute of the National Research Council of Canada (<http://www.pbi.nrc.gc.ca/ENGLISH/technology-platforms/plant-hormone-profiling.htm>) by high-performance liquid chromatography electrospray tandem mass spectrometry (HPLC-ES-MS/MS) using deuterated internal standards, as described in Owen and Abrams (2009).

Neighbor joining tree

NCED mRNA sequences from *Arabidopsis*, rice, maize, tomato and potato were obtained from NCBI (www.ncbi.nlm.nih.gov) or Dana-Farber Cancer Institute (DFCI) Tomato Gene Index (<http://compbio.dfci.harvard.edu/tgi>). The sequences were translated to protein and aligned with ClustalW (www.ebi.ac.uk/clustalw/). A neighbor joining tree was generated using the PHYLIP Protdist program (Phylogeny Inference Package version 3.5c; J. Felsenstein, Department of Genetics, University of Washington,

Seattle) available on <http://bioweb.pasteur.fr/seqanal/phylogeny/phylip-uk.html>. Default parameters were used and 100 bootstrap replicates were generated. The consensus tree was drawn in TreeView (free available from <http://taxonomy.zoology.gla.ac.uk/rod/treeview.html>). An unrooted tree was drawn from output data without bootstrap values, bootstrap values (calculated by Prodist) were manually added afterwards.

Isolation and cloning of *SICYP707A1* and *CYP707* like cDNAs

The full-length *SICYP707A1* sequence was isolated by plaque screening of a phage cDNA library (HybriZAP[®] 2.1, Stratagene, La Jolla, CA, USA) with a probe corresponding to base 1,024–1,387 of *SICYP707A1*, following the manufacturer's protocol. TC177455, AI484420 and TC186477 sequences were obtained from the DFCI Tomato Gene Index.

RNA isolation, cDNA synthesis and Q-PCR data analysis

RNA was isolated with the RNeasy kit (Qiagen, Valencia, CA, USA). Photometric RNA measurements were done to equilibrate the RNA concentrations of different samples. Equal amounts of RNA were DNase treated (RQ1, Promega, Madison, WI, USA). RNA (0.5 μg) was reverse transcribed (RT) in a total volume of 10 μL using a cDNA synthesis kit (iScript[™], Bio-Rad Laboratories, Hercules, CA, USA) following manufacturer's protocol.

Real-time quantitative PCR (Q-PCR) primers were designed using a computer program (Beacon Designer Software, Premier Biosoft International, CA, USA). Alignments of genes that were part of a gene family were made and gene specific DNA pieces were used for primer design. Primer pairs are depicted in Table 1. Specificity was tested by sequencing the amplicon. Q-PCRs were done using SYBR green mix (iQ-SYBR Green Supermix, Bio-Rad Laboratories). PCR reactions were performed in a 96-well thermocycler (Bio-Rad iCycler) using a temperature program starting with 3 min at 95°C then 40 cycles consisting of 15 s at 95°C and 45 s at 57°C and finally the melting temperature of the amplified product was determined to verify the presence of a single product. Five microliters of 25-fold diluted cDNA was used per sample. Technical and biological replicates were always performed. Both actin 2/7 and ubiquitin 7 were used as internal control genes, to correct for difference in cDNA amounts. Diluted DNase-treated RNA was also included in the Q-PCR as a control for genomic DNA contamination.

Relative mRNA levels were calculated following the Bio-Rad outlined methodology based on Vandesompe et al. (2002) and corrected for PCR efficiencies, which were

Table 1 Primer sequences used for semi- and real-time quantitative PCR

Gene name or number (annotation)	Primer sequences
LeNCED1 (9- <i>cis</i> -epoxy-carotenoid dioxygenase)	5'-CTTATTTGGCTATCGCTGAACC-3' 5'-CCTCCAACTTCAAACCTCATTGC-3'
SICYP707A1 (ABA 8'-hydroxylase)	5'-AGAGAGGCTGTAGCTGAGTGG-3' 5'-TTGGCAAGTTCATTCCCTGGAC-3'
TC177455 (ABA 8'-hydroxylase)	5'-GCAATGAAAAGCGAGGAAAGAGC-3' 5'-TCGAGCTGCAAAGATGACTCC-3'
AI484420 (Cytochrome P450)	5'-CTAAGGTGGCAAGGAGGAAGC-3' 5'-GTGTCCTGGGCAGCAAAGAG-3'
TC186477 (ABA 8'-hydroxylase)	5'-GAGCATTCAAACCCGAAGCC-3' 5'-AATTGTACCCTGTTTCGAGCAC-3'
ACT2/7 (actin 2/7)	5'-GGACTCTGGTGTGGTGTAG-3' 5'-CCGTTTCAGCAGTAGTGGTG-3'
UBQ7 (ubiquitin 7)	5'-CCCTGGCTGATTACAACATTC-3' 5'-TGGTGTCAAGTGGGTTCAATG-3'

determined using dilution ranges (0.1–10 pg/μL). The average of two biological repeats and two technical repeats is depicted together with the SE. When error bars were omitted in graphs, biological replicates showed the same trend but the absolute amounts were different—in these cases, only one biological replicate is shown.

Semi-quantitative PCR

Semi-quantitative PCRs were performed using 5 μL of 25-fold diluted cDNA, buffer IV, 2.5 mM MgCl₂, 0.5 unit Red Hot Taq DNA polymerase (all from ABgene Limited, Epsom, Surrey, UK), 0.4 mM dNTPs (Fermentas, St. Leon-Rot, Germany), and 0.1 μM primers (Table 1) in a reaction of 30 cycles, each comprising of 15 s at 95°C (denaturation), 30 s at 57°C (primer annealing) and 10 s at 72°C (extension time).

In situ hybridization

The RNA probes were synthesized by transcribing bases 1,467–1,648 of the *CYP707A1* cDNA using T7 (sense) and SP6 (anti-sense) RNA polymerase and digoxigenin (DIG)-labeled UTP (Roche Applied Science, Basel, Switzerland). DIG labeling efficiencies were tested by a spot assay. Tissue fixation was done as described by Bereterbide et al. (2002). Embedded tissues were sliced into 8 μm sections. Paraplast Plus (Sigma-Aldrich, St. Louis, MO, USA) was removed with Histoclear (National Diagnostics, Atlanta, GA, USA), sections were hydrated, immersed in TE (Tris-HCl pH 7.5, 10 mM EDTA) and a proteinase K treatment was performed. Sections were then dehydrated again, and air dried for 2–3 h. The slides were incubated in a humidified box overnight at 50°C, in 150 μL hybridization mixture

(50% formamide, 0.5 μg/mL tRNA, 10% (w/v) dextran sulfate, 300 mM NaCl, 5 mM EDTA, 10 mM Tris-HCl pH 7.5) supplemented with 100 ng of probe. The next morning slides were immersed in a series of SSC (Na₂citrate, NaCl) washing buffers until 0.5× SSC. Anti-DIG alkaline phosphatase-coupled antibody (Roche Applied Science) was diluted 500-fold in 1% (w/v) BSA solution and after blocking a volume of 150 μL was added to the slides and incubated in a humid chamber at 37°C for 1.5 h. After two washes, the slides were put in color substrate reaction buffer [100 mM NaCl, 5 mM MgCl₂, 100 mM Tris-HCl pH 9.0, 10% (w/v) polyvinylalcohol, 0.24 mg/mL Levamisole (Sigma-Aldrich), 5 μL/mL NBT and 3.75 μL/mL BCIP (Roche Applied Science)] in the dark until purple staining appeared. Staining was stopped by immersing the slides in TE.

Overexpression lines

To generate transgenic overexpression *SICYP707A1* lines, the coding region (base 6–1,478) was PCR amplified and cloned in the Gateway entry vector pENTR/D-TOPO vector (Invitrogen, Carlsbad, CA, USA). *SICYP707A1* coding region was recombined between the Cauliflower Mosaic Virus 35S promoter in the pGD625 vector (Chalfun-Junior et al. 2005) and the *NOPALINE SYNTHASE* terminator. Transgenic plants were generated by *Agrobacterium tumefaciens*-mediated transformation. Cotyledons of 10-day-old seedlings were cut and pre-incubated overnight on co-cultivation MS medium containing vitamin Gamborg B5, 1% (v/w) sucrose, 0.5 mg/L Mes buffer, 0.8% (v/w) Daishin agar, 0.05 mg/L 2,4-D, 0.1 mg/L IAA, 2.0 mg/L zeatin, 200 μM acetosyringone, pH 5.8. *A. tumefaciens* (EHA 105) were grown to an OD₆₀₀ of 1.0 and diluted 25

times in liquid LB media and grown for another 4 h to an OD600 between 0.2 and 0.3. Bacteria were pelleted and resuspended in liquid MS medium containing vitamin Gamborg B5 mixture, 3% (w/v) sucrose, 0.5 mg/L Mes buffer, 200 μ M acetosyringone. Cotyledons were incubated 30 min in the bacterial suspension, washed and placed back on the co-cultivation media. Two days after transformation, cotyledons were placed on MS medium containing vitamin Gamborg B5 mixture, 3% (v/w) sucrose, 0.5 mg/L Mes buffer, 0.8% (v/w) Daishin agar, 0.1 mg/L IAA, 2.0 mg/L zeatin, 200 mg/L cefotaxime and 50 mg/L vancomycin and incubated for another 2 days. To induce callus and shoot formation, cotyledons were transferred to shoot inducing MS medium containing vitamin Gamborg B5 mixture, 1% (w/v) glucose, 0.5 mg/L Mes buffer, 0.8% (w/v) Daishin agar, 0.1 mg/L IAA, 2.0 mg/L zeatin, 500 mg/L carbenicillin and 100 mg/L kanamycin. Cotyledons were transferred to fresh medium every 3 weeks. Elongated shoots of 2–4 cm were excised from the callus and transferred to rooting MS culture medium containing 1.5% (w/v) sucrose, 4% (w/v) purified agar (Oxoid Ltd, Hampshire, UK), 0.25 mg/L IBA, 100 mg/L cefotaxime and 100 mg/L vancomycin. Rooted plantlets were transferred to soil for further selection including, ploidy test and PCR with primers specific for the kanamycin resistance gene. Media components and antibiotics were obtained from Duchefa Biochemie BV unless otherwise specified.

Water stress experiment

Pots of wild type and *SICYP707A1* overexpression lines of the same age and size were saturated with water at the start of the experiment. Plants were withheld water from then on. Photographs were taken 72 h after start of the experiment.

Leaf surface area

Leaf surface area was measured for four compound leaves of 2 months old tomato plants on a Li-3100 area meter (Li-Cor, Lincoln, NE, USA). Average leaf surface is depicted with SD. Leaves were always picked at the same height.

Accession details

GenBank accession numbers: *LeNCED1* (Z97215), *AtNCED2* (NM117945), *AtNCED3* (NM112304), *AtNCED4* (NM118036), *AtNCED5* (NM102749), *AtNCED6* (NM113327), *AtNCED9* (NM106486), *SlNCED1* (AY662343), *OsNCED1* (AY838897), *OSNCED2* (AY838898), *OsNCED3* (AY838899), *OsNCED4* (AY838900), *OsNCED5* (AY838901), *VP14* (ZMU95953), *AhNCED1* (AJ574819), *SICYP707A1* (EU183406), *SICYP707A3* (AI484420), *StCY*

P707A2 (DQ206631), *AtCYP707A1* (NM202845), *AtCYP707A2* (NM128466), *AtCYP707A3* (NM180805), *AtCYP707A4* (NM112814), *ACT2/7* (BT013707), *UBQ7* (AK246454).

DFCI Tomato Gene Index accession numbers: *TC175556*, *TC154638*, *TC154637*, *TC156234*, *SICYP707A2* (TC177455), *SICYP707A4* (TC186477).

Results

ABA content

Abscisic acid signaling and ABA response genes are highly expressed in unpollinated ovaries and their expression decreases upon pollination (Vriezen et al. 2008), suggesting a concomitant decrease in ABA concentration. To confirm this assumption, ABA concentration was determined in ovaries at several time points before and after pollination. Figure 1 shows that the free-ABA level in ovaries significantly decreased from 6.7 ng/mg DW before pollination (C0) to 2.4 ng/mg DW 3 days after pollination (dap). In contrast, if ovaries were kept unpollinated during the same period, the free-ABA level increased to 7.6 ng/mg DW (C3).

Regulation of ABA biosynthesis

It is generally accepted that the cleavage reaction catalyzed by 9-*cis*-epoxy-carotenoid dioxygenases (NCED) is a major rate-limiting step and point of regulation in ABA biosynthesis (Qin and Zeevaart 2002). In other species, such as *Arabidopsis* and rice, NCED enzymes are encoded by small gene families composed of five and three members, respectively (Tan et al. 2003). However, in tomato




Fig. 1 ABA concentration decreases in ovaries after pollination. ABA concentration in unpollinated ovaries of emasculated flowers at full bloom (C0) and in unpollinated ovaries 3 days after full bloom (C3), and 1–3 dap: 1, 2, and 3 days after pollination. Mean values ($n = 3$) \pm SE are depicted

only one NCED (*LeNCED1*) has been identified so far, and shown to have 9-*cis*-epoxy-carotenoid cleavage activity. In an attempt to obtain a complete overview of the transcriptional regulation of the whole NCED gene family in tomato, we searched the DFCI Tomato Gene Index and NCBI EST databases for sequences homologous to *LeNCED1* or to one of the five known *Arabidopsis* genes. The neighbor joining tree in Fig. 2 shows that the four tomato sequences most similar to *LeNCED1* did not group together with the *LeNCED1* and the other functional NCED genes from *Arabidopsis*, rice, potato and maize. Three of the four tomato sequences (TC175556, TC154638, and TC154637) formed a separate group, while the other sequence (TC156234) grouped together with other carotenoid cleavage dioxygenases (CCDs) that are not known to be related to ABA biosynthesis (Tan et al. 2003). In addition, the deduced amino acid sequences of the four tomato sequences were less than 35% identical to the functional *Arabidopsis* or tomato NCED amino acid sequences. Therefore, we think it is highly likely that no additional functional NCED genes are present in the tomato genome.

LeNCED1 mRNA levels decreased after pollination in both the ovules and placenta tissue and the pericarp (Fig. 3a). Semi-quantitative PCR analysis, with specific *LeNCED1* primers, revealed that it is expressed equally in wall, ovules and placenta in unpollinated ovaries (Fig. 4a). After pollination, there is a decrease in expression in wall and ovules. Because of its equal distribution throughout the

unpollinated ovary, no in situ hybridization was performed for *LeNCED1*.

Regulation of ABA catabolism

Abscisic acid concentration is regulated by catabolism as well as synthesis. In plant cells, ABA is mainly inactivated by 8'-hydroxylation (Nambara and Marion-Poll 2005), which is in *Arabidopsis* catalyzed by four cytochrome P450 mono-oxygenases, *AtCYP707A1* to *AtCYP707A4* (Kushiro et al. 2004). We previously identified a tomato homolog of *AtCYP707A4*, which was strongly induced in the ovary after pollination (Vriezen et al. 2008). We have now isolated and characterized the corresponding full-length (1,742 bp) cDNA clone from a tomato ovary cDNA library. The deduced amino acid sequence is highly homologous to the *Arabidopsis* CYP707A4 protein (65% amino acid identity) and *Solanum tuberosum* CYP707A2 gene (95% identity). The tomato protein (designated *SICYP707A1*) contains the highly conserved cysteine residue (within the PFGNGTHSC_{CPG} motif), which is the putative heme-iron-ligand, common to all P450s and essential for catalytic activity (Kushiro et al. 2004). A database search yielded three more putative CYP707A ESTs (TC177455, AI484420, and TC186477) that share between 61 and 68% sequence identity with *SICYP707A1* at nucleotide level. Their deduced protein sequences are each 70–74% identical with one of the four *Arabidopsis* *AtCYP707A* proteins. We, therefore, consider the three genes to encode putative ABA 8'-hydroxylases, and designated them *SICYP707A2*, *SICYP707A3* and *SICYP707A4*.

The mRNA levels of the four putative tomato *CYP707A* genes were determined in emasculated flowers 3 days after full bloom (control) and 3 days after pollination in tomato ovaries. Figure 3b shows that the mRNA concentration of *SICYP707A1* strongly increased after pollination in whole ovaries and it reached a much higher level than the mRNA levels of the other *CYP707A* genes. In addition, the mRNA levels of all other putative tomato *CYP707A* genes were reduced after pollination. This suggests that *SICYP707A1* is the most important gene responsible for ABA catabolism in pollinated ovaries. Expression of *SICYP707A1* was also analyzed in other tissues and it is expressed in sepals, petals and pedicels (Fig. 3c). The accumulation of *SICYP707A1* mRNA in ovaries was determined more precisely in separated ovule/placenta and pericarp tissue during the first 3 days after pollination. *SICYP707A1* mRNA levels were several hundredfold up-regulated after pollination, specifically in ovules/placenta and not in pericarp (Fig. 3d).

Furthermore, mRNA localization was analyzed in ovules and placenta separately. In unpollinated (control) ovaries *SICYP707A1* mRNA is present in the placenta, but not in





Fig. 2 Neighbor joining tree of NCED and NCED-like sequences with bootstrap values indicated at junctions. *LeNCED1* groups together with known NCED sequences such as *AtNCED2,3,5,6,9*, *VP14* and *OsNCED3,5,6*. Two other groups are formed including TC175556, TC154638, TC154637, TC156234, *OsNCED1,2* and *AtNCED4*. Tomato sequences are *underlined*. For accession details see “Materials and methods”

Fig. 3 **a** Relative mRNA levels of *LeNCEDI*. *LeNCEDI* mRNA levels decrease in both pericarp and ovules/placenta after pollination. **b** mRNA levels of *SICYP707A1*, *SICYP707A2*, *SICYP707A3*, and *SICYP707A4* (CYP1–CYP4) in ovaries 3 days after full bloom (control) or after pollination. **c** mRNA levels of *SICYP707A1* in different generative and vegetative tissues of tomato; hypocotyl (H) and root (R) tissue from 10-day-old seedlings, and pedicel (P), sepal (Se), petal (Pt), anther (A) and ovary (O) tissue from flowers 3 days before full bloom. **d** mRNA levels of *SICYP707A1* in more detail in separated ovules/placenta and pericarp tissue. *SICYP707A1* mRNA levels are increased after pollination specifically in ovule/placenta tissue. Mean values ($n = 4$) \pm SE



the ovules and pericarp (Fig. 4a). In pollinated ovaries, *SLCYP707A1* mRNA level is generally higher and present in both placenta and ovule tissue. In situ hybridization with a gene-specific probe confirmed this localization of *SICYP707A1* in pollinated ovaries. The purple staining suggests relatively high *SICYP707A1* mRNA levels in the ovules and in the periphery of the placenta but low levels in the pericarp (Fig. 4b, c).

Gene expression data suggest that the reduction in ABA levels that we measured after pollination could be due to increased ABA 8'-hydroxylation activity. To strengthen this hypothesis, we analyzed the levels of the ABA catabolic products in ovaries 3 days after pollination and 3 days after anthesis in unpollinated ovaries. These tissues are

comparable to the C3 and 3 dap samples, respectively, which were used to measure ABA (Fig. 1). Table 2 shows that the level of ABA in unpollinated ovaries was two-and-half-fold higher than in pollinated ovaries, comparable to the measurements shown in Fig. 1. The product of ABA 8'-hydroxylation is (–)-PA which can be reduced to (–)-DPA (Zeevaert 1999). PA levels did not change, but DPA levels increased from 0.44 to 1.20 ng/mg DW in ovaries after pollination. 7'- and 9'-ABA hydroxylation can also inactivate ABA in several plant species (Nambara and Marion-Poll 2005). The product of ABA 9'-hydroxylation is 9'-hydroxy ABA which is converted to neo-phaseic acid (neo-PA). In tomato ovaries, a considerable amount of neo-PA was present, but this was not affected by pollination. The presence




Fig. 4 Localization of *LeNCED1* and *SICYP707A1* mRNA in ovaries. **a** Semi-quantitative PCR with *LeNCED1* and *SICYP707A1* specific primers, and actin 2/7 as an internal control gene. *LeNCED1* is equally expressed in wall, placenta and ovules of unpollinated ovaries and expressed lower after pollination in wall and ovules. *SICYP707A1* is weakly expressed in the placenta of unpollinated ovaries and expressed higher in both ovules and placenta in pollinated ovaries. In situ hybridization with a sense (**b**) and anti-sense (**c**) RNA probe of *SICYP707A1*. Alkaline phosphatase activity gives a purple staining. Bar represents 100 μ m

of 7'-hydroxy ABA was also detected, but its level was low and could not be quantified significantly. The amount of a conjugated form of ABA, ABA-glucosyl ester (ABA-GE), decreased from 1.32 to 0.65 ng/mg DW in ovaries after pollination (Table 2).

Functional analysis of *SICYP707A1*

To prove that the *SICYP707A1* gene is coding for a functional ABA 8'-hydroxylase, we overexpressed its coding region under control of the 35S-promoter in tomato. Several transgenic lines were obtained, of which the line with the highest overexpression level was further analyzed. Figure 5a shows the *SICYP707A1* mRNA levels in unpollinated mature ovaries. The *SICYP707A1* overexpression line has approximately 45-fold higher mRNA levels than wild type. Consequently, the free-ABA level in unpollinated ovaries was 46% lower in *SICYP707A1* overexpression plants than in wild-type plants (Fig. 5b), confirming the ABA 8'-hydroxylase activity of *SICYP707A1*. The overexpression plants have a smaller leaf surface area and wilt




Fig. 5 **a** Relative mRNA levels of the *SICYP707A1* gene in unpollinated mature ovaries of overexpression (OE) plants are 45 times higher than wild type (wt). **b** Relative ABA levels in unpollinated mature ovaries of overexpression plants is 56% of wild type. **c** Overexpression plants (right) wilt stronger after 72 h of water with holding than wild-type plants (left). **d** The leaf surface area of the compound leaves of overexpression plants is smaller than wild type. **e** On the stem of the overexpression plants small adventitious roots are visible

stronger 72 h after water with holding than wild-type plants (Fig. 5c, d). In addition, we also observed the initiation of adventitious root growth on the stem of the *SICYP707A1* overexpression plants (Fig. 5e).

Hormone crosstalk

Because fruit set is induced by GA and auxin, we were interested in the effect of these hormones on the activity of both, the *LeNCED1* gene and the *SICYP707A1* gene. The *LeNCED1* mRNA level in ovaries was lower after GA₃ treatment or 4-Cl-IAA treatment and comparable to the levels found after pollination (Fig. 6a). ABA treatment, however, increased the mRNA level of *LeNCED1*. *SICYP707A1* mRNA level was not affected by GA₃ treatment in contrast to pollination, 4-Cl-IAA treatment, or ABA treatment, which all had a stimulating effect (Fig. 6b).

Table 2 ABA and catabolite levels in unpollinated and pollinated ovaries

	ABA (ng/mg DW)	PA (ng/mg DW)	DPA (ng/mg DW)	ABA-GE (ng/mg DW)	neo-PA (ng/mg DW)
Unpollinated	10.76 (\pm 0.12)	0.10 (\pm 0.05)	0.44 (\pm 0.05)	1.32 (\pm 0.04)	0.46 (\pm 0.09)
Pollinated	4.26 (\pm 0.31)	0.10 (\pm 0.06)	1.20 (\pm 0.08)	0.65 (\pm 0.05)	0.46 (\pm 0.01)

Mean values ($n = 2$) \pm SE are depicted




Fig. 6 mRNA levels of *LeNCED1* and *SICYP707A1* before (C = control) and after hormone treatment or after pollination. **a** *LeNCED1* expression is induced by ABA treatment and reduced after pollination, GA₃ and 4-Cl-IAA (IAA) treatment. **b** *SICYP707A1* expression is induced after pollination, 4-Cl-IAA and ABA treatment. Mean values ($n = 6$) \pm SE

Discussion

ABA content

Fruit set is an agro-economically important process, in which hormones play a pivotal role. Gibberellins and auxins were known previously to induce fruit growth and, by demonstrating that ABA signaling is high in mature ovaries and decreases after pollination, we have recently suggested a role for ABA in fruit set (Vriezen et al. 2008). In addition, ABA has been immuno-localized in the ovary of *Arabidopsis* and cucumber, indicating a function for ABA in these tissues (Peng et al. 2006). Previously, Kojima et al. (1993) had measured relatively high levels of ABA in unpollinated pistils compared to 10-day-old tomato fruits. Here, we demonstrate that within 3 days the free-ABA content of pollinated tomato ovaries decreases to approximately 30% of that in unpollinated ovaries. A decrease in ABA concentration to 50 or 25% of the original concentration has also been measured in other physiological systems, such as in dormancy breakage in seeds (Chiwocha et al. 2005) and tubers (Destefano-Beltran et al. 2006). The relative decrease in ABA content in the tomato ovary is comparable to those observed in other processes in which ABA level is of physiological importance.

Regulation of ABA biosynthesis and catabolism

Abscisic acid concentration is actively regulated in ovaries, as we found biosynthesis genes to be down-regulated and ABA catabolism genes to be induced in the ovary after fruit initiation. NCED genes are often encoded by a small gene family. However, we were not able to find putative NCED genes other than *LeNCED1* in the tomato sequence databases, which contain 213,947 ESTs and 41,425 unique sequences. The three other CCDs described are probably

not related to ABA biosynthesis as they do not group with *LeNCED1*. Moreover, TC154637 and TC154638 have already been described as *carotenoid cleavage dioxygenase 1* genes involved in volatile terpenoid production (Simkin et al. 2004). Therefore, *LeNCED1* appears to produce the most represented NCED transcript in ovary and possibly is the only NCED gene in tomato. In addition, the fact that the *notabilis* mutant with a mutation in the *LeNCED1* gene has a strong ABA-deficient phenotype (Burbidge et al. 1999; Thompson et al. 2004) indicates that this gene has a prominent role in ABA biosynthesis. Transcript levels for this gene decreased after pollination concomitant with the decrease in ABA concentration, indicating that *LeNCED1* regulation directly affects biosynthesis of ABA in the mature ovary.

Transcription levels of a gene very homologous to an *Arabidopsis* ABA 8'-hydroxylase (*AtCYP707A4*) were strongly induced specifically in ovules and placenta tissue, in contrast to the mRNA levels of the three other putative tomato *CYP707A* genes. Similarly, the *CYP707A* genes in *Arabidopsis* are also differentially expressed, for instance during dehydration and rehydration responses and during seed dormancy (Kushiro et al. 2004; Okamoto et al. 2006). The deduced protein sequence of *SICYP707A1* has very high homology to *AtCYP707A4*. Moreover, we have indirectly proven that the *SICYP707A1* gene encodes a functional ABA 8'-hydroxylase by overexpressing it in tomato. This resulted in reduced ABA levels and ABA-deficient phenotypes, such as adventitious rooting, reduced leaf surface area and increased wilting. These characteristics have also been described for the tomato ABA-deficient mutants *sitiens*, *notabilis* and *flacca* (Taylor and Tarr 1984; Burbidge et al. 1999). The remarkably large increase we found in *SICYP707A1* mRNA levels suggests a strong induction of ABA 8'-hydroxylation after pollination. In addition, we showed that after pollination a higher level of DPA, the end product of the ABA 8'-hydroxylation pathway, was present in the ovary, suggesting that ABA 8'-hydroxylation was indeed induced after pollination. Interestingly, 7'- and 9'-hydroxylation of ABA also takes place in tomato. However, no differences were found between 7'-hydroxy ABA and neo-PA levels in ovaries after pollination, suggesting that these conversions take place, but do not attribute to the decrease in ABA levels after pollination. Thus, ABA 8'-hydroxylation is the major ABA catabolic pathway in tomato ovaries (our work) as was suggested for *Arabidopsis* (Kushiro et al. 2004). ABA can be transported and stored as ABA-GE (Nambara and Marion-Poll 2005) and the high levels in unpollinated ovaries could mean that ABA-GE is transported to and stored in the ovary. Altogether, these data strengthen our hypothesis that after pollination ABA levels are reduced mainly through ABA 8'-hydroxylation. The specific localization of

SICYP707A1 in ovules and placenta might indicate that embryo or seed formation requires rapid removal of ABA. The localization of *SICYP707A1* mRNA is comparable to the localization of the hormone ABA in the ovules of *Arabidopsis* and cucumber (Peng et al. 2006). The decrease in mRNA levels of the three other putative tomato CYP707A genes might be explained by the reduction in ABA concentration, since positive feedback from ABA on CYP707A expression has been found in *Arabidopsis* (Kushiro et al. 2004; Umezawa et al. 2006), potato (Destefano-Beltran et al. 2006) and by us in tomato (Fig. 6b).

Hormone crosstalk

Auxin has been shown to interact with several hormones via regulation of expression of genes encoding biosynthetic or catabolic enzymes (Ross and O'Neill 2001). A fertilization-induced increase in auxin signaling most likely regulates the expression of ABA biosynthesis and catabolism genes during tomato fruit set, since we could show that auxin treatment reduced *LeNCED1* transcript levels and increased *SICYP707A1* transcript levels. GA₃ treatment also reduced *LeNCED1* gene expression but it had no effect on *SICYP707A1* gene expression. Therefore, auxin, via GA, or both hormones together regulate *LeNCED1* gene expression, whereas *SICYP707A1* expression seems to be regulated by auxin alone. This would be consistent with auxin inducing GA biosynthesis, but not vice versa (Vriezen et al. 2008; de Jong et al. 2009). It seems likely that a pollination-induced auxin increase precedes the GA increase, as summarized in Fig. 7. This hormonal signaling cascade may also lead to a reduction in the ABA content via changes in *LeNCED1* and *SICYP707A1* expression. In the promoter of *LeNCED1*, auxin, GA, and ABA response elements were identified (Thompson et al. 2004), consistent with *LeNCED1* being regulated by all three hormones.




Fig. 7 Model of the hormonal interactions after pollination. Pollination induces an increase in auxin concentration and subsequently an increase in GA concentration. Both an increase in GA and IAA levels inhibit the expression of the *LeNCED1* gene (ABA biosynthesis), while only increased IAA levels can stimulate the expression of the *SICYP707A1* gene (catabolism), ABA levels therefore decrease

Since we found that *LeNCED1* is positively regulated by ABA, a decrease of ABA content after pollination might also contribute to the decrease in its mRNA levels. A positive feedback of ABA levels on *NCED* mRNA levels has been shown in *Arabidopsis* (Wan and Li 2006). However, Thompson et al. (2000) showed that ABA had no effect on *LeNCED1* expression in a detached leaf assay. Different tissues might thus have different regulatory mechanisms of *LeNCED1* expression. The increase in *SICYP707A1* levels cannot be caused by a lower content of ABA, because this gene is also positively regulated by ABA.

Function of ABA during fruit set

In a mature unpollinated ovary, many cell cycle genes, such as *LeCDKB1,2*, *CycA1,2* and *CycD3*, are relative lowly expressed (Vriezen et al. 2008). In addition, the relative expression level in the mature tomato ovary of two genes homologous to *PsDRM1*, which proved to be excellent dormancy (DRM) markers in pea (Stafstrom et al. 1998), are high. Together, this suggests that the mature ovary before pollination has entered a temporarily dormant state, during which the tissue is quiescent. It is known that ABA can directly influence the cell cycle via suppression of *CDKA1* expression (Smalle et al. 2003) or stimulation of the *ICK1* (an inhibitor of CDK action) expression (Wang et al. 1998), thereby inhibiting cell division and inducing an quiescent state. We know that the ABA signaling and response pathway are active in the unpollinated mature ovary (Vriezen et al. 2008). In addition, in the research presented here, we have shown that ABA levels are relatively high in mature ovaries and are down-regulated after pollination. Based on these observations, a hypothesis is that ABA inhibits growth until fruit set. The post-pollination increase of auxin and GA levels may directly or indirectly repress ABA biosynthesis. However, we were unable to induce fruit set by application of an ABA biosynthesis inhibitor, fluridone. Neither could we inhibit fruit set by ABA application to pollinated ovaries (data not shown). This might indicate that not only the concentration of ABA but rather the hormonal balance of ABA and other hormones, such as auxin and GA, is important. Similarly, the potential of axillary bud outgrowth is determined by a balance of the hormones such as cytokinin, auxin and ABA (Shimizu-Sato and Mori 2001). Besides inhibition of growth, ABA can also induce tolerance to several abiotic stresses, such as heat stress, cold stress and drought stress (Bartels and Sunkar 2005; Gusta et al. 2005; Bonham-Smith et al. 2008). For example, ABA induces dehydrins which are proteins with a potential in vivo role in stabilizing cells under abiotic stress (Kalemba and Pukacka 2007). Several dehydrins are indeed higher expressed in the mature unpollinated ovary (Vriezen et al. 2008). A putative protective function for ABA might

become more apparent during unfavorable environmental conditions when ABA protects the mature ovary during its temporal dormant state.

Acknowledgments We thank Dr. Ivo Rieu (University of Freiburg, Germany) for his cooperation and fruitful scientific discussions. Part of the research was funded by a grant from the Netherlands Organization for Scientific Research (NWO, R 89-182).

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Bartels D, Sunkar R (2005) Drought and salt tolerance in plants. *Crit Rev Plant Sci* 24:23–58
- Bereterbide A, Hernould M, Farbos I, Glimelius K, Mouras A (2002) Restoration of stamen development and production of functional pollen in an alloplasmic CMS tobacco line by ectopic expression of the *Arabidopsis thaliana* *SUPERMAN* gene. *Plant J* 29:607–615
- Bonham-Smith PC, Kapoor M, Bewley JD (2008) Exogenous application of abscisic acid or triadimefon affects the recovery of *Zea mays* seedlings from heat shock. *Physiol Plant* 73:27–30
- Burbidge A, Grieve TM, Jackson A, Thompson A, McCarty DR, Taylor IB (1999) Characterization of the ABA-deficient tomato mutant *notabilis* and its relationship with maize *Vp14*. *Plant J* 17:427–431
- Chalfun-Junior A, Franken J, Mes JJ, Marsch-Martinez N, Pereira A, Angenent GC (2005) *ASYMMETRIC LEAVES2-LIKE1* gene, a member of the AS2/LOB family, controls proximal–distal patterning in *Arabidopsis* petals. *Plant Mol Biol* 57:559–575
- Chiwocha SDS, Cutler AJ, Abrams SR, Ambrose SJ, Yang J, Ross ARS, Kermod AR (2005) The *etr1-2* mutation in *Arabidopsis thaliana* affects the abscisic acid, auxin, cytokinin and gibberellin metabolic pathways during maintenance of seed dormancy, moist-chilling and germination. *Plant J* 42:35–48
- de Jong M, Wolters-Arts M, Feron R, Mariani C, Vriezen WH (2009) The *Solanum lycopersicum* auxin response factor 7 (*SIARF7*) regulates auxin signaling during tomato fruit set and development. *Plant J* 57:160–170
- Destefano-Beltran L, Knauber D, Huckle L, Suttle JC (2006) Effects of postharvest storage and dormancy status on ABA content, metabolism, and expression of genes involved in ABA biosynthesis and metabolism in potato tuber tissues. *Plant Mol Biol* 61:687–697
- Finkelstein R, Reeves W, Ariizumi T, Steber C (2008) Molecular aspects of seed dormancy. *Annu Rev Plant Biol* 59:387–415
- Fos M, Nuez F, Garcia-Martinez JL (2000) The gene *pat-2*, which induces natural parthenocarpy, alters the gibberellin content in unpollinated tomato ovaries. *Plant Physiol* 122:471–479
- Fos M, Proano K, Nuez F, Garcia-Martinez JL (2001) Role of gibberellins in parthenocarpic fruit development induced by the genetic system *pat-3/pat-4* in tomato. *Physiol Plant* 111:545–550
- Garcia-Martinez JL, Carbonell J (1980) Fruit-set of unpollinated ovaries of *Pisum sativum* L. influence of plant-growth regulators. *Planta* 147:451–456
- Gillaspy G, Ben-David H, Grissem W (1993) Fruits—a developmental perspective. *Plant Cell* 5:1439–1451
- Goetz M, Hooper LC, Johnson SD, Macedo Rodrigues JC, Koltunow AM (2007) Expression of aberrant forms of *AUXIN RESPONSE FACTOR8* stimulates parthenocarpy in *Arabidopsis* and tomato. *Plant Physiol* 145:351–366
- Gorguet B, van Heusden AW, Lindhout P (2005) Parthenocarpic fruit development in tomato. *Plant Biol* 7:131–139
- Griffiths J, Murase K, Rieu I, Zentella R, Zhang ZL, Powers SJ, Gong F, Phillips AL, Hedden P, Sun TP, Thomas SG (2006) Genetic characterization and functional analysis of the *GID1* gibberellin receptors in *Arabidopsis*. *Plant Cell* 12:3399–3414
- Gusta LV, Trischuk R, Weiser CJ (2005) Plant cold acclimation: the role of abscisic acid. *J Plant Growth Regul* 24:308–318
- Horvath DP, Anderson JV, Chao WS, Foley ME (2003) Knowing when to grow: signals regulating bud dormancy. *Trends Plant Sci* 8:534–540
- Kalemba EM, Pukacka S (2007) Possible roles of LEA proteins and sHSPs in seed protection: a short review. *Biol Lett* 44:3–16
- Kojima K, Kuraishi S, Sakurai N, Fusao K (1993) Distribution of abscisic-acid in different parts of the reproductive-organs of tomato. *Sci Hort* 56:23–30
- Kushiro T, Okamoto M, Nakabayashi K, Yamagishi K, Kitamura S, Asami T, Hirai N, Koshiha T, Kamiya Y, Nambara E (2004) The *Arabidopsis* cytochrome P450 *CYP707A* encodes ABA 8'-hydroxylases: key enzymes in ABA catabolism. *EMBO J* 23:1647–1656
- Marti C, Orzaez D, Ellul P, Moreno V, Carbonell J, Granell A (2007) Silencing of *DELLA* induces facultative parthenocarpy in tomato fruits. *Plant J* 52:865–876
- Mazzucato A, Taddei AR, Soressi GP (1998) The parthenocarpic fruit (*pat*) mutant of tomato (*Lycopersicon esculentum* Mill.) sets seedless fruits and has aberrant anther and ovule development. *Development* 125:107–114
- Mezzetti B, Landi L, Pandolfini T, Spena A (2004) The *defH9-iaaM* auxin-synthesizing gene increases plant fecundity and fruit production in strawberry and raspberry. *BMC Biotechnol* 4:4–13
- Nambara E, Marion-Poll A (2005) Abscisic acid biosynthesis and catabolism. *Annu Rev Plant Biol* 56:165–185
- Okamoto M, Kuwahara A, Seo M, Kushiro T, Asami T, Hirai N, Kamiya Y, Koshiha T, Nambara E (2006) *CYP707A1* and *CYP707A2*, which encode abscisic acid 8'-hydroxylases, are indispensable for proper control of seed dormancy and germination in *Arabidopsis*. *Plant Physiol* 141:97–107
- Olimpieri I, Siligato F, Caccia R, Mariotti L, Ceccarelli N, Soressi GP, Mazzucato A (2007) Tomato fruit set driven by pollination or by the parthenocarpic fruit allele are mediated by transcriptionally regulated gibberellin biosynthesis. *Planta* 226:877–888
- Owen SJ, Abrams SR (2009) Measurement of plant hormones by liquid chromatography–mass spectrometry. In: Cutler S, Bonetta D (eds) *Plant hormones: methods and protocols*. Humana Press (in press)
- Peng YB, Cheng Z, Wang DH, Gong HQ, Xu ZH, Bai SN (2006) Preferential localization of abscisic acid in primordial and nursing cells of reproductive organs of *Arabidopsis* and cucumber. *New Phytol* 170:459–466
- Qin XQ, Zeevaert JAD (2002) Overexpression of a 9-*cis*-epoxycarotenoid dioxygenase gene in *Nicotiana plumbaginifolia* increases abscisic acid and phaseic acid levels and enhances drought tolerance. *Plant Physiol* 128:544–551
- Rebers M, Kaneta T, Kawaide H, Yamaguchi S, Yang YY, Imai R, Sekimoto H, Kamiya Y (1999) Regulation of gibberellin biosynthesis genes during flower and early fruit development of tomato. *Plant J* 17:241–250
- Rodrigo MJ, Garcia-Martinez JL (1998) Hormonal control of parthenocarpic ovary growth by the apical shoot in pea. *Plant Physiol* 116:511–518
- Ross J, O'Neill D (2001) New interactions between classical plant hormones. *Trends Plant Sci* 6:2–4

- Rotino GL, Perri E, Zottini M, Sommer H, Spena A (1997) Genetic engineering of parthenocarpic plants. *Nat Biotechnol* 15:1398–1401
- Serrani JC, Sanjuan R, Ruiz-Rivero O, Fos M, Garcia-Martinez JL (2007) Gibberellin regulation of fruit set and growth in tomato. *Plant Physiol* 145:246–257
- Shimizu-Sato S, Mori H (2001) Control of outgrowth and dormancy in axillary buds. *Plant Physiol* 127:1405–1413
- Simkin AJ, Schwartz SH, Aldridge M, Taylor MG, Klee H (2004) The tomato carotenoid cleavage dioxygenase 1 genes contribute to the formation of the flavor volatiles β -ionone pseudoionone, and geranylacetone. *Plant J* 40:882–892
- Smalle J, Kurepa J, Yang PZ, Emborg TJ, Babiychuk E, Kushnir S, Vierstra RD (2003) The pleiotropic role of the 26S proteasome subunit RPN10 in *Arabidopsis* growth and development supports a substrate-specific function in abscisic acid signaling. *Plant Cell* 15:965–980
- Stafstrom JP, Ripley BD, Devitt ML, Drake B (1998) Dormancy-associated gene expression in pea axillary buds. *Planta* 205:547–552
- Tan BC, Joseph LM, Deng WT, Liu LJ, Li QB, Cline K, McCarty DR (2003) Molecular characterization of the *Arabidopsis* 9-*cis*-epoxycarotenoid dioxygenase gene family. *Plant J* 35:44–56
- Taylor IB, Tarr AR (1984) Phenotypic interactions between abscisic acid deficient tomato mutants. *Theor Appl Gen* 68:115–119
- Thompson AJ, Jackson AC, Parker RA, Morpeth DR, Burbidge A, Taylor IB (2000) Abscisic acid biosynthesis in tomato: regulation of zeaxanthin epoxidase and 9-*cis*-epoxycarotenoid dioxygenase mRNAs by light/dark cycles, water stress and abscisic acid. *Plant Mol Biol* 42:833–845
- Thompson AJ, Thorne ET, Burbidge A, Jackson AC, Sharp RE, Taylor IB (2004) Complementation of *notabilis*, an abscisic acid-deficient mutant of tomato: importance of sequence context and utility of partial complementation. *Plant Cell Environ* 27:459–471
- Umezawa T, Okamoto M, Kushiro T, Nambara E, Oono Y, Seki M, Kobayashi M, Koshiba T, Kamiya Y, Shinozaki K (2006) *CYP707A3*, a major ABA 8'-hydroxylase involved in dehydration and rehydration response in *Arabidopsis thaliana*. *Plant J* 46:171–182
- Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paep A, Speleman F (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 3:research0034.1–research0034.11
- Vriezen WH, Feron R, Maretto F, Keijman J, Mariani C (2008) Changes in tomato ovary transcriptome demonstrate complex hormonal regulation of fruit set. *New Phytol* 177:60–76
- Wan XR, Li L (2006) Regulation of ABA level and water-stress tolerance of *Arabidopsis* by ectopic expression of a peanut 9-*cis*-epoxycarotenoid dioxygenase gene. *Biochem Biophys Res Commun* 347:1030–1038
- Wang H, Qi QG, Schorr P, Cutler AJ, Crosby WL, Fowke LC (1998) ICK1, a cyclin-dependent protein kinase inhibitor from *Arabidopsis thaliana* interacts with both Cdc2a and CycD3, and its expression is induced by abscisic acid. *Plant J* 15:501–510
- Wang H, Jones B, Li ZG, Frasse P, Delalande C, Regad F, Chaabouni S, Latche A, Pech JC, Bouzayen M (2005) The tomato Aux/IAA transcription factor IAA9 is involved in fruit development and leaf morphogenesis. *Plant Cell* 17:2676–2692
- Zeevaart JAD (1999) Abscisic acid metabolism and its regulation. In: Hooykaas MA, Hall MA, Libbenga KR (eds) *Biochemistry and molecular biology of plant hormones*. Elsevier, New York, pp 189–207