



KAR₁-induced dormancy release in *Avena fatua* caryopses involves reduction of coleorhiza sensitivity to ABA and ABA/GA_s ratio in coleorhiza and radicle

Jan Kępczyński¹ · Michał Dziurka² · Agata Wójcik¹

Received: 16 February 2024 / Accepted: 10 March 2024
© The Author(s) 2024

Abstract

Main conclusion The dormancy release by KAR₁ is associated with a reduction of coleorhiza and radicle sensitivity to ABA as well as with reduction the ABA/GA_s ratio in the coleorhiza, by a decrease content of ABA, and in the radicle, by a decrease the ABA and an increase of the GA_s contents.

Abstract Both, karrikin 1 (KAR₁) and gibberellin A₃ (GA₃), release dormancy in *Avena fatua* caryopses, resulting in the emergence of coleorhiza (CE) and radicle (RE). Moreover, KAR₁ and GA₃ stimulate CE and RE in the presence of abscisic acid (ABA), the stimulation being more effective in CE. The stimulatory effects of KAR₁ and GA₃ involve also the CE and RE rates. A similar effect was observed at KAR₁ concentrations much lower than those of GA₃. KAR₁ increased the levels of bioactive GA₅ and GA₆ in embryos and the levels of GA₁, GA₅, GA₃, GA₆ and GA₄ in radicles. The stimulatory effect of KAR₁ on germination, associated with increased levels of gibberellins (GA_s) and reduced levels of ABA in embryos, was counteracted by paclobutrazol (PAC), commonly regarded as a GA_s biosynthesis inhibitor. Consequently, KAR₁ decreased the ABA/GA_s ratio, whereas PAC, used alone or in combination with KAR₁, increased it. The ABA/GA_s ratio was reduced by KAR₁ in both coleorhiza and radicle, the effect being stronger in the latter. We present the first evidence that KAR₁-induced dormancy release requires a decreased ABA/GA_s ratio in coleorhiza and radicle. It is concluded that the dormancy-releasing effect of KAR₁ in *A. fatua* caryopses includes (i) a reduction of the coleorhiza and radicle sensitivity to ABA, and (2) a reduction of the ABA/GA_s ratio (i) in the coleorhiza, by decreasing the ABA content, and (ii) in the radicle, by decreasing the ABA and increasing the content GA_s, particularly GA₁. The results may suggest different mechanisms of dormancy release by KAR₁ in monocot and dicot seeds.

Keywords Abscisic acid · Coleorhiza · Dormancy · Gibberellins · Karrikin 1 · Radicle · Wild oat

Abbreviations

CE	Coleorhiza emergence
GA _s	Gibberellins
KAR ₁	Karrikin 1
NO	Nitric oxide
PAC	Pacllobutrazol
RE	Radicle emergence

Communicated by Dorothea Bartels.

✉ Jan Kępczyński
jan.kepczynski@usz.edu.pl


¹ Institute of Biology, University of Szczecin, Waska 13, 71-415 Szczecin, Poland

² Institute of Plant Physiology, Polish Academy of Sciences, Niezapominajek 21, 20-239 Krakow, Poland

Introduction

Harvested intact viable seeds of many plant species from both monocotyledonous and dicot plants are not able to complete germination under suitable species-specific conditions (water, temperature, air, light). Such seeds are regarded as primarily dormant (Bewley 2013). Primary dormancy established during seed maturation is very important in the wild plant life, as it facilitates survival and dispersal of the species. On the other hand, seed dormancy in agricultural crop weeds makes their control difficult. It is widely accepted that the balance between ABA and GA_s, which are synthesized in cytosol (Fig. 1), is responsible for the state of seed dormancy. ABA plays a key role in the induction and maintenance of dormancy, and GA_s participate in dormancy release and/or germination (Finkelstein et al. 2008). *Avena fatua*

Fig. 1 Diagram illustrating the metabolic pathways of ABA and GA_s. The compounds analyzed in this work are shown as grey fields. Adapted from Kępczyńska and Orłowska (2021)



is an example of an annual weed grass which infests major cereals in the world, including Poland, and produces primarily dormant caryopses which show physiological dormancy. This dormancy is expressed as the absence of complete germination at warmer temperatures at which non-dormant caryopses can germinate. Dormant caryopses of the grass in question, able to remain viable in a soil bank for several years, have been used as a model material in the study of the dormancy release mechanism in monocots (Simpson 1990; Kępczyński 2018, 2023).

Dormancy of *A. fatua* caryopses/florets can be removed by various treatments. It is released by after-ripening during dry storage of florets (Kępczyński et al. 2021) as well as by treatments involving GA₃ (Kępczyński 2018) or nitric oxide (NO) (Kępczyński et al. 2021). The NO treatment was observed to be ineffective with respect to the content of GA_s from non-13-hydroxylation and 13-hydroxylation pathways considered as bioactive (Hedden and Phillips 2000), but strongly decreased the content of ABA (Kępczyński et al. 2021). In addition, although the dormancy-releasing effect of NO was counteracted by paclobutrazol (PAC), a GA_s biosynthesis inhibitor which blocks the GA_s biosynthesis by inhibiting ent-kaurene oxidation (Desti and Amare 2021), PAC did not affect the GA_s contents, however it increased the ABA content (Kępczyński et al. 2021). The dormancy release by NO has been concluded to involve a decrease in the ABA/GA_s ratios and a reduction of caryopsis sensitivity to ABA. Other factors such as plant-derived smoke, smoke water and KAR₁ isolated from smoke are also known to stimulate seed germination in numerous species, including

dormant caryopses of *A. fatua* (Kępczyński 2023). KAR₁ has been found to be more effective in releasing dormancy than GA₃. Interaction between KAR₁ and PAC suggested that induction of germination in dormant *A. fatua* caryopses involves endogenous GA_s (Ruduś et al. 2019). As shown by Kępczyński and Van Staden (2012), although exogenous ethylene did not release caryopsis dormancy completely, to remove dormancy due to KAR₁ endogenous ethylene action was required. Also, as reported by Ruduś et al. (2019), the effect of KAR₁ was associated with non-transcriptional and transcriptional activation of ACC synthase and ACC oxidase and with modulation of the sensitivity to ethylene by regulation synthesis of ethylene receptors.

It is known that, following dormancy release, the sensu stricto germination of both monocot and dicot non-dormant seeds has been completed when the radicle or other embryonic tissue emerges through the structure covering it (Bewley et al. 2013). In the case of caryopses of, e.g., barley, *Brachypodium distachyon* and *A. fatua*, the radicle is sheathed by coleorhiza and germination is considered to involve two stages (González-Calle et al. 2015; Holloway et al. 2020; Kępczyński et al. 2021). During the first stage, the coleorhiza breaks through the surrounding structures, the second stage being associated with radicle emergence through the coleorhiza. Therefore, the sensu stricto germination is completed when the coleorhiza in monocots is punctured by radicle. So far, studies involving caryopses have used different germination criteria: the coleorhiza emergence from the surrounding tissues (Gubler et al. 2008; Kępczyński 2023), the radicle emergence through

the coleorhiza (Gendreau et al. 2008) or both criteria simultaneously (Jacobsen et al. 2013; González-Calle et al. 2015; Holloway et al. 2020; Kępczyński et al. 2021). Previously, the coleorhiza was considered as responsible for protecting the emerging radicle (Sargent and Osborne 1980), whereas at present it is recognized as playing also a key role in grass caryopsis dormancy (Millar et al. 2006; Barrero et al. 2009), including in *A. fatua* (Holloway et al. 2020). Moreover, it was postulated that ABA reduction in the coleorhiza is very important in controlling caryopsis dormancy and germination of barley (Millar et al. 2006; Barrero et al. 2009) and *A. fatua* (Kępczyński et al. 2021).

The present study was aimed at highlighting the relationship between KAR₁ and gibberellins (GA_s) as well as ABA in dormancy release in *A. fatua* caryopses. This was done by studying the effect of KAR₁ and GA₃ on the final percentage and speed of the dormant caryopses' coleorhiza and radicle emergence in the absence or in the presence of ABA. The relationship between KAR₁ and endogenous GA_s was investigated by determining contents of bioactive GA_s from the non-13-hydroxylation pathway (GA₄ and GA₇) and from the 13-hydroxylation pathway (GA₁, GA₅, GA₃ and GA₆) in embryos from caryopses germinated in the presence of KAR₁. In addition, effects of PAC, KAR₁ and PAC + KAR₁ on the GA_s and ABA contents were explored. Moreover, the GA_s level was determined in the coleorhiza and radicle of embryos isolated from caryopses treated with KAR₁. Further, changes in the ABA/GA_s ratio in embryos, coleorhiza and radicle in KAR₁-treated caryopses were investigated. The results provide new information on the role of KAR₁ in releasing caryopsis dormancy of grasses in relation to ABA/GA_s ratios in the coleorhiza and radicle before germination is completed.

Material and methods

Avena fatua L. spikelets, containing florets, collected in 2015, were dried to a constant moisture content (ca. 11%) and stored at -20 °C until used.

Germination assays

Air-dried dormant caryopses (florets without the lemma and palea) (25 in 3 replicates) were incubated in the dark at 20 °C in Petri dishes (6 cm diameter) on a single layer of filter paper (Whatman No.1) moistened with 1.5 ml distilled water, KAR₁ (10⁻⁹, 3 × 10⁻⁹, 10⁻⁸ M), GA₃ (10⁻⁵, 10⁻⁴, 10⁻³ M), ABA (10⁻⁵, 3 × 10⁻⁵, 10⁻³ M), KAR₁ + ABA, GA₃ + ABA, PAC (10⁻⁴ M), and KAR₁ (3 × 10⁻⁹ M) + PAC. Chemicals, except for PAC which was dissolved in acetone, were dissolved in water at room temperature (KAR₁) or in water heated to ca. 40 °C. Caryopses

with CE over the coat and with RE through the coleorhiza were counted every day until day 7 of germination (Fig. 2). All manipulations were performed under green light which did not affect germination. Effects of the compounds used on dormancy release were characterized by the final percentage of caryopses and Timson's index (Σ₇), calculated by summing up the CE or RE percentages over 7 days (Timson 1965).

Determination of GA_s and ABA contents

Dormant caryopses (25 in 3 replicates) were incubated in the dark at 20 °C in Petri dishes (6 cm diameter) on a single layer of filter paper (Whatman No. 1) moistened with 1.5 ml distilled water, KAR₁ (3 × 10⁻⁹ M) for 18, 24, 30 or 36 h or KAR₁ + PAC (10⁻⁴ M) for 30 h. Upon completion of incubation, the embryos (after 18, 24 and 36 h) or coleorhizae and radicles (after 24 h) were isolated, and GA_s and ABA were analyzed as described previously (Kępczyński et al. 2021, 2023). In Fig. 4 and 5 the GA_s contents in embryos incubated in water or PAC, demonstrated previously, were used. Likewise, the data of ABA contents obtained in previous studies (Kępczyński et al. 2021, 2023) were used for calculating the ABA/GA_s ratios.




Fig. 2 Germination of *A. fatua* caryopses. Photograph show ungerminated caryopsis, caryopsis with emerged coleorhiza and caryopsis with emerged radicle (germinated caryopsis)

Statistical treatment

The mean \pm standard deviation (SD) of three replicates was calculated. Significance of differences between the means was tested using one- or two-way analysis of variance (ANOVA; Statistica for Windows v. 10.0, Stat-Soft Inc., Tulsa, OK, USA). Duncan's multiple range test was used to identify significantly different ($P \leq 0.05$) mean values.

Results

Effects of KAR₁ and GA₃ on the emergence of coleorhiza and radicles of *A. fatua* caryopses in the absence or in the presence of ABA

Both, coleorhizae and radicles, of the primarily dormant *A. fatua* caryopses incubated in water or ABA solutions were almost or totally unable to emerge (0–5%) (Fig. 3). KAR₁ stimulated the coleorhiza emergence (CE) in all the concentrations used (Fig. 3A). The highest and similar levels

of stimulation were found after KAR₁ was applied at concentrations of 3×10^{-9} and 10^{-8} M; ca. 90% of caryopses observed show CE. KAR₁ applied at concentrations of 10^{-9} – 10^{-8} M produced a similar CE despite the presence of ABA at 10^{-5} M (ca. 80%). Application of KAR₁ at 3×10^{-9} or 10^{-8} M markedly enhanced CE in the presence of 3×10^{-5} and 10^{-4} M ABA; ca. 50% of coleorhizae emerged. KAR₁ used 3×10^{-9} and 10^{-8} M resulted in ca. 90% RE (Fig. 3B). The highest antagonizing KAR₁ effect was found when ABA was used at the lowest concentration (10^{-5} M). KAR₁ at all concentrations did not affect germination in the presence of ABA at the highest concentration of 10^{-4} M (ca. 3%). KAR₁ at 10^{-8} M increased Timson's index (Σ_7) by the factor of 21 and 23 in the coleorhiza and radicle, respectively (Table 1A). In the presence of ABA at 10^{-5} M, KAR₁ at 10^{-8} M did enhance Timson's index in both the coleorhiza and radicle by the factor of 39 and 31, respectively.

Likewise, GA₃ stimulated CE and RE at all the concentrations used (Fig. 3C, D). The highest stimulatory effect on CE and RE of dormant caryopses was observed at GA₃ concentrations of 10^{-4} and 10^{-3} M; almost all the




Fig. 3 Effects of KAR₁ and GA₃ on the emergence of coleorhiza (A, C) and radicle (B, D) of *A. fatua* in the absence or in the presence of ABA after 7 days of germination. One way ANOVA with Duncan's

post hoc test was used to test for significance of differences. Means denoted by different letters differ significantly ($P < 0.05$, $n = 3$)

Table 1 Effects of KAR₁ (A) and GA₃ (B) on the speed (Timson’s index) of coleorhiza and radicle emergence in *A. fatua* in the presence of ABA

	ABA, M			
	Coleorhiza		Radicle	
	0	10 ⁻⁵	0	10 ⁻⁵
(A) KAR₁, M				
0	21.3 ± 20.1a	10.7 ± 12.2a	17.3 ± 16.2a	10.7 ± 12.2a
10 ⁻⁹	373.3 ± 56.6d	289.3 ± 35.9b	336.0 ± 52.9c	216.0 ± 32.7b
10 ⁻⁸	448.0 ± 32.7d	414.7 ± 46.0 cd	405.3 ± 38.0d	337.3 ± 23.1c
(B) GA₃, M				
0	21.3 ± 20.1a	10.7 ± 12.2a	17.3 ± 16.2a	10.7 ± 12.2a
10 ⁻⁵	381.3 ± 44.2c	244.0 ± 46.1b	353.3 ± 28.1d	121.3 ± 15.1b
10 ⁻⁴	500.0 ± 41.8d	360.0 ± 32.0c	465.3 ± 47.7e	216.0 ± 38.2c

One way ANOVA with Duncan’s post hoc test was used to test for significance of differences. Means denoted by different letters differ significantly ($P < 0.05, n = 3$)

caryopses (90–100%) showed CE and RE. When used at the concentrations mentioned, GA₃ resulted in 90% of the caryopses showing CE, despite the presence of ABA at 10⁻⁵-3 × 10⁻⁵ M (Fig. 3C). When used at the two highest concentrations in the presence of 10⁻⁵ M ABA, GA₃ produced an almost complete RE (90–95%) (Fig. 3D). The stimulatory effect of GA₃ was not evident in the presence of 10⁻⁴ M ABA (ca. 3%). GA₃ at 10⁻⁴ M increased Timson’s index calculated for the coleorhiza and radicle by the factor of 24 and 27, respectively (Table 1B). When used together with ABA, GA₃ resulted in a Timson’s index increase by the factor of 34 and 20 in the coleorhiza and radicle, respectively.

Effects of KAR₁ on GA_s contents in embryos from germinated caryopses

Contents of GA_s from non-13-hydroxylation, GA₄ and GA₇, and 13-hydroxylation pathway, GA₁, GA₅, GA₃ and GA₆ in embryos isolated from caryopses germinated for various periods of time in water or KAR₁ solutions was determined (Fig. 4). After 18 h of germination, KAR₁ did not affect the contents of GA₁, GA₄ and GA₇ (not shown) and decreased the GA₅ content, whereas the contents of GA₃ and GA₆ increased (ca. 2 times). When the period of germination in the presence of KAR₁ was extended to 36 h, the GA₅ and GA₆ contents in embryos were higher than in embryos from caryopses germinated in water; the increase was 1.5–2.2 times.

Effects of PAC, KAR₁ and PAC + KAR₁ on radicle emergence as well as ABA and GA_s contents and ABA/GA_s ratio in embryos

Radicles of dormant caryopses were almost unable to emerge when kept in water or in PAC solutions; ca. 2–5% of the radicles emerged (Fig. 5A). In contrast, KAR₁-treated caryopses germinated nearly completely (ca. 90% showed




Fig. 4 Effects of KAR₁ on the GA_s contents in embryos of *A. fatua* caryopses after 18 and 36 h of germination. After 36 h, coleorhiza emerged in 49% of caryopses. Vertical bars indicate ±SD. One way ANOVA with Duncan’s post hoc test was used to test for significance of differences. Means denoted by different letters differ significantly ($P < 0.05, n = 3$)

emerged radicles). PAC applied in combination with KAR₁ almost totally counteracted the stimulatory effect of the latter; ca. 6% of the caryopses showed emerging radicles. Contents of ABA, GA₅, GA₃ and GA₆ (the 13-hydroxylation pathway) were determined in embryos isolated from caryopses germinated for 30 h in water, in the presence of PAC, KAR₁ and the combination of PAC and KAR₁. PAC was found to enhance, by 30%, the ABA content compared to embryos from water-germinated caryopses (Fig. 5B). In contrast, KAR₁ applied alone decreased (by 60%) the ABA content. When PAC was applied simultaneously with KAR₁, the ABA level was lower (by 30%) than that in embryos from water-germinated caryopses, but was higher than that in KAR₁-treated ones. PAC did not affect the GA_s content,




Fig. 5 Effects of KAR₁, PAC and KAR₁+PAC on the final radicle emergence of *A. fatua* caryopses (A) as well as on GA_s (GA₅, GA₃, GA₆) (C) and ABA (B) contents in embryos after 30 h of caryopsis germination (presented as % of control content) and the ABA/GA_s

ratios (D) in embryos. One way ANOVA with Duncan's post hoc test was used to test for significance of differences. Means denoted by different letters differ significantly ($P < 0.05$, $n = 3$)

whereas KAR₁ increased it by 40% (Fig. 5C). The GA_s content was similar in embryos from caryopses incubated in water, in the presence of PAC or KAR₁ applied simultaneously with PAC. The ABA/GA_s ratios showed large treatment-dependent differences; the ratio of 1.6 to 2.9 was associated with the absence of germination, the ratio of 0.6 being characteristic of germination resulting from KAR₁ treatment (Fig. 5D).

Effects of KAR₁ on the GA_s content and ABA/GA_s ratio in coleorhiza and radicle

KAR₁ did not affect the GA₉, GA₄, GA₇, GA₂₀, GA₁, GA₈, GA₅ and GA₆ contents in the coleorhiza of caryopses germinated for 24 h, but decreased the GA₃ content (Fig. 6A). The content of all GA_s except for GA₉ and GA₇ was increased in radicles of KAR₁-treated caryopses, the highest effect being seen in GA₁; the GA₁ content increased by the factor of ca. 6. The total content of all the bioactive GA_s from both GA_s biosynthesis pathways was similar in the coleorhiza from untreated and KAR₁-treated caryopses (Fig. 6B). When

caryopses were treated with KAR₁, the bioactive GA_s content in the radicles was 1.7 times higher than that in the radicles of untreated caryopses. The ABA/GA_s ratios in the coleorhiza and radicles showed KAR₁ to decrease the ratio by the factor 1.5 and 3, respectively (Fig. 6C).

Discussion

Relationship between exogenous KAR₁ or GA₃ and exogenous ABA

The results presented confirm the findings, reported by previous studies, that KAR₁ applied at very low concentrations is able to remove dormancy in caryopses of *A. fatua* (Kępczyński 2018, 2023; Fig. 3), thus enabling almost all the caryopses to complete germination. The time course of germination in grasses, e.g. *Brachypodium distachyon* (González-Calle et al. 2015) and *A. fatua* (Kępczyński et al. 2021) involves two stages: the coleorhiza emergence (CE) at the first stage and the radicle emergence




Fig. 6 Effects of KAR₁ on the GA_s (GA₉, GA₄, GA₇, GA₂₀, GA₁, GA₈, GA₅) (A) contents, total GA_s content (B) and the ABA/GA_s ratios (C) in coleorhiza and radicle after 24 h germination of *A. fatua*

caryopses. One way ANOVA with Duncan’s post hoc test was used to test for significance of differences. Means denoted by different letters differ significantly ($P < 0.05$, $n = 3$)

(RE) at the second, either stage or the two combined being used as a criterion for caryopsis germination (Gubler et al. 2008; Gendreau et al. 2008; Jacobsen et al. 2013; González-Calle et al. 2015). The coleorhiza is recognized as being mainly responsible for dormancy control in caryopses of barley (Barrero et al. 2009), *B. distachyon* (González-Calle et al. 2015) and *A. fatua* (Holloway et al. 2020; Kępczyński et al. 2021), and plays a role similar to that of the endosperm in dicot seeds. Radicle emergence through the endosperm or coleorhiza is regarded as a completed germination (Bewley et al. 2013). KAR₁ turned out to be very effective in inducing CE and RE of dormant caryopses, not only when applied alone (Kępczyński et al. 2021; Fig. 3A, B), but also in the presence of ABA (Fig. 3A, B), a compound which inhibits germination of non-dormant *A. fatua* caryopses (Kępczyński et al. 2021). However, in an

experiment with *Arabidopsis* seeds, KAR₁ was unable to overcome ABA inhibition of seed germination (Nelson et al. 2009). Like in previous studies, GA₃, similarly to KAR₁, released dormancy in *A. fatua* caryopses, but much higher concentrations were needed (Kępczyński 2018; Fig. 3C, D). This is consistent with findings from other seeds, e.g. those of *Arabidopsis* and *Brassica tournefortii*, more sensitive to KAR₁ than to gibberellin (Daws et al. 2007; Stevens et al. 2007; Nelson et al. 2009). In the presence of ABA, a higher concentration of GA₃ than KAR₁ was required for dormant *A. fatua* caryopses to germinate (Fig. 3). Antagonistic effects of both KAR₁ and GA₃ towards ABA were more evident in CE than in RE. This indicates that, in the presence of ABA, the coleorhiza is more sensitive to KAR₁ and GA₃ than the radicle, which is consistent with previous results showing a stronger response of dormant caryopses to KAR₁

in the coleorhiza than in the radicle (Kępczyński et al. 2021). Not only did both, KAR₁ and GA₃, increase the percentage germination in the presence of ABA, but they also accelerated caryopsis germination, even when the final germination was not affected (Table 1). Taking into account the final germination and the germination speed, it can be concluded that KAR₁ and GA₃ reduced the sensitivity of both the coleorhiza and radicle to ABA; however, the effect was more effective in the coleorhiza. It was demonstrated earlier (Kępczyński et al. 2023) that NO, another dormancy-releasing agent, was able to reduce the sensitivity of *A. fatua* caryopses to ABA.

The relationship between exogenous KAR₁ and endogenous GA_s and ABA

Embryos

A. fatua embryos showed the presence of GA_s from the non-13-hydroxylation and 13-hydroxylation pathways (Kępczyński et al. 2021; Fig. 4), recognized as bioactive (Hedden 2016). The dormancy releasing effect of KAR₁ was associated only with an increase in the 13-hydroxylation pathway GA_s (Fig. 4), in contrast to the dormancy releasing effect of NO, which did not affect the contents of GA_s from both pathways in the embryos (Kępczyński et al. 2023). This suggests a difference between these compounds in the dormancy release mechanisms. The stimulating effect of KAR₁ and NO on germination of dormant caryopses was strongly counteracted by PAC (Kępczyński 2018; Ruduś et al. 2019; Kępczyński et al. 2023; Fig. 5A) regarded as a GA_s biosynthesis inhibitor (Desta and Amare 2021), which suggests a requirement for endogenous GA_s in response to both compounds. PAC applied alone to caryopses was reported to not affect GA_s content in embryos (Kępczyński et al. 2023; Fig. 5C), nor did it affect the GA_s content when caryopses were treated with NO (Kępczyński et al. 2023). However, PAC reduced the GA_s content when applied simultaneously with KAR₁, the decline being associated with a reduced stimulatory effect of KAR₁ on germination of dormant caryopses (Fig. 5A, C). It was also reported earlier that PAC, in addition to influencing the GA_s content, can increase the content of ABA by increasing its synthesis and/or inhibiting its catabolism (Yamaguchi et al. 2007; Desta and Amare 2021). The ABA content, which was reduced by KAR₁ (Kępczyński et al. 2021; Fig. 5B) as a result of degradation to phaseic acid (Kępczyński 2023), and by NO (Kępczyński et al. 2023), was increased due to PAC being used alone (Kępczyński et al. 2023) or simultaneously with KAR₁ (Fig. 5B). Taken together, this might confirm different mechanisms involved in dormancy release in *A. fatua* caryopses by KAR and NO.

It is widely accepted that the ABA/GA_s ratio is mainly responsible for the dormancy level and seed germination (Rodriguez et al. 2015; Tuan et al. 2018). A high ABA/GA_s ratio associated with a high ABA and low GA_s contents is responsible for dormancy. A low ratio, associated with low ABA and high GA_s levels, allows germination. ABA plays a crucial role in the induction and maintenance of seed dormancy (Rodriguez-Gacio et al. 2009). GA_s act antagonistically to ABA, promote dormancy release and are required for germination (Kucera et al. 2005; Bewley et al. 2013). Taking into account the ABA/GA_s ratio in *A. fatua* embryos, it can be seen that the KAR₁-effected dormancy release is associated with a marked decrease in the ratio (Fig. 5D). Its high level in embryos from caryopses germinated in water or in the presence of PAC or PAC + KAR₁ is presumably responsible for inability of the caryopses to transit from dormancy to germination. Reduction of the ABA/GA_s ratio by KAR₁ involves a decrease in the ABA content and a simultaneous increase in GA_s contents (Fig. 5B, C), while in the case of NO the effect is associated only with a decrease in the ABA content (Kępczyński et al. 2023). Interestingly, the ABA levels in *Arabidopsis* seeds were not affected by KAR₁ prior to radicle emergence, although KAR₁ did effectively remove dormancy (Nelson et al. 2009). Moreover, KAR₁ only slightly increased the GA₄ level. It has also been shown that KAR₁ can delay germination of non-dormant soybean seeds by a change in the ABA/GA₄ ratio due to the biosynthesis of ABA and GA₄ being enhanced and impaired, respectively (Meng et al. 2016).

Coleorhiza

Since the coleorhiza is considered to be responsible for the dormancy state of barley (Millar et al. 2006; Barrero et al. 2009) and *A. fatua* (Holloway et al. 2020) caryopses, GA_s from the non-13-hydroxylation and 13-hydroxylation pathways were determined not only in embryos, but also in the coleorhiza. KAR₁ was found to not increase the contents of bioactive GA_s from both pathways (Fig. 6A, B). Likewise, the GA₉, substrate for GA₄ and GA₇, as well as GA₂₀, the substrate for GA₁ and GA₅, were not affected by KAR₁. KAR₁ was earlier demonstrated to be capable of reducing the ABA content in the coleorhiza (Kępczyński et al. 2021). Taking these results into account, it can be concluded that the KAR₁-effected control of the ABA but not GA_s contents in the coleorhiza plays a vital role in releasing caryopsis dormancy. It has been previously postulated that endogenous GA_s in the coleorhiza are not involved in dormancy release of barley caryopses since expression of the genes involved in the GA_s synthesis (*KAURENOIC ACID OXIDASE1*) was upregulated only in the coleorhiza of dormant caryopses, and expression of *GIBBERELLIN 2-OXIDASE1* responsible

for GA_s inactivation was upregulated in coleorhiza of non-dormant caryopses (Barrero et al. 2009). Thus, ABA in the coleorhiza plays an essential role in the control of caryopsis dormancy in barley (Barrero et al. 2009) and *A. fatua* (Holloway et al. 2020; Kępczyński et al. 2021), GA_s not being involved. The ABA/GA_s ratio showed that, like in the embryos, KAR₁ reduces the ratio in the coleorhiza (Fig. 6C), but in contrast to embryos the effect is associated only with a reduction in the ABA content, which probably could allow an increase in the activity of enzymes responsible for weakening the coleorhiza, as was postulated for barley (Barrero et al. 2009), *Brachypodium distachyon* (Gonzalez-Calle et al. 2015) and *A. fatua* (Holloway et al. 2020).

Radicle

KAR₁ increased the total content of bioactive GA_s from both pathways (Fig. 6B), which was related to an increase in the content of four gibberellins (GA₁, GA₅, GA₃ and GA₆) from the 13-hydroxylation pathway and one, GA₄, from the non-13-hydroxylation pathway (Fig. 6A). The largest impact of KAR₁ was recorded in GA₁ the content of which was increased by ca. 470%, suggesting that it is mainly GA₁ that is required for radicle growth. The contents of GA₄, GA₅, GA₃ and GA₆ increased by 50–70%. Moreover, GA₂₀—a substrate for GA₁ biosynthesis—was markedly, by ca. 40%, increased by KAR₁, which may suggest the potential to synthesize GA₁, enabling a further increase of its content and, perhaps, also the remaining GA_s of this pathway. The increase of the content of GA₈, a GA₁ catabolite, by KAR₁ may indicate the regulation of the bioactive gibberellin concentrations also through deactivation. The presented data could indicate that GA_s from the 13-hydroxylation pathway play the main role in radicle growth; GA₄ from the non-13-hydroxylation pathway seems to be less important (Fig. 6A). In contrast to KAR₁ effect on the GA_s content, KAR₁ reduced the content of ABA (Kępczyński et al. 2021), a GA_s antagonist, considered as a positive regulator of dormancy and negatively affecting seed germination (Hilhorst 1995). Thus, the balance between ABA and GA_s was altered by KAR₁, like in the coleorhiza. It is worth emphasizing that the reduction of the ABA/GA_s ratio by KAR₁ in the coleorhiza is associated only with a reduction in the ABA content, whereas in the radicle, in addition to the reduction of the ABA content, there is also an increase in the GA_s content. Considering the antagonistic effects of ABA and GA_s, a change in the ABA/GA_s ratio possibly enables the radicle to grow and break through the coleorhiza, allowing germination to be completed. Thus, KAR₁-associated dormancy release, involves—in addition to a reduction of the ABA/GA_s ratio in the coleorhiza—probably also an increase of the radicle's expansive force by a decrease

of the ABA/GA_s ratio, making it easier for the radicle to penetrate the coleorhiza. It was earlier proposed that RE may depend not only on weakening of the coleorhiza, but also on the expansive force of the radicle (Barrero et al. 2009). Our previous study (Kępczyński et al. 2021) allowed to suggest that the inability of dormant caryopses to complete germination, probably mainly due to the endogenous ABA concentration being too high, might involve inhibition of the cell-cycle activation.

To summarize, like in previous studies (Kępczyński 2018, 2023), KAR₁ and GA₃, very actively induced dormancy release in *A. fatua* caryopses. Both compounds reduced the sensitivity of the coleorhiza and radicle to exogenous ABA. The dormancy releasing KAR₁ effect was associated with a decrease in the ABA/GA_s ratios in embryos, coleorhiza and radicle before germination was completed. The mechanism of dormancy release by KAR₁ is related to a reduction of the coleorhiza sensitivity to ABA and a decrease in the ABA/GA_s ratios in the coleorhiza, regarded as playing a key role in maintaining caryopsis dormancy, by a decrease in the ABA level. A KAR₁-induced reduction of the radicle ABA/GA_s ratio, involving a large increase in the content of bioactive GA_s, particularly GA₁, as well as a reduction of the ABA level are probably required for the radicle to grow and break through the coleorhiza. Thus, caryopsis dormancy release under the influence of KAR₁ requires a reduction of the ABA/GA_s ratio in the coleorhiza and also in the radicle, which probably allows the radicle expansion force to increase, thus facilitating the coleorhiza puncture. It has been shown for the first time here that a reduction of the ABA/GA_s ratio is necessary for dormancy release by KAR₁. The results presented may also indicate that the mode of KAR₁-effected dormancy release differs between monocot and dicot seeds.

Acknowledgements We would like to express our gratitude to Prof. Ewa Kępczyńska for reading and commenting on the manuscript. The authors thank Emilia Jesionowska, M.Sc. for technical assistance. We are indebted to Dr Teresa Radziejewska for linguistic assistance.

Author contributions JK: conceived and designed the research, interpreted results and wrote the manuscript; AW: conducted the physiological experiments; MD: carried out the GC analysis. All authors read, reviewed and approved the manuscript.

Data availability The data sets generated and/or analysed during the current study are available from the corresponding author on a reasonable request.

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing,

adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Barrero JM, Talbot MJ, White RG, Jacobsen JV, Gubler F (2009) Anatomical and transcriptomic studies of the coleorhiza reveal the importance of this tissue in regulating dormancy in barley. *Plant Physiol* 150:1006–1021
- Bewley JD, Bradford KJ, Hilhorst HWM, Nonogaki H (2013) Germination. In: Bewley JD, Bradford KJ, Hilhorst HWM, Nonogaki H (eds) *Physiology of development, germination and dormancy*. Springer, New York, pp 133–181
- Daws MI, Davies J, Pritchard HW, Brown NAC, Van Staden J (2007) Butenolide from plant-derived smoke enhances germination and seedling growth of arable weed species. *Plant Growth Regul* 51:73–82
- Desta B, Amare G (2021) Paclobutrazol as a plant growth regulator. *Chem Biol Technol Agric* 8:1
- Finkelstein R, Reeves W, Ariizumi T, Steber C (2008) Molecular aspects of seed dormancy. *Annu Rev Plant Biol* 59:387–415
- Gendreau E, Romaniello S, Barad S, Leymarie J, Benech-Arnold R, Corbineau F (2008) Regulation of cell cycle activity in the embryo of barley seeds during germination as related to grain hydration. *J Exp Bot* 59:203–212
- González-Calle V, Barrero-Sicilia C, Carbonero P, Iglesias-Fernández R (2015) Mannans and endo- β -mannanases (MAN) in *Brachypodium distachyon*: expression profiling and possible role of the *BdMAN* genes during coleorhiza-limited seed germination. *J Exp Bot* 66:3753–3764
- Gubler F, Hughes T, Waterhouse P, Jacobsen J (2008) Regulation of dormancy in barley by blue light and after-ripening: effects on abscisic acid and gibberellin metabolism. *Plant Physiol* 147:886–896
- Hedden P (2016) Gibberellin biosynthesis in higher plants. *Annu Plant Rev* 49:73–94
- Hilhorst WM (1995) A critical update on seed dormancy. *Seed Sci Res* 5:61–73
- Holloway T, Steinbrecher T, Perez M, Seville A, Stock D, Nakabashi K, Leubner-Metzger G (2020) Coleorhiza-enforced seed dormancy: a novel mechanism to control germination in grasses. *New Phytol* 229:2179–2191
- Jacobsen JV, Barrero JM, Hughes T, Julkowska M, Taylor JM, Xu Q, Gubler F (2013) Roles for blue light, jasmonate and nitric oxide in the regulation of dormancy and germination in wheat grain (*Triticum aestivum* L.). *Planta* 238:121–138
- Kępczyńska E, Orłowska A (2021) Profiles of endogenous ABA, bioactive GAs, IAA and their metabolites in *Medicago truncatula* Gaertn. non-embryonic and embryonic tissues during induction phase in relation to somatic embryo formation. *Planta* 253:67
- Kępczyński J (2018) Induction of agricultural weed seed germination by smoke and smoke-derived karrikin (KAR₁), with a particular reference to *Avena fatua* L. *Acta Physiol Plant* 40:87
- Kępczyński J (2023) Induction of dormancy release in agricultural weed seeds by plant-derived smoke and smoke-derived Karrikin1 (KAR1). A relationship with plant hormones. In: Soumya M, Aftab T (eds) *Strigolactones karrikins and alkaloids in plants*. Taylor & Francis, Boca Raton, pp 225–240
- Kępczyński J, Van Staden J (2012) Interaction of karrikinolide and ethylene in controlling germination of dormant *Avena fatua* L. caryopses. *Plant Growth Regul* 67:185–190
- Kępczyński J, Wójcik A, Dziurka M (2021) *Avena fatua* caryopsis dormancy release is associated with changes in KAR₁ and ABA sensitivity as well as with ABA reduction in coleorhiza and radicle. *Planta* 253:52
- Kępczyński J, Wójcik A, Dziurka M (2023) NO-mediated dormancy release of *Avena fatua* caryopses is associated with decrease in abscisic acid sensitivity, content and ABA/GA_s ratios. *Planta* 257:101
- Kucera B, Cohn MA, Leubner-Metzger G (2005) Plant hormone interactions during seed dormancy release and germination. *Seed Sci Res* 15:281–307
- Meng Y, Chen F, Shuai H, Luo X, Ding J et al (2016) Karrikins delay soybean seed germination by mediating abscisic acid and gibberellin biogenesis under shaded conditions. *Sci Rep* 6:22073
- Millar AA, Jacobsen JV, Ross JJ, Helliwell CA, Poole AT, Scofield G, Reid JB, Gubler F (2006) Seed dormancy and ABA metabolism in Arabidopsis and barley: the role of ABA 80—hydroxylase. *Plant J* 45:942–954
- Nelson DC, Riseborough JA, Flematti GR, Stevens J, Ghisalberti EL, Dixon KW, Smith SM (2009) Karrikins discovered in smoke trigger Arabidopsis seed germination by a mechanism requiring gibberellin acid synthesis and light. *Plant Physiol* 149:863–873
- Rodríguez MV, Barrero JM, Corbineau F, Gubler F, Benech-Arnold RL (2015) Dormancy in cereals (not too much, not so little): about the mechanisms behind this trait. *Seed Sci Res* 25:99–119
- Rodríguez-Gacio MC, Matilla-Vázquez MA, Matilla AJ (2009) Seed dormancy and ABA signaling. *Plant Signal Behav* 4:1035–1048
- Ruduś I, Cembrowska-Lech D, Jaworska A, Kępczyński J (2019) Involvement of ethylene biosynthesis and perception during germination of dormant *Avena fatua* L. caryopses induced by KAR₁ or GA₃. *Planta* 249:719–738
- Sargent JA, Osborne DJ (1980) A comparative study of the fine structure of coleorhiza and root during the early hours of germination of rye embryos. *Protoplasma* 104:91–103
- Simpson GM (1990) *Seed dormancy in grasses*. Cambridge University Press, Cambridge
- Stevens JC, Merritt DJ, Flematti GR, Ghisalberti EL, Dixon KW (2007) Seed germination of agricultural weeds is promoted by the butenolide 3-methyl-2H-furo[2,3-c]pyran-2-one under laboratory and field conditions. *Plant Soil* 298:113–124
- Timson J (1965) New method of recording germination data. *Nature* 207:216–217
- Tuan PA, Kumar R, Rehal PK, Toora PK, Ayle BT (2018) Molecular mechanism underlying abscisic acid/gibberellin balance in the control of seed dormancy and germination in cereals. *Front Plant Sci* 9:668
- Yamaguchi S, Kamiya Y, Nambara E (2007) Regulation of ABA and GA levels during seed development and germination in Arabidopsis. In: Bradford KJ, Nonogaki H (eds) *Seed development, dormancy and germination*. Wiley-Blackwell, Hoboken, pp 224–247

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.