

Burkhard Helpap

Nonepithelial neoplasms of the urinary bladder

Published online: 26 January 2002
© Springer-Verlag 2002

Virchows Arch (2001) 439:497–503

The credit line under Table 2 was omitted, so Table 2 is reprinted here in its entirety.

Table 2 Five-year survival rates in sarcomas of the bladder

Tumor type	Grade 1	Grade 2	Grade 3	Total
Malignant fibrous histiocytoma	89.8% (33)	76.5% (160)	48.1% (156)	349
Leiomyosarcoma	92.9% (16)	66.6% (65)	44.7% (67)	148
Malignant schwannoma	77.8% (5)	56.1% (26)	52.1% (41)	72
Rhabdomyosarcoma	0% (0)	74.9% (15)	42.1% (45)	60
Undifferentiated sarcomas	0% (0)	66.9% (43)	40.3% (97)	140

Reprinted with permission from J.M. Coindre, Verh Dtsch Ges Pathol 82:59–63 (1998) (modified)

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s004280100479>

B. Helpap (✉)
Department of Pathology, Hegau-Klinikum Singen,
University of Freiburg, Germany
e-mail: pathologie@hegau-klinikum.de
Tel.: +49-7731-892100, Fax: +49-7731-892105