Correlations between reduced expression of the metastasis suppressor gene KAI-1 and accumulation of p53 in uterine carcinomas and sarcomas

**Juliane Briesse · Heinrich M. Schulte · Maria Sajin ·
Christoph Bamberger · Katja Redlin ·
Karin Milde-Langosch · Thomas Löning ·
Ana-Maria Bamberger**

Published online: 5 July 2008
© Springer-Verlag 2008

The author's affiliations were given incorrectly. The correct affiliations are given here.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00428-008-0608-7>

J. Briesse · C. Bamberger · A.-M. Bamberger
Section on Endocrinology and Metabolism of Ageing,
University Clinic Hamburg-Eppendorf,
Hamburg, Germany

J. Briesse (✉)
Department of Pathology and Tumor Biology,
University of Leeds, St. James's Hospital,
Beckett Street,
LS9 7TF Leeds, UK
e-mail: j.briesse@leeds.ac.uk

H. M. Schulte · K. Redlin
Endokrinologikum,
Hamburg, Germany

M. Sajin
Department of Morphopathology,
University of Bucharest,
Bucharest, Romania

K. Milde-Langosch
Department of Gynecology,
University Clinic Hamburg-Eppendorf,
Hamburg, Germany

T. Löning
Department of Pathology, Albertinen-Hospital,
Hamburg, Germany