Correction to: The effect of the peritoneal tumor microenvironment on invasion of peritoneal metastases of high-grade serous ovarian cancer and the impact of neoadjuvant chemotherapy

J. O. A. M. van Baal¹ · C. A. R. Lok¹ · E. S. Jordanova¹ · H. Horlings² · W. J. van Driel¹ · F. C. Amant¹ · K. K. Van de Vijver^{2,3}

Published online: 2 December 2020

© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: Virchows Archiv (2020) 477:535–544
<https://doi.org/10.1007/s00428-020-02795-8>

The authors regret that in the Abstract section of the original published version of the above article contained errors. They requested that these be noted [Bold text used to highlight problem area].

In the abstract originally written as “After NACT, **increased ieCD3+ cells** were shown, compared to no-NACT (p=0.044).” However, this should be “After NACT, **decreased stromal (s)CD3+ cells** were shown, compared to no-NACT (p=0.044).”

Furthermore, “NACT **increases abundance of ieCD3+ cells** in peritoneal metastases or HGSOC.” should be adjusted to “NACT **decreases abundance of sCD3+ cells** in peritoneal metastases or HGSOC.”

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00428-020-02795-8>

✉ J. O. A. M. van Baal
j.v.baal@nki.nl

¹ Department of Gynecology, Center for Gynecologic Oncology
Amsterdam, P.O. Box 90203, Amsterdam 1006, BE,
The Netherlands

² Division of Diagnostic Oncology & Molecular Pathology,
The Netherlands Cancer Institute, Antoni van Leeuwenhoek
Hospital, Amsterdam, The Netherlands

³ Department of Pathology, Ghent University Hospital, Cancer
Research Institute Ghent (CRIG), Ghent, Belgium