Correction to: Quantification of neurons in the hippocampal formation of chimpanzees: comparison to rhesus monkeys and humans

Christina N. Rogers Flattery¹ · Rebecca F. Rosen² · Aaron S. Farberg³ · Jeromy M. Dooyema³ · Patrick R. Hof⁴ · Chet C. Sherwood⁵ · Lary C. Walker^{3,6} · Todd M. Preuss^{3,7}

Published online: 29 October 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to:
Brain Structure and Function (2020) 225:2521–2531
<https://doi.org/10.1007/s00429-020-02139-x>

The original version of this article contained a mistake in Figs. 3 and 4: the colors corresponding to CA1 and the subiculum were reversed. The correct versions of Figs. 3 and 4 are provided here.

The original article can be found online at <https://doi.org/10.1007/s00429-020-02139-x>.

✉ Christina N. Rogers Flattery
cnrogers@fas.harvard.edu

¹ Department of Human Evolutionary Biology, Harvard University, 11 Divinity Ave, Cambridge, MA 02138, USA

² National Institute of Mental Health, National Institutes of Health, Bethesda, MD 20814, USA

³ Yerkes National Primate Research Center, Emory University, Atlanta, GA 30329, USA

⁴ Nash Family Department of Neuroscience and Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

⁵ Department of Anthropology and Center for the Advanced Study of Human Paleobiology, The George Washington University, Washington, DC 20052, USA

⁶ Department of Neurology, Emory University School of Medicine, Atlanta, GA 30322, USA

⁷ Department of Pathology and Laboratory Medicine, Emory University School of Medicine, Atlanta, GA 30322, USA

Fig. 3 Schematic representation of the average proportion of neurons, volume of each subregion, and density of neurons in 5 subdivisions of the hippocampal formation in 6 adult chimpanzees

Fig. 4 Schematic representation of the proportions of neurons in five hippocampal subregions of adult rhesus monkeys (Keuker et al. 2003), chimpanzees (present study) and humans (mean values of West 1993; West and Gundersen 1990; and Šimić et al. 1997) relative

to total hippocampal number. The proportion of neurons in CA1 and subiculum increases, whereas the proportion of GCL (granule cell layer of the dentate gyrus) neurons decreases, from rhesus monkeys to chimpanzees to humans

References

- Keuker JJ, Luiten PG, Fuchs E (2003) Preservation of hippocampal neuron numbers in aged rhesus monkeys. *Neurobiol Aging* 24(1):157–165
- Šimić G, Kostović I, Winblad B, Bogdanović N (1997) Volume and number of neurons of the human hippocampal formation in normal aging and Alzheimer's disease. *J Comp Neurol* 379(4):482–494
- West MJ (1993) Regionally specific loss of neurons in the aging human hippocampus. *Neurobiol Aging* 14(4):287–293

- West MJ, Gundersen HJG (1990) Unbiased stereological estimation of the number of neurons in the human hippocampus. *J Comp Neurol* 296(1):1–22

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.