CORRESPONDENCE

Left ventricular dysfunction and subclinical atherosclerosis in children with classic congenital adrenal hyperplasia: a single-center study from Upper Egypt

Kotb Abbass Metwalley¹

Received: 20 November 2015 / Accepted: 27 November 2015 / Published online: 10 December 2015
© Springer-Verlag Berlin Heidelberg 2015

Reply

"We agree with Dr. Ozkan et al. that minimization of variation or error in carotid intima-media thickness (cIMT) measurement is crucial for both research and clinical applications. Technical matters like dynamic range, output and receiver gains, transducer frequency, and number of measurements were taken into consider-

ation while measuring the cIMT of our studied cases; we did not mention these details because of brevity and because they are well known to pediatric cardiologists. All measurements of cIMT of our studied cases were done at end diastole gated by R wave on electrocardiography."

Regards,

Kotb Abbass Metwalley, M.D.

Communicated by Beat Steinmann

This is a correspondence to an article which can be found at <http://dx.doi.org/10.1007/s00431-015-2677-3>

✉ Kotb Abbass Metwalley
kotb72@gmail.com

¹ Pediatric Endocrinology Unit, Department of Pediatrics, Faculty of Medicine, Assiut University, B.O. 71111, Assiut, Egypt