

Improved characterisation of solitary solid pancreatic tumours using contrast enhanced transabdominal ultrasound

C. F. Dietrich · B. Braden · M. Hocke · M. Ott · A. Ignee

Published online: 1 December 2007
© Springer-Verlag 2007

Erratum to: J Cancer Res Clin Oncol DOI 10.1007/s00432-007-0326-6

Owing to a systemic error, some of the data in Table 1 were incorrect. The text referring to Table 1 presents the correct values. The revised version of Table 1 is given here.

The online version of the original article can be found under
doi:10.1007/s00432-007-0326-6.

C. F. Dietrich (✉) · A. Ignee
Medical and Imaging Department,
Caritas-Krankenhaus, Innere Medizin 2,
Caritaskrankenhaus Bad Mergentheim,
Uhlandstr. 7, 97980 Bad Mergentheim, Germany
e-mail: christoph.dietrich@ckbm.de

B. Braden
Medical Department I, University Hospital Frankfurt,
Theodor Stern Kai 7, 60590 Frankfurt am Main, Germany

B. Braden
Medical Department, John Radcliffe Hospital,
Headley Way, Oxford OX3 9DU, UK

M. Ott
Department of Pathology, Caritas-Krankenhaus,
Uhlandstr. 7, 97980 Bad Mergentheim, Germany

M. Hocke
Department of Internal Medicine II,
Friedrich Schiller University Jena,
Erlanger Allee 101, 07740 Jena, Germany

Table 1 Included and excluded patients according to the final diagnosis

Final diagnosis	CEUS performed	Excluded from study evaluation	Included in study evaluation
Pancreatic cancer	154	84	70
Neuroendocrine tumour	82	57	25
Serous microcystic adenoma	16	2	14
Lymphoma	8	8	0
Acinus cell carcinoma	3	3	0
Mucinous cystadenoma or cystadenocarcinoma	16	16	0
Intraductal papillary mucinous neoplasia	5	5	0
Autoimmune pancreatitis	3	0	3

Exclusion criteria were: no sign of chronic pancreatitis, cystic lesion > 20 mm, no active hormone production of the tumour, multiple lesions, metastasis, allergic reaction to SonoVue