

Robert Y. L. Zee · Nancy R. Cook · Chung-Ah Kim  
Arturo Fernandez-Cruz · Klaus Lindpaintner

## ***TP53* haplotype-based analysis and incidence of post-angioplasty restenosis**

Published online: 13 October 2004  
© Springer-Verlag 2004

---

### **Hum Genet (2004) 114:386–390**

In the Abstract section, the haplotypes (which appear twice in the abstract) should be 16bp–P72-G13494, and 16bp+ -P72-A13494 (not R72).

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00439-003-1080-8>

---

R. Y. L. Zee (✉) · N. R. Cook  
Division of Preventive Medicine,  
Brigham and Women's Hospital,  
Harvard Medical School,  
900 Commonwealth Avenue East,  
Boston, MA 02215, USA  
E-mail: rzee@rics.bwh.harvard.edu  
Tel.: +1-617-7328175  
Fax: +1-617-7839212

C.-A. Kim  
Division of Endocrine-Hypertension,  
Brigham and Women's Hospital,  
Harvard Medical School, Boston,  
Massachusetts, USA

A. Fernandez-Cruz  
Hospital Universitario San Carlos,  
Ciudad Universitaria, Madrid, Spain

K. Lindpaintner  
F. Hoffmann-La Roche Ltd.,  
Basel, Switzerland