

Itai Benjamini · Elchanan Mossel

On the mixing time of a simple random walk on the super critical percolation cluster

Received: 16 December 2001 / Revised version: 13 August 2002 /
Published online: 19 December 2002 – © Springer-Verlag 2002

Abstract. We study the robustness under perturbations of mixing times, by studying mixing times of random walks in percolation clusters inside boxes in \mathbf{Z}^d . We show that for $d \geq 2$ and $p > p_c(\mathbf{Z}^d)$, the mixing time of simple random walk on the largest cluster inside $\{-n, \dots, n\}^d$ is $\Theta(n^2)$ – thus the mixing time is robust up to a constant factor. The mixing time bound utilizes the Lovász-Kannan average conductance method. This is the first non-trivial application of this method which yields a tight result.

1. Introduction

An important parameter of random walks on finite graph is the mixing time of the random walk. We refer the reader to [1] for background and many references, or to subsection 1.2 for terminology used in this paper.

It is natural to study the robustness of the mixing time under perturbations. In this paper we address this issue by studying the effect of *random* perturbations of the underlying graph on the mixing times of simple random walk inside boxes in \mathbf{Z}^d .

A classical way to perturb the lattice \mathbf{Z}^d is by performing super-critical percolation. See [7] for introduction and many references, or subsection 1.2 for terminology.

In this paper we study the mixing time of simple random walk on the largest super-critical percolation cluster inside $\{-n, \dots, n\}^d$. We show that for all $d \geq 2$, the mixing time of the random walk on the perturbed box, is up to constant, the same as the mixing time of simple random walk on the original box.

Below we use the notation $g(n) = O(f(n))$ to indicate that $\limsup g(n)/f(n) < \infty$. We will write $f(n) = \Omega(g(n))$ when $g(n) = O(f(n))$, and $f(n) = \Theta(g(n))$ when both $f(n) = O(g(n))$ and $g(n) = O(f(n))$.

1.1. Main results

Let $G = (V, E)$ be a finite graph with no loops. Consider a simple random walk on G . In order to avoid periodicity of the random walk, we work with the contin-

uous time random walk. Look at the markov semi-group generated by the matrix $(Q_{x,y})_{x \in V, y \in V}$ where

$$Q_{x,y} = \begin{cases} \frac{1}{\deg(x)} & \text{if } x \sim y, \\ -1 & \text{if } x = y, \\ 0 & \text{otherwise.} \end{cases} \tag{1}$$

We denote by $\tau_1 = \tau_1(G)$ the mixing time of the random walk in total variation, by $\varphi = \varphi(G)$ the Cheeger constant of the graph G (see subsection 1.2 or [1]). We will denote by $B_d(n)$ the graph $G = (V, E)$ where

$$V = \{-n, \dots, n\}^d, \\ E = \{((x_1, \dots, x_d), (y_1, \dots, y_d)) \in V \times V : \sum_{i=1}^d |x_i - y_i| = 1\}.$$

It is well known that $\tau_1(B_d(n)) = \Theta(n^2)$ and $\varphi(B_d(n)) = \Theta(n^{-1})$.

Let $p_c(\mathbf{Z}^d)$ be the critical parameter for bond percolation in \mathbf{Z}^d (see [7] or subsection 1.2). Fix $p > p_c(\mathbf{Z}^d)$, and denote by $\mathcal{C} = \mathcal{C}_d(n) = \mathcal{C}_d(n, p)$ the largest cluster inside $B_d(n)$. Thus $\mathcal{C}_d(n)$ is the open component of $B_d(n)$ which has the maximal number of edges.

The following result proves the stability of the mixing time and the Cheeger constant under percolation:

Theorem 1.1. *If $d \geq 2$ and $p > p_c(\mathbf{Z}^d)$, then there exist constants $0 < c = c(p) < C = C(p)$ such that*

$$\lim_{n \rightarrow \infty} \mathbf{P}_p[cn^2 < \tau_1(\mathcal{C}_d(n)) < Cn^2] = 1. \tag{2}$$

Similarly, if $d \geq 2$ and $p > p_c(\mathbf{Z}^d)$, then there exist constants $0 < c = c(p) < C = C(p)$ such that

$$\lim_{n \rightarrow \infty} \mathbf{P}_p\left[\frac{c}{n} < \varphi(\mathcal{C}_d(n)) < \frac{C}{n}\right] = 1. \tag{3}$$

The upper bound on the mixing time is achieved via an estimate of a weighted variant of the Cheeger constant which was introduced by Lovász and Kannan [9]. In order to obtain tight bounds by the Lovász-Kannan method, we study geometrical properties of the percolation cluster, using in particular bootstrap and renormalization. One of the novel features of our result is that it is the first non-trivial application of the Lovász-Kannan method which yields a tight result. We conclude this section by recalling some background from percolation theory and from the theory of finite markov chains. Section 2 contains the proofs of these theorems, and Section 3 contains some remarks, conjectures and open problems.

We find it useful to use the following terminology (analogously to that used in the theory of random graphs). Let \mathcal{G}_n be an event describing a property of $\mathcal{C}_d(n)$. We will say that the event holds asymptotically almost surely (abbreviated a.a.s.), if $\lim_{n \rightarrow \infty} \mathbf{P}_p[\mathcal{G}_n] = 1$.

1.2. Background

Bernoulli bond percolation. In Bernoulli bond percolation on \mathbf{Z}^d , the edges of \mathbf{Z}^d are open (respectively closed) with probability p (respectively $1 - p$) independently. The corresponding product measure on the configurations of edges is denoted by \mathbf{P}_p or just \mathbf{P} . Let $\mathcal{C}(v)$ be the (open) cluster of v . In other words, $\mathcal{C}(v)$ is the maximal connected component of open edges in \mathbf{Z}^d containing v .

We write

$$\theta^v(p) = \mathbf{P}_p\{\mathcal{C}(v) \text{ is infinite}\}.$$

Since \mathbf{Z}^d is transitive, we may write $\theta(p)$ for $\theta^v(p)$. If $\mathcal{C}(v)$ is infinite for some v , we say that *percolation* occurs. We refer the reader to [7] for more background.

A particular property of super-critical percolation that we use below is the following

Proposition 1.2. *If $p > p_c(\mathbf{Z}^d)$ then there exists $p' = p'(p, d) > 0$ such that a.a.s. the number of open edges in $\mathcal{C}_d(n)$ is at least $p'd(2n + 1)^d$. Moreover, there exists a constant $c > 0$ s.t. a.a.s. $\mathcal{C}_d(n) \cap B_d(n - c \log n)$ is the intersection of the infinite percolation cluster with $B_d(n - c \log n)$.*

Proof. Let $\theta_e(p)$ be the probability that an edge e belongs to the infinite cluster (clearly the definition does not depend on the specific edge e). It is clear that since $\theta(p) > 0$ so does $\theta_e(p) > 0$. Consider $B_d(n)$ as a subgraph of \mathbf{Z}^d . By ergodicity it follows that a.a.s. there are at least $0.5\theta_e(p)d(2n + 1)^d$ edges in $B_d(n)$ which belong to the infinite percolation cluster. By [2] it follows that there exists some constant $c > 0$ such that a.a.s. if two edges $e_1, e_2 \in B_d(n - c \log n)$ belong to the infinite percolation cluster, then they are connected inside $B_d(n)$.

It remains to be shown that in $B_d(n)$ there is at most one connected component of size larger than $0.5\theta_e(p)d(2n + 1)^d$. This follows from the fact (see e.g. [7]) that when $p > p_c$ the probability that a vertex v which is not in the infinite cluster belongs to a connected cluster of size larger than k is bounded by $\exp(-\alpha k^{(d-1)/d})$ for some $\alpha > 0$. \square

Mixing and relaxation times; Cheeger constant. We follow [1] for basic notations and definitions. Consider the random walk on the graph $G = (V, E)$ with transition kernel (1) as a reversible markov chain. Note that π , the stationary distribution for the chain, satisfies

$$\pi(x) = \frac{\deg(x)}{\sum_{y \in C_n^d} \deg(y)}. \tag{4}$$

where $\deg(y)$ is the degree of y in G . Similarly, the probability of an edge e , denoted by $Q(e)$, is uniform for all edges of G . For two sets A and B write $Q(A, B) = \sum_{e=(x,y), x \in A, y \in B} Q(e)$. Let the eigenvalues of Q (1) be $0 = \lambda_1 > \lambda_2 \geq \dots$. We let the *spectral gap* of the random walk on G be $-\lambda_2$, and the *relaxation time* of the random walk be $\tau_2 = -\lambda_2^{-1}$.

For two distribution measures μ and ν on the same discrete space. The *total-variation distance*, $d_V(\mu, \nu)$, between μ and ν is defined as

$$d_V(\mu, \nu) = \frac{1}{2} \sum_x |\mu(x) - \nu(x)| = \sup_A |\mu(A) - \nu(A)|.$$

Consider again the random walk on G . Denote by π_x^t the distribution measure of the walk started at x at time t . The *mixing time* of the random walk, τ_1 , is defined as

$$\tau_1 = \inf\{t : \sup_{x,y} d_V(\pi_x^t, \pi_y^t) \leq e^{-1}\}.$$

Usually it is harder to estimate τ_1 than it is to estimate τ_2 . However, in general, the following relation holds:

$$\tau_2 \leq \tau_1 \leq \tau_2 \left(1 + \frac{1}{2} \log \frac{1}{\min_x \pi(x)}\right) \tag{5}$$

(see e.g. [1] Lemma 23).

A geometric tool which is used in order to bound relaxation times is Cheeger inequality. For a set A we define its conductance as

$$\varphi_A = \frac{Q(A, A^c)}{\pi(A)\pi(A^c)}. \tag{6}$$

Let φ be the Cheeger constant:

$$\varphi = \inf_A \varphi_A = \inf_{A:0 < \pi(A) \leq 1/2} \varphi_A. \tag{7}$$

Cheeger inequality states that

$$\tau_2 \leq 8\varphi^{-2}. \tag{8}$$

(see e.g. [1] Theorem 40).

In [9], Lovász and Kannan introduced the following variant of the Cheeger constant. For $0 < x \leq 1/2$, let

$$\varphi(x) = \inf_{A:0 < \pi(A) \leq x} \varphi_A. \tag{9}$$

Then it is shown in [9] (see [10] where the formula in [9] is corrected to include the second term) that:

$$\tau_1 \leq 32 \int_{\min_i \pi_i}^{1/2} \frac{1}{x\varphi^2(x)} + \frac{8}{\varphi}. \tag{10}$$

2. Mixing times

2.1. Lower bounds

We start by proving the upper bound on the Cheeger constant and the lower bound on the mixing time.

Lemma 2.1. *For all $d \geq 2$ and $p > p_c(\mathbf{Z}^d)$ there exists a constant $c = c(p, d) > 0$ such that a.a.s. it holds for $\mathcal{C}_d(n)$ that if x satisfies $\min_v \pi(v) \leq x \leq 1/2$, then*

$$\varphi(x) < \frac{c}{nx^{1/d}}. \tag{11}$$

Proof. By Proposition 1.2. there exists $p'(p)$ such that a.a.s. there are at least $p'd(2n+1)^d$ edges belonging to $\mathcal{C}_d(n)$. We will prove that (11) holds for $\min_v \pi(v) \leq x \leq q$, for some constant q . By the monotonicity of φ in x , it then follows that by increasing the value of c , (11) holds for all $\min_v \pi(v) \leq x \leq 1/2$. Let r be such for all $k < n/r + 1$ it is possible to cover at least a $(1 - p'/2)$ fraction of the edges of $B_d(n)$ by disjoint translations of $B_d(k)$. Let $k < n/r + 1$. It is clear that at least one of the translations $v + B_d(k)$ satisfies

$$\frac{2}{p'} \left(\frac{2k+1}{2n+1} \right)^d \geq \pi(B_d(k) + v) \geq \frac{p'}{2} \left(\frac{2k+1}{2n+1} \right)^d.$$

(the inequality on the left is satisfied by all translations).

On the other hand, since in \mathbf{Z}^d there are $2d(2k+1)^{d-1}$ edges going out of $v + B_d(k)$,

$$Q(v + B_d(k), \mathcal{C}_d(n) \setminus v + B_d(k)) \leq \frac{2d(2k+1)^{d-1}}{p'd(2n+1)^d} = \frac{2(2k+1)^{d-1}}{p'(2n+1)^d}.$$

It therefore follows that

$$\varphi_{v+B_d(k)} \leq \frac{8}{p'^2(2k+1)} \leq \frac{c}{n\pi^{1/d}(v + B_d(k))}$$

for some constants c . Choosing k to be the maximal such that

$$x \geq \frac{2}{p'} \left(\frac{2k+1}{2n+1} \right)^d$$

we obtain the required result (taking $q = p'r^{-d}/4$). \square

In order to prove the lower bound on the mixing time, we will use the following lemma. Let π be the stationary distribution for the simple random walk on the graph G (see (4)). Let $f : V \rightarrow \mathbb{R}$ be a function. We write $\pi[f] = \sum_{v \in V} \pi(v) f(v)$ for the expected value of f with respect to π .

Lemma 2.2. *Let $G = (V, E)$ be a finite graph. For each $v \in V$, let $D_v : V \rightarrow \mathbb{R}$ be defined by $D_v(x) = D(v, x)$ where D is the graph metric distance. Then*

$$\tau_2 \geq \max_v \left(\pi[D_v^2] - \pi^2[D_v] \right).$$

Proof. Take D_v as a test function in the extremal characterization of τ_2 (see e.g. [1]):

$$\tau_2 = \sup \left\{ \frac{\pi[g^2] - \pi^2[g]}{\mathcal{E}(g, g)} : \mathcal{E}(g, g) \neq 0 \right\},$$

where \mathcal{E} is the Dirichlet form:

$$\mathcal{E}(g, g) = \frac{1}{2} \sum_u \sum_{w \neq u} \pi[u] Q_{u,w} (g(u) - g(w))^2,$$

and note that if $Q_{u,w} \neq 0$, then $D_v(u) - D_v(w) \in \{-1, 0, 1\}$. \square

Lemma 2.3. *For all $d \geq 2$, there exists a constant $c > 0$ such that $\lim_{n \rightarrow \infty} \mathbf{P}_p[\tau_1 \geq cn^2] = 1$.*

Proof. Without loss of generality assume that 0 belong to the largest open cluster inside $B_d(n)$. From [2] it follows that there exist $a > 0, b > 0$ such that a.s. if $x, y \in B_d(n - b \log n) \cap \mathcal{C}_d(n)$, then $D(x, y) \leq a|x - y|_1$.

From proposition 1.2 it follows that a.s. there are at least $c'n^d$ vertices x with $|x|_1 \leq n/4a$ and at least $c'n^d$ vertices with $(1 - 1/(8a))n \geq |x|_1 \geq (1 - 1/(4a))n$ for some constant $c > 0$. Therefore, there are at least $c'n^d$ vertices x with $D_0(x) \leq n/4$ and at least $c'n^d$ vertices with $D_0(x) \geq 3n/4$.

Applying lemma 2.2 with the function D_0 it follows that a.s. $\tau_2 \geq cn^2$ for some constant $c > 0$. Using the lower bound (5) on τ_1 in terms of τ_2 we achieve the desired conclusion. \square

2.2. Bootstrap

In this subsection we show how Theorem 2.4 below implies the upper bound in Theorem 1.1. The proof of Theorem 2.4 is given in the following subsections.

Theorem 2.4. *For all $d \geq 2$, and all $p > p_c(\mathbf{Z}^d)$, there exist constants $c_1 = c_1(d, p) > 0$ and $c_2 = c_2(d, p) > 0$ such that $\mathcal{C}_d(n)$ satisfies a.s. that for sets A such that A and A^c are connected and $\frac{1}{2} \geq \pi(A) \geq \frac{c_1 \log^{\frac{d-1}{d}} n}{n^d}$,*

$$\varphi_A \geq \frac{c_2}{n\pi^{1/d}(A)}. \tag{12}$$

We will also utilize the following lemma which has the same proof as Lemma 36 of [1].

Lemma 2.5. *For all x , the minimum in (9) is obtained for a set A such that A is connected. The minimum at (7) is obtained for a set A such that A and A^c are connected.*

For x , there exists a set A for which the value $\varphi(x)$ is obtained and such that A is connected. However, it may be the case that the complement of the set is not connected. We bootstrap in the lemma below in order to prove that the estimates in Theorem 2.4 suffice for our purposes.

Lemma 2.6. *For all $d \geq 2$, and all $p > p_c(\mathbf{Z}^d)$, there exist constants $c_1 = c_1(d, p) > 0$ and $c_2 = c_2(d, p) > 0$ such that $\mathcal{C}_d(n)$ satisfies a.a.s. that for all $\frac{1}{2} \geq x \geq \frac{c_1 \log^{\frac{d-1}{d}} n}{n^d}$,*

$$\varphi(x) \geq \frac{c_2}{nx^{1/d}}. \tag{13}$$

Proof. By Lemma 2.5 the claim holds for $x = 1/2$. Thus by the monotonicity of φ in x it follows that for any constant $q > 0$ by increasing the value of c_2 , we obtain that a.a.s. (13) holds for all $q \leq x \leq 1/2$. We will therefore prove that (13) holds a.a.s. for all $q \geq x \geq \frac{c_1 \log^{\frac{d-1}{d}} n}{n^d}$, where q is determined below. Assume that A is a set for which $\varphi(x) = \varphi_A$. By Lemma 2.5 we may assume that the set A is connected. If $\mathcal{C}_d(n) \setminus A$ is connected, we are done, so we assume the contrary. Note that by Lemma 2.1 a.a.s. $\varphi_A \leq \frac{c'}{nx^{1/d}}$ for some constant c' . In particular $Q(A, \mathcal{C}_d(n) \setminus A) \leq \frac{c'x^{1-1/d}}{n}$. By the assumption that $\mathcal{C}_d(n) \setminus A$ is not connected we may write $\mathcal{C}_d(n) \setminus A$ as the union of disconnected components A_1, \dots, A_r , where $r \geq 2$. Let i be the index for which $\pi(A_i)$ is maximized. Without loss of generality we may assume that $\pi(A_i) \leq 1/2$ (otherwise repeat the argument below for $\mathcal{C}_d(n) \setminus A$).

Claim 2.7.

$$\pi(A_i) \geq \pi(A).$$

Proof. Assume the contrary and that $q \leq 1/4$. Note that by Lemma 2.5 a set for which the value of φ is obtained is connected and has a connected complement. In particular, for some constant c'' , a.a.s. $\varphi \geq c''/n$. By the assumptions that $\pi(A) \leq 1/4$, and that $\pi(A_i) < \pi(A)$ for all i , we may find a sub-collection $I \subset \{1, \dots, r\}$ such that if $B = A \cup_I A_i$, then $3/4 > \pi(B) > 1/4$. Note that $Q(B, \mathcal{C}_d(n) \setminus B) \leq Q(A, \mathcal{C}_d(n) \setminus A)$. This implies that

$$\frac{c''}{n} \leq \varphi \leq \varphi_B \leq 8Q(A, \mathcal{C}_d(n) \setminus A) \leq \frac{8c'}{n}x^{1-1/d} \leq \frac{8c'}{n}q^{1-1/d}.$$

Thus, when q is sufficiently small, we obtain a contradiction and the proof follows. \square

Proof of Lemma 2.6 continued. Recall that i is the index for which $\pi(A_i)$ is maximized and let A' be the (π) smallest set among A_i and $\mathcal{C}_d(n) \setminus A_i$, so that $\pi(A') \leq 1/2$. Note that A' is connected, and has a connected complement. Moreover by claim 2.7 $1 - \pi(A) \geq \pi(A') \geq \pi(A)$. It follows by Theorem 2.4 that $\varphi_{A'} \geq \frac{c_2}{n\pi^{1/d}(A')}$. However, this implies that

$$\begin{aligned} Q(A, \mathcal{C}_d(n) \setminus A) &\geq Q(A', \mathcal{C}_d(n) \setminus A') \geq c_2 \frac{\pi^{1-1/d}(A')(1 - \pi(A'))}{n} \\ &\geq c_2 \frac{\pi^{1-1/d}(A)(1 - \pi(A))}{n} \end{aligned}$$

and we obtain that

$$\varphi_A \geq \frac{c_2}{nx^{1/d}},$$

as needed. \square

Proof of the upper bound in theorem 1.1. We assume that (13) holds. If the set A satisfies $\pi(A) \leq \frac{c_1 \log^{\frac{d}{d-1}} n}{n^d}$, then since $C_d(n)$ is connected it follows that $Q(A, A^c) \geq 1/(2dn^d)$ and therefore

$$\varphi_A \geq \frac{Q(A, A^c)}{\pi(A)} \geq \frac{1}{2dc_1 \log^{\frac{d}{d-1}} n}. \tag{14}$$

Since by Lemma 2.5 the set A which achieves the minimum at the definition of the Cheeger constant (7) is connected, we obtain by (14) and (12) that

$$\lim_{n \rightarrow \infty} \mathbf{P}_p[\varphi \geq \frac{2^{1/d} c_2}{n}] = 1.$$

Moreover, by (14) and (13) we obtain that a.a.s. for all $\frac{1}{2} \geq x \geq \frac{c_1 \log^{\frac{d}{d-1}} n}{n^d}$,

$$\varphi(x) \geq \min\left\{\frac{1}{2dc_1 \log^{\frac{d}{d-1}} n}, \frac{c_2}{nx^{1/d}}\right\}.$$

Thus, by (10), we see that a.a.s.

$$\int_{\min_v \pi(v)}^{\frac{1}{2}} \frac{1}{t\varphi^2(t)} dt \leq \int_{\frac{1}{2dn^d}}^{\frac{1}{2}} \frac{4d^2 c_1^2 \log^{\frac{2d}{d-1}} n}{t} dt + \int_0^{\frac{1}{2}} \frac{n^2}{c_2^2 t^{1-2/d}} dt \leq C'n^2,$$

for $C' = C'(d, p)$, and therefore a.a.s.

$$\tau_1 \leq 32 \int_{\min_v \pi(v)}^{\frac{1}{2}} \frac{1}{t\varphi^2(t)} dt + \frac{8}{\varphi} \leq 32C'n^2 + \frac{8n}{2^{1/d} c_2} \leq Cn^2,$$

for some constant $C = C(d, p)$ as needed. \square

2.3. Upper bound for $d = 2$

In this subsection we present a short proof of Theorem 2.4 in the planar ($d = 2$) case. An independent proof for general $d \geq 2$ is given in Subsections 2.4 and 2.5 below.

The main tool in the proof will be the following large deviation result by Kesten [8]:

Lemma 2.8. *Consider i.i.d. percolation $\{X(e)\}$ on the edges of \mathbf{Z}^2 where*

$$\mathbf{P}_p[X(e) = 0] = 1 - \mathbf{P}_p[X(e) = 1] = p < 1/2 (= p_c(\mathbf{Z}^2)).$$

For two point x and y , let $D(x, y)$ be their distance in the first passage percolation model:

$$D(x, y) = \min\left\{\sum_{i=1}^k X(e_i) : e_1, \dots, e_k \text{ is a path connecting } x \text{ to } y\right\}.$$

Then, there exist constants $a > 0$ and $b > 0$ such that for all points x and y ,

$$\mathbf{P}_p[D(x, y) \leq a|x - y|_1] \leq \exp(-b|x - y|_1),$$

where $|(x_1, y_1) - (x_2, y_2)|_1 = |x_1 - x_2| + |y_1 - y_2|$.

Proof of theorem 2.4, $d = 2$. We will use the following dual first-passage percolation model. Take \mathbf{Z}^2 and draw the dual lattice \mathbf{Z}_*^2 . Each edge e of \mathbf{Z}^2 crosses a unique edge e^* of \mathbf{Z}_*^2 . If the edge e is closed, set $X(e^*) = 0$; otherwise, set $X(e^*) = 1$. Note that $\mathbf{P}_p[X(e^*) = 0] = 1 - p < 1/2$ and we may therefore apply Lemma 2.8 to $\{X(e^*)\}$. In particular, we obtain that if c' is large, then a.a.s. for all pairs of points x^* and y^* in the dual of $\mathcal{C}_2(n)$ such that $|x^* - y^*|_1 \geq c' \log n$, we have,

$$D(x^*, y^*) \geq a|x^* - y^*|_1. \tag{15}$$

Let A be a connected set in $\mathcal{C}_2(n)$ such that $1/2 \geq t = \pi(A) \geq \frac{c_1 \log^2 n}{n^2}$ and such that $\mathcal{C}_2(n) \setminus A$ is connected. $B_2(n) \setminus A$ is the union of (\mathbf{Z}^2) disconnected components A_1, \dots, A_r where $\mathcal{C}_2(n) \setminus A \subset A_r$. Let $A' = A \cup \cup_{i=1}^{r-1} A_i$. Then A' and $B_2(n) \setminus A'$ are both connected. Moreover by Proposition 1.2 a.a.s. both A' and $B_2(n) \setminus A'$ contain at least qtn^2 vertices for some constant q .

Let γ be the boundary of A' in $B_2(n)$. In other words, γ consists of all the edges (x, y) such that $x \in A', y \notin A'$. We let γ^* be the path which is obtained by taking the dual edges of the edges of γ . Since both A' and $B_2(n) \setminus A'$ are connected, γ^* is connected.

A set of l_1 diameter g in \mathbf{Z}^2 contains at most $(2g + 1)^2$ points. Therefore A' has l_1 diameter at least $c_1 \sqrt{qtn}$ for some constant $c_1 > 0$. In other words, there exist two point x^* and y^* on γ^* such that $|x^* - y^*|_1 \geq c_1 \sqrt{qtn}$. Let $|\gamma^*|$ be the number of edges e on γ^* for which $X(e) = 1$. Taking c_1 sufficiently large it follows by (15) that a.a.s. $|\gamma^*| \geq ac_1 \sqrt{qtn}$. Since $10n^2 Q(A, \mathcal{C}_2(n) \setminus A) \geq |\gamma^*|$, we obtain that

$$\varphi_A \geq \frac{ac_1 \sqrt{qt}}{10nt(1-t)} \geq \frac{c_2}{\sqrt{tn}},$$

for some constant $c_2 = c_2(d, p) > 0$ as needed. \square

2.4. Upper bound for $d \geq 2$ and p close to 1

We now prove the theorem for $d \geq 2$ assuming that p is close to 1. For two vertices v and w in $B_d(n)$, a *cutset* separating v from w is a set B of edges of $B_d(n)$ such that any path in $B_d(n)$ which connects v to w intersects B at least in one edge. A *minimal* cutset separating v from w , is a cutset which has no proper subset which is also a cutset separating v from w .

The following fact is probably well known. We refer the reader to Babson and Benjamini [3] for a proof (in a more general setting).

Lemma 2.9. *For $v, w \in B_d(n)$ the number of minimal cutsets of size m separating v from w is bounded by $c(d)^m$ for some constant $c(d)$.*

Proof of Theorem 2.4, $d \geq 2$ and p close to 1. By Proposition 1.2 a.a.s. the π measure of a subset $A \subset \mathcal{C}_d(n)$ is up to constant the same as the number of vertices in the set divided by $(2n + 1)^d$. Therefore, in order to show that (12) holds, it suffices to show that there exist constants $\hat{c} > 0$ and $\hat{c}_1 > 0$ such that a.a.s. all connected sets $A \subset \mathcal{C}_d(n)$ such that $\mathcal{C}_d(n) \setminus A$ is also connected and such that the size of $|A|$ is at least $\hat{c}_1 \log^{\frac{d}{d-1}} n$ and at most $|\mathcal{C}_d(n)|/2$, satisfy that the number of open edges going from A to $\mathcal{C}_d(n) \setminus A$ is at least $\hat{c}|A|^{\frac{d-1}{d}}$.

For such a set A , the set $B_d(n) \setminus A$ is a union of disconnected components A_1, \dots, A_r where $\mathcal{C}_d(n) \setminus A \subset A_r$. Fix a point $v \in A$ and $w \in A_r$ and look at the set B of edges connecting A to A_r (= the set of edges connecting $A \cup \cup_{i=1}^{r-1} A_i$ to A_r). This is a minimal cutset of edges separating v from w . Moreover, since $|A_r| \geq |A|$, it follows by the isoperimetric inequality [4] that $|B| \geq c_I |A|^{\frac{d-1}{d}} \geq c_I \hat{c}_1^{\frac{d-1}{d}} \log n$ for some constant c_I .

It follows that in order to prove the theorem, it suffices to show that there exist constants $c_1 > 0, c_2 > 0$ such that for all minimal cut-sets $|B|$ of size at least $c_1 \log n$, the number of open edges in $|B|$ is at least $c_2|B|$. We denote by \tilde{B} the subset of open edges of set B .

We will use a first moment argument. Applying large deviation estimates, we see that if $c_2 > 0$ is sufficiently small, and $p < 1$ is sufficiently large, then $\mathbf{P}_p[|\tilde{B}| \leq c_2|B|] \leq (c(d) + 1)^{-|\tilde{B}|}$.

Summing up, and using Lemma 2.9 we see that the probability that there exists any cut-set B with $|\tilde{B}| \leq c_2|B|$ and $|B| \geq c_1 \log n$ is bounded by

$$(2n + 1)^{2d} \sum_{s \geq c_1 \log n} c(d)^s (c(d) + 1)^{-s} = o(1),$$

provided that c_1 is sufficiently large. \square

2.5. Upper bound for $d \geq 2$ and $p > p_c(\mathbf{Z}^d)$

The proof uses renormalization and the result for p close to 1. The renormalization will produce site percolation with high density of “good” sites. We will need the following fact

Proposition 2.10. *There exists constants $p^* < 1, a > 0$ and $c > 0$ such that a.a.s. for site percolation with parameter $p > p^*$ all connected sets A in $B_d(n)$ of size at least $c \log n$ have at least $a|A|$ open sites.*

Proof. It is well known that the number of connected sets of size m containing a specified vertex v is bounded by $c(d)^m$ for some constant $c(d)$. If $p^* < 1$ is sufficiently large, and $a > 0$ is sufficiently small, then for each set A , the number of open sites in A denoted $|\tilde{A}|$ satisfies

$$\mathbf{P}_p[|\tilde{A}| \leq a|A|] \leq (c(d) + 1)^{-|A|}.$$

Summing over all sets, we see that the probability that there exists any connected set of size greater than $c \log n$ for which $|\tilde{A}| < a|A|$ is bounded by

$$(2n + 1)^d \sum_{s \geq c \log n} c(d)^s (c(d) + 1)^{-s} = o(1),$$

provided that c is sufficiently large. \square

We have proved Theorem 2.4 and therefore Lemma 2.6 for large $p < 1$. An analogous proof implies the analogous result for site percolation for large $p < 1$. For convenience we state this result below:

Lemma 2.11. *For all $d \geq 2$, there exists $p^* < 1$ such that for $p > p^*$, there exist constants $c_1 = c_1(d, p) > 0$ and $c_2 = c_2(d, p) > 0$ such that a.s. site percolation with parameter p on $B_d(n)$ satisfies for all $\frac{1}{2} \geq x \geq \frac{c_1 \log^{\frac{d-1}{d}} n}{n^d}$ that*

$$\varphi(x) \geq \frac{c_2}{nx^{1/d}}. \tag{16}$$

Proof of Theorem 2.4 $p > pc(\mathbf{Z}^d)$

For $v \in \mathbf{Z}^d$, we let

$$Q_N(v) = v + B_d(N) = \{w : |w - v|_\infty \leq N\}.$$

We will slightly abuse the notation by writing $Q_N(v)$ for the induced subgraph on $Q_N(v)$. We call $v \in (N\mathbf{Z})^d$ a *good vertex* if the following conditions hold:

- There exists an open cluster which intersects all $d - 1$ dimensional faces of the box $Q_{5N/4}(v)$.
- All connected components of diameter more than $N/10$ inside the box $Q_{5N/4}(v)$ intersect the above cluster.

By standard renormalization results (see Proposition 2.1 in Antal and Pisztora [2]) it follows that for any $p > pc(\mathbf{Z}^d)$, the set of good vertices stochastically dominates site percolation with parameter $p^*(N)$ on $(N\mathbf{Z})^d$, with $\lim_{N \rightarrow \infty} p^*(N) = 1$.

We take a connected set A in $\mathcal{C}_d(n)$ such that both A and $\mathcal{C}_d(n) \setminus A$ are connected and such that $C \log^{\frac{d}{d-1}} n \leq |A| \leq |\mathcal{C}_d(n) \setminus A|$. We will show that a.s. the number of open edges between A and $\mathcal{C}_d(n)$ is at least $c|A|^{\frac{d-1}{d}}$, where c and C are positive constants to be determined later. We will thus obtain the required result.

We let $A' = \{v \in (N\mathbf{Z})^d : |Q_{5N/4}(v) \cap A| \geq N/10\}$. Since A is connected, A' is a connected set in $(N\mathbf{Z})^d$. Moreover, it is clear that $|A|/(2N)^d \leq |A'| \leq |A|$. Let A_g be the set of good sites in A' . By Proposition 2.10 when $p^*(N)$ is sufficiently large, a.s. $|A_g| \geq a|A'| \geq \frac{a}{(2N)^d} |A|$.

It now follows by Lemma 2.11 that if C is sufficiently large, then a.s. there are at least $c''|A_g|^{\frac{d-1}{d}} \geq c'|A|^{\frac{d-1}{d}}$ pairs of good neighbors u and w such that $u \in A_g$ and $w \notin A_g$, where $c' > 0$.

We note that each such pair defines an open edge going from $A \cap Q_N(u)$ to $(\mathcal{C}_d(n) \setminus A) \cap Q_N(w)$. Moreover each such edge is defined by at most $4d^2$ pairs (u, w) . It now follows that a.s. the edge boundary of A is of size at least $\frac{c'}{4d^2} |A|^{\frac{d-1}{d}}$ as needed. \square

3. Further remarks

3.1. Coupling

For simple random graph models it is easy to bound from above the mixing time by constructing explicit coupling. We give two examples below

1. Let $G(n, p)$ be the random graph model for fixed p . It is easy to see that every two vertices have at least $\frac{p^2 n}{2}$ joint neighbors with probability going to 1 as $n \rightarrow \infty$. Therefore, by coupling we see that $\sup_{x,y} D_V(\mathbf{P}_x^t, \mathbf{P}_y^t) < (1 - \frac{p^2}{2})^t$. So the mixing time is $\Theta(1)$.
2. Consider the following perturbation of $B_2(n)$. To each vertex v of the square attach a pipe of length X_v where X_v are independent random variables taking the values 0, 1. As noted to us by Amir Dembo, one can use the usual reflecting coupling on the square $B_2(n)$, in order to show that the mixing time is $\Theta(n^2)$. Indeed, let x and y be two vertices and consider random walks starting at x and y . Always delay one of the walks in order that the two walks make steps in $B_2(n)$ simultaneously. Whenever the two walks make steps in $B_2(n)$, use the usual reflecting coupling. This example can be generalized to any dimension and the assumption on X_v may be replaced by a weaker moment assumption. In a previous draft of this paper, we had a more complicated result in the same spirit.

We think it is an interesting challenge to try a variant of the last argument showing via a coupling argument that the mixing time on C_n^d is $O(n^2)$ for all d . One approach of implicit construction of such coupling is to use some kind of central limit theorem in order to bring the walks closer and closer. Unfortunately the present form of the CLT on super critical percolation cluster (De Masi et al. (1989) [6]) provides no estimates on the convergence rate and therefore no bound on the coupling time of $C_d(n)$.

3.2. Final comments

1. In this note $p > p_c(\mathbf{Z}^d)$, is fixed, what is the dependence on p of the mixing time? What is the mixing times on the critical cluster?
2. One can consider the mixing time of random walks on percolation clusters on other graphs. With Nick Wormald (in preparation), it is shown that the mixing time for simple random walk on $G(n, c/n)$, $c > 1$ is poly-logarithmic in n .

Acknowledgement. Thanks to Oded Schramm and Prasad Tetali for helpful discussions and to the anonymous referee for helpful suggestions.

References

- [1] Aldous, D., Fill, J.A.: Reversible Markov chains and random walks on graphs, *book in preparation*. 2000

- [2] Antal, P., Pisztor, A.: On the chemical distance in supercritical Bernoulli percolation, *Ann. Probab.* **24**, 1036–1048 (1996)
- [3] Babson, E., Benjamini, I.: Cut sets and normed cohomology with applications to percolation. *Proc. Amer. Math. Soc.* **127**, 589–597 (1999)
- [4] Bollobás, B., Leader, Imre.: Edge-isoperimetric inequalities in the grid. *Combinatorica* **11**, 299–314 (1991)
- [5] Liggett, T., Schonmann, R., Stacey, A.: Domination by product measures. *Ann. Probab.* **25** 1, 71–95 (1997)
- [6] De Masi, A., Ferrari, P., Goldstein, S., Wick, W.: An invariance principle for reversible Markov processes. Applications to random motions in random environments. *J. Statist. Phys.* **55**, 787–855 (1989)
- [7] Grimmett, G.: Percolation. Second edition. Springer-Verlag, Berlin, 1999. xiv+444
- [8] Kesten, H.: On the time constant and path length of first-passage percolation. *Adv. in Appl. Probab.* **12**, 848–863 (1980)
- [9] Lovász, L., Kannan, R.: Faster mixing via average conductance *Proc. 1995 ACM STOC*.
- [10] Montenegro, R., Son, J.B.: Edge Isoperimetry and Rapid Mixing on Matroids and Geometric Markov Chains, *Proc. 2001 ACM STOC*