

Nikolai Dokuchaev

Estimates for distances between first exit times via parabolic equations in unbounded cylinders

Received: 5 June 2003 / Revised version: 14 January 2004 /
Published online: 25 March 2004 – © Springer-Verlag 2004

Abstract. First exit times and their dependence on variations of parameters are studied for diffusion processes with non-stationary coefficients. Estimates of L_p -distances and some other distances between two exit times are obtained. These estimates are based on some new prior estimates for solutions of parabolic Kolmogorov's equations with infinite horizon without Cauchy conditions.

1. Introduction

It is known that first exit times from a region for smooth solutions of ordinary equations do not depend continuously on variations of the initial data or the coefficients. However, first exit times for non-smooth trajectories of diffusion processes have some path-wise regularity with respect to these variations (some related results can be found in author's papers (1987),(1992)). This paper studies path-wise dependence on variations of initial data and coefficients for first exit times of diffusion processes from a domain $D \subset \mathbf{R}^n$. We present an effective estimate of distances between exit times via estimates for solutions $v(x, t)$ of backward Kolmogorov's parabolic equations in the unbounded cylinder $D \times [0, +\infty)$, when the Cauchy boundary condition is replaced by the condition $\sup_{t \geq 0} \|v(\cdot, t)\|_{L_2(D)} < +\infty$. These problems are sometimes called Fourier problems. Similar problems were studied earlier by Bokalo (1993), Oleinik and Yosifian (1976), and others. Nakao (1984) and Shelukhin (1993) considered related problems for periodic solutions. Oleinik and Yosifian (1976) and Bokalo (1993) consider non-linear equations, but under some additional condition. For our special case of Kolmogorov's equations, this condition requires that the generating differential operator is self-adjoint (i.e., $\mathcal{A} = \mathcal{A}^*$ in our notations below). This condition is too restrictive for applications in stochastic analysis, and we do not require it in the present paper. We obtained solvability and uniqueness results as well as new prior estimates for norms in Sobolev spaces and Hölder spaces for solutions of Kolmogorov's equations with infinite horizon. Using these estimates, we found effective estimates of $\mathbf{E}|\tau_1 - \tau_2|^p$ and

N. Dokuchaev: Department of Mathematics and Statistics, University of Limerick, Ireland.
e-mail: Nikolai.Dokuchaev@ul.ie, or N.Dokuchaev@yahoo.com

Mathematics Subject Classifications (2000): 60G17, 60G40, 60J50, 60J60, 60J65

Key words or phrases: Diffusion processes – First exit times – Parabolic Kolmogorov's equations

$\mathbf{E}\{\exp(\lambda|\tau_1 - \tau_2|) - 1\}$ for exit times τ_1 and τ_2 of two diffusion processes, where $p \geq 1$ and $\lambda \in (0, \lambda_{\max})$, and where $\lambda_{\max} > 0$ depends on the class of coefficients of the Itô's equations.

Some definitions

Assume that we are given an open bounded domain $D \subset \mathbf{R}^n$ with C^2 -smooth boundary ∂D . We denote Euclidean norm as $|\cdot|$, and \bar{D} denotes the closure of a region D .

We denote by $\|\cdot\|_X$ the norm in a linear normed space X , and $(\cdot, \cdot)_X$ denotes the scalar product in a Hilbert space X .

Let us introduce some spaces of functions. Let $G \subset \mathbf{R}^k$ be an open domain, then $W_q^m(G)$ denotes the Sobolev space of functions that belong $L_q(G)$ together with first m derivatives, $q \geq 1$.

Let $H^0 \triangleq L_2(D)$, i.e., it is a Hilbert space, and let $H^1 \triangleq \bar{W}_2^1(D)$ be the closure in the $W_2^1(D)$ -norm of the set of all smooth functions that vanish in a neighborhood of ∂D , $k = 1, 2$. Let $H^2 = W_2^2(D) \cap H^1$ be the space equipped with the norm of $W_2^2(D)$. Let H^{-1} be the dual space to H^1 , with the norm $\|\cdot\|_{H^{-1}}$ such that if $u \in H^0$ then $\|u\|_{H^{-1}}$ is the supremum of $(u, v)_{H^0}$ over all $v \in H^0$ such that $\|v\|_{H^1} \leq 1$.

The scalar product $(u, v)_{H^0}$ is assumed to be well defined for $u \in H^{-1}$ and $v \in H^1$ as well, meaning the natural extending from $u \in H^0$ and $v \in H^1$.

Let ℓ_m be the Lebesgue measure in \mathbf{R}^m , and let $\bar{\mathcal{B}}_m$ be the σ -algebra of the Lebesgue sets in \mathbf{R}^m .

Let $0 \leq s < T \leq +\infty$ and let $\mathcal{Q} \triangleq D \times (s, T)$.

Let us introduce the following spaces:

$$C^k(s, T) \triangleq C([s, T]; H^k), \quad X_r^k(s, T) \triangleq L_r([s, T], \bar{\mathcal{B}}_1, \ell_1; H^k),$$

$$k = -1, 0, 1, 2.$$

Let $X_{r,loc}^k(s, T)$ be the set of all functions $u(x, t) : \mathcal{Q} \rightarrow \mathbf{R}$ such that $u|_{D \times [\theta, t]} \in X_r^k(\theta, t)$ for all θ, t such that $s < \theta < t < T$, $k = -1, 0, 1, 2$.

Let $Y^k(s, +\infty)$ be linear normed spaces of functions $u : [s, +\infty) \rightarrow H^k$ with finite norm

$$\|u\|_{Y^k(s, +\infty)} = \sup_{m=0,1,2,\dots} \left(\int_{s+m}^{s+m+1} \|u(\cdot, t)\|_{H^k}^2 dt \right)^{1/2} \quad k = -1, 0, 1, 2.$$

For $\gamma \geq 1$, let $W_\gamma^{2,1}(\mathcal{Q})$ be a Banach space of functions $u : \mathcal{Q} \rightarrow \mathbf{R}$ that belong to $L_\gamma(\mathcal{Q})$ together with all derivatives $\frac{\partial u}{\partial t}, \frac{\partial u}{\partial x_k}, \frac{\partial^2 u}{\partial x_k \partial x_m}, k, m = 1, \dots, n$, with finite norm

$$\|u\|_{W_\gamma^{2,1}(\mathcal{Q})} \triangleq \|u\|_{L_\gamma(\mathcal{Q})} + \left\| \frac{\partial u}{\partial t} \right\|_{L_\gamma(\mathcal{Q})} + \sum_{k=1}^n \left\| \frac{\partial u}{\partial x_k} \right\|_{L_\gamma(\mathcal{Q})} + \sum_{k,m=1}^n \left\| \frac{\partial^2 u}{\partial x_k \partial x_m} \right\|_{L_\gamma(\mathcal{Q})}.$$

Let $\alpha \in (0, 1)$ be a non-integer number. We will say that a function $u : Q \rightarrow \mathbf{R}$ belongs to the class $\mathcal{H}^{1+\alpha, (1+\alpha)/2}(Q)$ if u and $\partial u/\partial x$ are continuous, and

$$\langle\langle u \rangle\rangle_Q^{(1+\alpha)} \triangleq \langle u \rangle_Q^{(\alpha)} + \sum_{k=1}^n \left\langle \frac{\partial u}{\partial x_k} \right\rangle_Q^{(\alpha)} < +\infty,$$

where

$$\begin{aligned} \langle u \rangle_Q^{(\alpha)} \triangleq & \max_{(x,t) \in Q} |u(x,t)| + \sup_{x,x' \in D, t \in [s,T]} \frac{|u(x,t) - u(x',t)|}{|x - x'|^\alpha} \\ & + \sup_{x \in D, t,t' \in [s,T]} \frac{|u(x,t) - u(x,t')|}{|t - t'|^{\alpha/2}}. \end{aligned}$$

This class is a special case of the Hölder space from Ladyzhenskaya *et al.* (1968), p.7.

2. Main results

2.1. Parabolic equations in unbounded cylinders

Let $Q \triangleq D \times [0, +\infty)$.

Let $f(x, t) : Q \rightarrow \mathbf{R}^n$ and $\beta(x, t) : Q \rightarrow \mathbf{R}^{n \times n}$ be measurable functions.

Let $(\Omega, \mathcal{F}, \mathbf{P})$ be a standard probability space. Consider n -dimensional Itô's equation

$$\begin{cases} dy(t) = f(y(t), t)dt + \beta(y(t), t)dw(t), & t > s, \\ y(s) = a. \end{cases} \tag{2.1}$$

Here $w(t)$ is a standard n -dimensional Wiener process. The random vectors a with values in \bar{D} does not depend on $w(t) - w(r)$ for all $t > r > s$.

We denote by $y^{a,s}(t)$ the solution of (2.1) given a and s .

Let $\tau^{a,s} \triangleq \inf\{t \geq s : y^{a,s}(t) \notin D\}$.

Let $b \triangleq \beta\beta^\top$ and $Q_t \triangleq D \times (t, t + 1)$. We assume that

$$\begin{cases} \beta \in C(Q; \mathbf{R}^{n \times n}), \quad \sup_{(x,t) \in Q_s} \left| \frac{\partial f}{\partial x}(x, t) \right| < +\infty \quad \forall s > 0, \\ c_\beta \triangleq \sup_{(x,t) \in Q} |\beta(x, t)| < +\infty, \quad c_f \triangleq \sup_{(x,t) \in Q} |f(x, t)| < +\infty, \\ \bar{c}_\beta \triangleq \text{ess sup}_{(x,t) \in Q} \left| \frac{\partial \beta}{\partial x}(x, t) \right| < +\infty, \quad \delta \triangleq \inf_{(x,t) \in Q, \xi \in \mathbf{R}^n} \frac{\xi^\top b(x,t)\xi}{|\xi|^2} > 0. \end{cases} \tag{2.2}$$

Under these assumptions, equation (2.1) has the unique strong solution.

Set

$$\mathcal{P}_0 \triangleq (n, D, c_\beta, \bar{c}_\beta, c_f, \delta).$$

Let $\Theta(\mathcal{P}_0)$ be the set of all pairs (f, β) of functions $f : Q \rightarrow \mathbf{R}^n, \beta : Q \rightarrow \mathbf{R}^{n \times n}$ such that conditions (2.2) are satisfied.

Lemma 2.1. *There exists $\nu \in (0, 1)$ such that $\nu = \nu(n, D, c_f, \delta)$ depends only on (n, D, c_f, δ) and $\mathbf{P}(\tau^{a,s} > s + 1) \leq \nu$ for all $(f, \beta) \in \Theta(\mathcal{P}_0)$, for all $s \geq 0$, and for any random vector a such that $a \in D$ a.s. and a does not depend on $w(t) - w(r)$ for all $t > r > s$.*

Note that if the functions $f(x, t)$ and $\beta(x, t)$ are either constant in t or periodic in t with the period 1, then existence of $\nu = \nu(f, \beta) \in (0, 1)$ such that $\mathbf{P}(\tau^{a,s} > s + 1) \leq \nu (\forall a, s)$ is obvious.

Introduce the differential operator

$$\mathcal{A}(t)v(x) \triangleq \sum_{k=1}^n f_k(x, t) \frac{\partial v}{\partial x_k}(x) + \frac{1}{2} \sum_{k,m=1}^n b_{km}(x, t) \frac{\partial^2 v}{\partial x_k \partial x_m}(x). \tag{2.3}$$

Here x_k, f_k , and b_{km} are the components of the vectors x, f and the matrix $b = \beta\beta^\top$.

Let $q(x, t) : Q \rightarrow \mathbf{R}$ be a bounded measurable function.

Consider the boundary value problem in the semi-infinite cylinder $Q = D \times [0, +\infty)$

$$\begin{cases} \frac{\partial v}{\partial t}(x, t) + \mathcal{A}(t)v(x, t) + q(x, t)v(x, t) = -\varphi(x, t), \\ v(x, t)|_{x \in \partial D} = 0, \\ \text{ess sup}_{t \geq 0} \|v(\cdot, t)\|_{H^0} < +\infty. \end{cases} \tag{2.4}$$

Let

$$\mathcal{P}_q \triangleq \left(\mathcal{P}_0, \sup_{(x,t) \in Q} q(x, t) \right), \quad \mathcal{P}_{|q|} \triangleq \left(\mathcal{P}_0, \sup_{(x,t) \in Q} |q(x, t)| \right).$$

Theorem 2.1. *Let $\sup_{(x,t) \in Q} q(x, t) < -\ln \nu$ for all (x, t) , where $\nu = \nu(n, D, c_f, \delta)$ is such as is defined in Lemma 2.1. Let $(f, \beta) \in \Theta(\mathcal{P}_0)$. Then the following holds.*

(i) *For any $\varphi \in Y^{-1}(0, +\infty)$, there exists the unique (up to equivalency) solution $v : D \times (0, +\infty) \rightarrow \mathbf{R}$ of problem (2.4) in the class $X_\infty^0(0, +\infty) \cap X_{2,loc}^1(0, +\infty)$.*

(ii) *$v \in C^0(0, +\infty)$, and there exists a constant $c = c(\mathcal{P}_q)$ such that*

$$\sup_{t \in [0, +\infty)} \|v(\cdot, t)\|_{H^0} + \|v\|_{Y^1(0, +\infty)} \leq c \|\varphi\|_{Y^{-1}(0, +\infty)}. \tag{2.5}$$

(iii) *If $\varphi \in Y^0(0, +\infty)$ is a Borel measurable function, then*

$$v(x, s) = \mathbf{E} \int_s^{\tau^{x,s}} \varphi(y^{x,s}(t), t) \exp\left(\int_s^t q(y^{x,s}(r), r) dr\right) dt, \tag{2.6}$$

and this equality holds for all $s \geq 0$ for a.e. $x \in D$.

(iv) *If $\sup_{t \geq 0} \|\varphi|_{Q_t}\|_{L_\gamma(Q_t)} < +\infty$ for $\gamma \geq 2$, then $v|_{Q_t} \in W_\gamma^{2,1}(Q_t)$ for all $t \geq 0$, and there exists a constant $c = c(\mathcal{P}_{|q|}, \gamma)$ such that*

$$\sup_{t \geq 0} \|v|_{Q_t}\|_{W_\gamma^{2,1}(Q_t)} \leq c \sup_{t \geq 0} \|\varphi\|_{L_\gamma(Q_t)}. \tag{2.7}$$

(v) If $\sup_{t \geq 0} \|\varphi|_{Q_t}\|_{L_\gamma(Q_t)} < +\infty$ for $\gamma > n + 2$, then the function $v(x, t)$ and its derivatives $\partial v(x, t)/\partial x_k$ are continuous, bounded, and belong to the Hölder class $\mathcal{H}^{1+\alpha, (1+\alpha)/2}(Q_t)$ for all $t \geq 0$, where $\alpha \triangleq 1 - (n + 2)/\gamma$, $k = 1, \dots, n$. Moreover, there exists a constant $c = c(\mathcal{P}_{|q|}, \gamma)$ such that

$$\sup_{t \geq 0} \langle \langle u|_{Q_t} \rangle \rangle_{Q_t}^{(1+\alpha)} \leq c \sup_{t \geq 0} \|\varphi\|_{L_\gamma(Q_t)}. \tag{2.8}$$

Note that the parabolic equation in (2.4) is in the sense of Sobolev generalized functions, and we assume that the boundary condition $v(x, t)|_{x \in \partial D} = 0$ is satisfied if $v(\cdot, t) \in H^1 = W_2^1(D)$ for a.e. t . The definitions of the classes of functions ensure that the solution of problem (2.4) has sense.

Remark 2.1. The Krylov’s estimates give estimation of $\sup_{x \in D} |v(x, s)|$ for $q(x, t) \leq 0$ via the norm of φ in $L_{n+1}(D \times (s, +\infty))$ or via $\|\varphi\|_{L_n(D)}$ for independent on t functions $\varphi(x, t) = \varphi(x)$ (see Theorem II.4.2 from Krylov (1980)).

Consider now the boundary value problem with quasi-periodical conditions in the cylinder $Q_0 \triangleq D \times (0, 1)$

$$\begin{cases} \frac{\partial V}{\partial t}(x, t) + \mathcal{A}(t)V(x, t) + q(x, t)V(x, t) = -\varphi(x, t), \\ V(x, t)|_{x \in \partial D} = 0, \\ \mu V(x, 0) - V(x, 1) \equiv 0. \end{cases} \tag{2.9}$$

Theorem 2.2. Let $\mu \neq 0$. Let $(f, \beta) \in \Theta(\mathcal{P}_0)$. Let $\sup_{(x,t) \in Q} q(x, t) + \ln |\mu| < -\ln v$ for all (x, t) , where $v = v(\mathcal{P}_0)$ is such as defined in Lemma 2.1. Then the following holds.

- (i) For any $\varphi \in X_2^{-1}(0, 1)$ there exists the unique (up to equivalency) solution $V : D \times (0, 1) \rightarrow \mathbf{R}$ of problem (2.9) in the class $C^0(0, 1) \cap X_2^1(0, 1)$.
- (ii) There exists a constant $c = c(\mathcal{P}_q, \mu)$ such that

$$\sup_{t \in [0, 1]} \|V(\cdot, t)\|_{H^0} + \|V\|_{X_2^1(0, 1)} \leq c \|\varphi\|_{X_2^{-1}(0, 1)}. \tag{2.10}$$

- (iii) If $\varphi \in L_\gamma(Q_0)$ for $\gamma > n + 2$, then the functions $V(x, t)$ and $\partial V(x, t)/\partial x_k$ are continuous, bounded, and belong to the Hölder class $\mathcal{H}^{1+\alpha, (1+\alpha)/2}(Q_0)$ with $\alpha \triangleq 1 - (n + 2)/\gamma$, $k = 1, \dots, n$. Moreover, there exists a constant $c = c(\mathcal{P}_{|q|}, \gamma, \mu)$ such that

$$\langle \langle V \rangle \rangle_{Q_0}^{(1+\alpha)} \leq c \|\varphi\|_{L_\gamma(Q_0)}.$$

2.2. Estimates of distances between first exit times

Let $(\Omega, \mathcal{F}, \mathbf{P})$ be a standard probability space. Consider two n -dimensional diffusion processes $y_i(t)$, $i = 1, 2$, such that

$$\begin{cases} dy_i(t) = f_i(y_i(t), t)dt + \beta_i(y_i(t), t)dw(t), & t > 0, \\ y_i(0) = a_i. \end{cases} \tag{2.11}$$

Here $w(t)$ is a standard n -dimensional Wiener process, $f_i : Q \rightarrow \mathbf{R}^n$ and $\beta_i : Q \rightarrow \mathbf{R}^{n \times n}$ are non-random functions. The random vectors a_i with values in \mathbf{R}^n do not depend on $w(\cdot)$. We assume that $(f_i, \beta_i) \in \Theta(\mathcal{P}_0)$, $i = 1, 2$. In particular, we assume that the functions f_i and β_i are continuous, and $b_i(x, t) \triangleq \beta_i(x, t)\beta_i(x, t)^\top \geq \delta I_n$, where I_n is the unit matrix in $\mathbf{R}^{n \times n}$ and $\delta > 0$.

Let us introduce differential operators

$$\begin{aligned} \mathcal{A}_i(t)v(x) &\triangleq \sum_{k=1}^n f_{i,k}(x, t) \frac{\partial v}{\partial x_k}(x) \\ &+ \frac{1}{2} \sum_{k,m=1}^n b_{i,km}(x, t) \frac{\partial^2 v}{\partial x_k \partial x_m}(x), \quad i = 1, 2. \end{aligned} \tag{2.12}$$

Here $x_k, f_{i,k}$, and $b_{i,km}$ are the components of the vectors x, f_i and the matrices $b_i \triangleq \beta_i \beta_i^\top$.

Consider the boundary value problems in Q for $i = 1, 2$

$$\begin{cases} \frac{\partial v_i}{\partial t}(x, t) + \mathcal{A}_i(t)v_i(x, t) + \lambda v_i(x, t) = -1, \\ v_i(x, t)|_{x \in \partial D} = 0, \\ \text{ess sup}_{t>0} \|v_i(\cdot, t)\|_{L_2(D)} < +\infty. \end{cases} \tag{2.13}$$

Let $\tau_i \triangleq \inf\{t \geq 0 : y_i(t) \notin D\}$, and let $\tilde{\tau} \triangleq \tau_1 \wedge \tau_2 = \min(\tau_1, \tau_2)$.

Theorem 2.3. *Let $0 < \lambda < \min(-\ln v_1, -\ln v_2)$, where $v = v_i$ are such as defined in Lemma 2.1 for $(f, \beta) = (f_i, \beta_i)$, $i = 1, 2$. Let v_i be the solutions of problems (2.13), $i = 1, 2$. Then*

$$\mathbf{E} \frac{1}{\lambda} \left[e^{\lambda|\tau_1 - \tau_2|} - 1 \right] \leq \max_{i=1,2} \sup_{(x,t) \in Q} \left| \frac{dv_i}{dx}(x, t) \right| \mathbf{E}|y_1(\tilde{\tau}) - y_2(\tilde{\tau})|. \tag{2.14}$$

Clearly, $|\tau_1 - \tau_2|^p \leq p! \lambda^{-p} [e^{\lambda|\tau_1 - \tau_2|} - 1]$ for $p = 1, 2, \dots$. Therefore, Theorem 2.3 gives an estimate for L_p -distances between two exit times for all $p \in [1, +\infty)$.

Remark 2.2. By Theorem 2.1 (v), the derivatives $\partial v_i(x, t)/\partial x$ are bounded, and the upper bound for $\partial v_i(x, t)/\partial x$ depends only on (\mathcal{P}_0, λ) . Some upper bound can be found following the proof of Theorem 2.1 given below and using constants from this proof with $q(x, t) \equiv \lambda$. The question how to find the minimal upper bound is still open.

Example. Let $n = 1, y_i(t) = a_i + w(t)$, where a_i are non-random, $a_i \in D$, and $D \subset \mathbf{R}$ is a given interval. We have that $\mathbf{E}|y_1(\tilde{\tau}) - y_2(\tilde{\tau})| = |a_1 - a_2|$. Then it follows from Theorem 2.3 that $\mathbf{E}|\tau_1 - \tau_2|^p \leq c_1(p)|a_1 - a_2|$ for all $p \geq 1$, and $\mathbf{E}\{e^{\lambda|\tau_1 - \tau_2|} - 1\} \leq c_2(\lambda)|a_1 - a_2|$ for $\lambda \in (0, -\ln v)$, where v is such as in Lemma 2.1, and where $c_1(p) > 0$ and $c_2(\lambda) > 0$ are constants.

We denote by $[t]$ the integer part of t .

Remark 2.3. If $f_i(x, t) \equiv f_i(x, t + 1), \beta_i(x, t) \equiv \beta_i(x, t + 1)$, then it follows from Theorem 2.2 that $v_i(x, t) \equiv V_i(x, t - [t])$, where V_i is the solution of

$$\begin{cases} \frac{\partial V_i}{\partial t}(x, t) + \mathcal{A}_i(t)V_i(x, t) + \lambda V_i(x, t) = -1, \\ V_i(x, t)|_{x \in \partial D} = 0, \\ V_i(x, 0) \equiv V_i(x, 1). \end{cases}$$

If $f_i(x, t) \equiv f_i(x)$ and $\beta_i(x, t) \equiv \beta_i(x)$ do not depend on t , then $V_i(x, t) \equiv v_i(x, t) \equiv v_i(x)$, and the equations for v_i and V_i became elliptic equations in D .

3. Proofs

Proof of Lemma 2.1. Let $\mathcal{M} \triangleq \Theta(\mathcal{P}_0) \times D \times [0, +\infty)$. Let $\mu = (f, \beta, x, s) \in \mathcal{M}$ be given. Clearly, there exists a finite interval $D_1 \triangleq (d_1, d_2) \subset \mathbf{R}$ and a bounded domain $D_{n-1} \subset \mathbf{R}^{n-1}$ such that $D \subset D_1 \times D_{n-1}$.

Let $\tau_1^{x,s} \triangleq \inf\{t \geq s : y_1^{x,s}(t) \notin D_1\}$, where $y_1^{x,s}(t)$ is the first component of the vector $y^{x,s}(t) = (y_1^{x,s}(t), \dots, y_n^{x,s}(t))$. We have that

$$\mathbf{P}(\tau^{x,s} > s + 1) \leq \mathbf{P}(\tau_1^{x,s} > s + 1) = \mathbf{P}(y_1^{x,s}(t) \in D_1 \ \forall t \in [s, s + 1]). \tag{3.1}$$

Let $M^\mu(t) \triangleq \int_s^t \beta_1(y^{x,s}(r), r)dw(r), t \geq s$, where β_1 is the first row of the matrix β . Let $\widehat{D}_1 \triangleq (d_1 + K_1, d_2 + K_2)$, where $K_1 \triangleq -d_2 - \sup_{x,t} |f_1(x, t)|, K_2 \triangleq -d_1 + \sup_{x,t} |f_1(x, t)|$. Clearly, \widehat{D}_1 depends only on n, D , and c_f . It is easy to see that

$$\mathbf{P}(y_1^{x,s}(t) \in D_1 \ \forall t \in [s, s + 1]) \leq \mathbf{P}(M^\mu(t) \in \widehat{D}_1 \ \forall t \in [s, s + 1]). \tag{3.2}$$

Further,

$$\beta_1(y^{x,s}(t), t)^\top \beta_1(y^{x,s}(t), t) = |\beta_1(y^{x,s}(t), t)|^2 \in [\delta, c_\beta], \tag{3.3}$$

where δ and c_β are such as defined in (2.2). Clearly, $M^\mu(t)$ is a martingale vanishing at s with quadratic variation process

$$[M^\mu]_t \triangleq \int_s^t |\beta_1(y^{x,s}(r), r)|^2 dr, \quad t \geq s.$$

Let $\theta^\mu(t) \triangleq \inf\{r \geq s : [M^\mu]_r > t - s\}$. Note that $\theta^\mu(s) = s$, and the function $\theta^\mu(t)$ is strictly increasing in $t > s$ given (x, s) . By Dambis–Dubins–Schwarz Theorem (see, e.g., Revuz and Yor (1999)), the process $B^\mu(t) \triangleq M(\theta^\mu(t))$ is a Brownian motion vanishing at s , i.e., $B^\mu(s) = 0$, and $M^\mu(t) = B^\mu(s + [M^\mu]_t)$. Clearly,

$$\begin{aligned} \mathbf{P}(M^\mu(t) \in \widehat{D}_1 \ \forall t \in [s, s + 1]) &= \mathbf{P}(B^\mu(s + [M^\mu]_t) \in \widehat{D}_1 \ \forall t \in [s, s + 1]) \\ &\leq \mathbf{P}(B^\mu(r) \in \widehat{D}_1 \ \forall r \in [s, s + [M^\mu]_{s+1}]). \end{aligned} \tag{3.4}$$

By (3.3), $[M^\mu]_{s+1} \geq \delta$ a.s. for all x, s . Hence

$$\mathbf{P}(B^\mu(r) \in \widehat{D}_1 \quad \forall r \in [s, s + [M^\mu]_{s+1}]) \leq \mathbf{P}(B^\mu(r) \in \widehat{D}_1 \quad \forall r \in [s, s + \delta]). \tag{3.5}$$

By (3.1)–(3.2) and (3.4)–(3.5), it follows that

$$\sup_\mu \mathbf{P}(\tau^{x,s} > s + 1) \leq \nu \stackrel{\Delta}{=} \sup_\mu \mathbf{P}(B^\mu(r) \in \widehat{D}_1 \quad \forall r \in [s, s + \delta]),$$

and $\nu = \nu(n, D, c_f, \delta) \in (0, 1)$. This completes the proof. □

Note that the idea to use the Dambis–Dubins–Schwarz Theorem for the proof of Lemma 2.1 was suggested to the author by Tom Salisbury. Originally, the proof was substantially longer.

Let $\mathcal{A}^*(t)$ be the operator formally adjoint to $\mathcal{A}(t)$, i.e.,

$$\mathcal{A}^*(t)u = - \sum_{k=1}^n \frac{\partial}{\partial x_k} \left(f_k(x, t)u(x) \right) + \frac{1}{2} \sum_{k,m=1}^n \frac{\partial^2}{\partial x_k \partial x_m} \left(b_{km}(x, t)u(x) \right). \tag{3.6}$$

Consider the boundary value problem

$$\begin{cases} \frac{\partial p}{\partial t}(x, t) = \mathcal{A}^*(t)p(x, t) + q(x, t)p(x, t) + \xi(x, t), \\ p(x, t)|_{x \in \partial D} = 0, \\ p(x, s) \equiv \rho(x). \end{cases} \tag{3.7}$$

Here $t \geq s$, $q : Q \rightarrow \mathbf{R}$ and $\rho : D \rightarrow \mathbf{R}$ are some functions, the function $q(x, t)$ is measurable and bounded, $\xi|_{D \times [s, T]} \in X_2^{-1}(s, T)$ for all $T \geq s$.

The following Proposition 3.1 presents some facts from Chapter III from Ladyzhenskaya *et al.* (1968) and from Chapter III from Ladyzhenskaya (1985). Estimate (3.8) is "the energy inequality" (3.14) from Ladyzhenskaya (1985).

Proposition 3.1. *Let $0 \leq s < T$, $T - s \leq d$, where $d > 0$ is given. Assume that $(f, \beta) \in \Theta(\mathcal{P}_0)$, $\rho \in H^0$, $\xi \in X_2^{-1}(s, T)$. Then there exists the unique solution $p \in X_2^1(s, T) \cap C^0(s, T)$ of the problem (3.7), and there exists a constant $C = C(\mathcal{P}_q, d)$ such that*

$$\sup_{t \in [s, T]} \|p(\cdot, t)\|_{H^0}^2 + \int_s^T \|p(\cdot, t)\|_{H^1}^2 dt \leq C \left(\|\rho\|_{H^0}^2 + \int_s^T \|\xi(\cdot, t)\|_{H^{-1}}^2 dt \right) \tag{3.8}$$

for all $(f, \beta) \in \Theta(\mathcal{P}_0)$.

In addition to Proposition 3.1, notice that $p^{(s)}(\cdot, T) = p^{(t)}(\cdot, T)$ for $s < t < T$ if $p^{(s)}(\cdot, t) = p^{(t)}(\cdot, t)$. Here $p^{(s)}$ denotes the corresponding solution of (3.7) with $\xi \equiv 0$ given s .

To proceed further, we need some auxiliary lemmas.

We assume below that conditions of Theorem 2.1 are satisfied for q , i.e., we assume that $\sup_{(x,t) \in Q} q(x, t) < -\ln \nu$, and ν is the same as in Lemma 2.1.

Lemma 3.1. *Let p be the solution of (3.7) with $\xi \equiv 0$. Then*

$$\int_D |p(x, t)| dx \leq C_0 e^{-\omega_*(t-s)} \|\rho\|_{H^0} \quad \forall t \in [s, +\infty), \tag{3.9}$$

$$|p(x, t)| \leq C_1 e^{-\omega_*(t-s)} \|\rho\|_{H^0} \quad \forall t \in [s + 1, +\infty), \tag{3.10}$$

where $\omega_* \triangleq -\ln v - \sup_{(x,t) \in Q} q(x, t)$, and $C_i = C_i(\mathcal{P}_q)$ are constants that do not depend on s, t, ρ and depend on \mathcal{P}_q only, $i = 0, 1$.

Proof. By linearity of the problem, it suffices to consider ρ such that $\rho(x) \geq 0$ and $\int_D \rho(x) dx = 1$. Let $p_0(x, t) \triangleq p(x, t)e^{-\lambda(t-s)}$ and $q_0(x, t) \triangleq q(x, t) - \lambda$, where $\lambda \triangleq \sup_{(x,t) \in Q} q(x, t)$. Clearly, $q_0(x, t) \leq 0$ and

$$\begin{cases} \frac{\partial p_0}{\partial t}(x, t) = \mathcal{A}^*(t)p_0(x, t) + q_0(x, t)p_0(x, t), \\ p_0(x, t)|_{x \in \partial D} = 0, \\ p_0(x, s) \equiv \rho(x). \end{cases}$$

Therefore, $p_0(x, t)$ is the probability density function of the process $y^{a,s}(t)$ under assumption that this process is absorbed at ∂D and is killed inside D with the rate $|q_0(x, t)|$, where a is a random vector independent on $w(\cdot)$ with the probability density function ρ . Hence $0 \leq p_0(x, t) \leq \pi(x, t)$, where $\pi(x, t)$ is the probability density function of the process $y^{a,s}(t)$ under assumption that this process is absorbed at ∂D without being killed inside D , i.e.,

$$\begin{cases} \frac{\partial \pi}{\partial t}(x, t) = \mathcal{A}^*(t)\pi(x, t), \\ \pi(x, t)|_{x \in \partial D} = 0, \\ \pi(x, s) \equiv \rho(x). \end{cases}$$

Because of absorption at ∂D , we have

$$\int_D \pi(x, t) dx \leq \int_D \pi(x, r) dx \quad \forall r, t \in \mathbf{R} : s \leq r \leq t.$$

By Lemma 2.1, it follows that

$$\int_D \pi(x, t + 1) dx \leq v \int_D \pi(x, t) dx \quad \forall t \geq s.$$

Hence

$$\begin{aligned} \int_D |p_0(x, t)| dx &= \int_D p_0(x, t) dx \leq \|\pi(\cdot, t)\|_{L_1(D)} \\ &\leq \|\pi(\cdot, s + \lfloor t - s \rfloor)\|_{L_1(D)} \\ &\leq v \|\pi(\cdot, s + \lfloor t - s \rfloor - 1)\|_{L_1(D)} \\ &\leq v^2 \|\pi(\cdot, s + \lfloor t - s \rfloor - 2)\|_{L_1(D)} \leq \dots \leq v^{\lfloor t - s \rfloor} \|\rho\|_{L_1(D)} \\ &= e^{\lfloor t - s \rfloor \ln v} \|\rho\|_{L_1(D)}, \end{aligned} \tag{3.11}$$

where $\lfloor t \rfloor$ denotes the integer part of t . Then (3.9) follows.

Let us prove (3.10). Let $\Delta \triangleq \{(t, s) : t \geq s \geq 0\}$, and let $g(\cdot) : D^2 \times \Delta \rightarrow \mathbf{R}$ be the Green's function for the equation (3.7) such that if $\xi \equiv 0$ then

$$p(x, t) = \int_D g(x, y, t, s)p(y, s)dy, \quad t \geq s \geq 0. \tag{3.12}$$

Let $G(x, y, t, s)$ be the fundamental solution of problem (3.7) without the boundary condition on ∂D (i.e., for $D = \mathbf{R}^n$); the order of independent variables for G is similar to (3.12). By Lemma 7 from Aronson (1968), it follows that $|g(x, y, t, s)| \leq |G(x, y, t, s)| (\forall x, y, t, s)$. Using estimates from Aronson (1967), we obtain

$$|g(x, y, t + 1, t)| \leq |G(x, y, t + 1, t)| \leq c \quad \forall x, y \in D, t \geq 0, \tag{3.13}$$

where $c = c(\mathcal{P}_q)$ is a constant. By (3.11) and (3.13), it follows (3.10). This completes the proof of Lemma 3.1. □

Let us introduce linear normed spaces $Z^k(s, +\infty)$ of functions $u : (s, +\infty) \rightarrow H^k$ with finite norm

$$\|u\|_{Z^k(s, +\infty)} = \sum_{m=0}^{+\infty} \left(\int_{s+m}^{s+m+1} \|u(\cdot, t)\|_{H^k}^2 dt \right)^{1/2}.$$

Lemma 3.2. *Let $s \geq 0$, let $\rho \in H^0$, and let $\xi \in X_1^0(s, +\infty) \cup X_2^{-1}(s, +\infty)$ be such that $\xi(\cdot, t) \equiv 0$ for $t > s + 1$. Then there exists the solution $p \in X_1^1(s, +\infty) \cap C^0(s, +\infty)$ of problem (3.7). This solution is unique up to equivalency, and*

$$\|p\|_{X_1^1(s, s+1)} + \|p\|_{C^0(s, s+1)} \leq c_1(\|\rho\|_{H^0} + \|\xi\|_{X_1^0(s, s+1)}), \tag{3.14}$$

$$\|p\|_{X_2^1(s, s+1)} + \|p\|_{C^0(s, s+1)} \leq c_2(\|\rho\|_{H^0} + \|\xi\|_{X_2^{-1}(s, s+1)}), \tag{3.15}$$

$$\|p\|_{X_2^1(s+1, +\infty)} \leq c_3\|p(\cdot, s + 1)\|_{H^0}, \tag{3.16}$$

$$\|p\|_{X_1^1(s+1, +\infty)} + \|p\|_{C^0(s+1, +\infty)} \leq c_4\|p(\cdot, s + 1)\|_{H^0}, \tag{3.17}$$

$$\|p\|_{Z^1(s+1, +\infty)} \leq c_5\|p(\cdot, s + 1)\|_{H^0}, \tag{3.18}$$

where $c_i = c_i(\mathcal{P}_q) > 0$ are constants that do not depend on s and depend on \mathcal{P}_q only, $i = 1, \dots, 5$.

Proof of Lemma 3.2. Let us prove (3.14). For any $T \geq s$ and any $\varepsilon \in (0, \delta)$, we have

$$\begin{aligned} \|p(\cdot, T)\|_{H^0} - \|p(\cdot, s)\|_{H^0} &= \int_s^T \|p(\cdot, t)\|_{H^0}^{-1}(p(\cdot, t), \mathcal{A}^*p(\cdot, t) \\ &\quad + q(\cdot, t)p(\cdot, t) + \xi(\cdot, t))_{H^0} dt \end{aligned}$$

$$\begin{aligned}
 &= \int_s^T \|p(\cdot, t)\|_{H^0}^{-1} \left\{ -\frac{1}{2} \sum_{i,j=1}^n \left[\left(\frac{\partial p}{\partial x_i}(\cdot, t), b_{ij}(\cdot, t) \frac{\partial p}{\partial x_j}(\cdot, t) \right)_{H^0} \right. \right. \\
 &\quad \left. \left. + \frac{1}{2} \left(p(\cdot, t), \frac{\partial b_{ij}}{\partial x_j}(\cdot, t) \frac{\partial p}{\partial x_i}(\cdot, t) \right)_{H^0} \right] \right. \\
 &\quad \left. + \sum_{i=1}^n \left(\frac{\partial p}{\partial x_i}(\cdot, t), f_i(\cdot, t) p(\cdot, t) \right)_{H^0} \right. \\
 &\quad \left. + \left(\xi(\cdot, t) + q(\cdot, t) p(\cdot, t), p(\cdot, t) \right)_{H^0} \right\} dt \\
 &\leq \int_s^T \|p(\cdot, t)\|_{H^0}^{-1} \sum_{i=1}^n \left\{ \frac{1}{2} (\varepsilon - \delta) \left\| \frac{\partial p}{\partial x_i}(\cdot, t) \right\|_{H^0}^2 \right. \\
 &\quad \left. + c(\varepsilon) \|p(\cdot, t)\|_{H^0}^2 \right. \\
 &\quad \left. + \|p(\cdot, t)\|_{H^0} \|\xi(\cdot, t)\|_{H^0} \right\} dt \tag{3.19}
 \end{aligned}$$

Here the constant $c(\varepsilon) = c(\varepsilon, \mathcal{P}_q) > 0$ depends only on ε and $\mathcal{P}_q \triangleq (\mathcal{P}_0, \sup q(x, t))$. We had used elementary inequality $2\alpha\beta \leq \varepsilon\alpha^2 + \varepsilon^{-1}\beta^2$ ($\forall \alpha, \beta, \varepsilon \in \mathbf{R}, \varepsilon > 0$), and inequality

$$\left(\frac{\partial p}{\partial x_i}(\cdot, t), F(\cdot) p(\cdot, t) \right)_{H^0} \leq \frac{\varepsilon}{2} \left\| \frac{\partial p}{\partial x_i}(\cdot, t) \right\|_{H^0}^2 + \frac{1}{2\varepsilon} \|F(\cdot)\|_{L_\infty(D)} \|p(\cdot, t)\|_{H^0},$$

where $F(\cdot) : D \rightarrow \mathbf{R}$ is an arbitrary measurable bounded function.

By Poincaré - Friedrichs inequality (see, e.g., Yosida (1965)), it follows that there exist a constant $\kappa = \kappa(D) > 0$ such that

$$\|p(\cdot, t)\|_{H^0}^{-1} \sum_{i=1}^n \left\| \frac{\partial p}{\partial x_i}(\cdot, t) \right\|_{H^0}^2 \geq \kappa \|p(\cdot, t)\|_{H^1}.$$

By (3.19), it follows that

$$\begin{aligned}
 &\|p(\cdot, T)\|_{H^0} + \bar{c}_1 \int_s^T \|p(\cdot, t)\|_{H^1} dt \\
 &\leq \|\rho\|_{H^0} + \bar{c}_2 \left(\int_s^T \|p(\cdot, t)\|_{H^0} dt + \int_s^T \|\xi(\cdot, t)\|_{H^0} dt \right) \quad \forall T \geq s. \tag{3.20}
 \end{aligned}$$

Here $\bar{c}_i = \bar{c}_i(\mathcal{P}_q)$ are constants that do not depend on $T \in [s, +\infty)$ for $i = 1, 2$. By Gronwall's inequality, inequality (3.20) applied for $T \in [s, s + 1]$ implies (3.14).

Similarly (3.19)–(3.20), one can derive

$$\begin{aligned}
 &\|p(\cdot, T)\|_{H^0}^2 + \hat{c}_1 \int_s^T \|p(\cdot, t)\|_{H^1}^2 dt \\
 &\leq \|p(\cdot, s)\|_{H^0}^2 + \hat{c}_2 \left(\int_s^T \|p(\cdot, t)\|_{H^0}^2 dt + \int_s^T \|\xi(\cdot, t)\|_{H^{-1}}^2 dt \right) \quad \forall T \geq s. \tag{3.21}
 \end{aligned}$$

Constants $\widehat{c}_i = \widehat{c}_i(\mathcal{P}_q) > 0$ do not depend on $T \in [s, +\infty)$. By Gronwall’s inequality again, inequality (3.21) with $T \in [s, s + 1]$ implies (3.15) (In fact, this is the estimate from Proposition 3.1, or a reformulation of “the energy inequality” (3.14) from Ladyzhenskaya (1985)).

Let us prove (3.16)–(3.18). Remind that $\xi(x, t) \equiv 0$ for $t > s + 1$. By Lemma 3.1,

$$|p(x, t)| \leq C_1 e^{-\omega_*(t-s-1)} \|p(\cdot, s + 1)\|_{H^0} \quad (\forall t \geq s + 1),$$

where $C_1 = C_1(\mathcal{P}_q) > 0$ is a constant from (3.10), $\omega_* = -\ln v - \max q(x, t)$. Then

$$\begin{aligned} \|p(\cdot, t)\|_{H^0} &\leq C_1 e^{-\omega_*(t-s-1)} \|p(\cdot, s + 1)\|_{H^0}, \\ \int_{s+1}^{+\infty} \|p(\cdot, t)\|_{H^0}^2 dt &\leq C_2 \|p(\cdot, s + 1)\|_{H^0}^2, \\ \int_{s+1}^{+\infty} \|p(\cdot, t)\|_{H^0} dt &\leq C_3 \|p(\cdot, s + 1)\|_{H^0}. \end{aligned} \tag{3.22}$$

Here $C_i = C_i(\mathcal{P}_q) > 0$ are constants. Then (3.16) follows from (3.21) and (3.22). Further, (3.17) follows from (3.20) and (3.22). By (3.21)–(3.22),

$$\begin{aligned} \widehat{c}_1 \int_{s+m}^{s+m+1} \|p(\cdot, t)\|_{H^1}^2 dt &\leq \|p(\cdot, s + m)\|_{H^0}^2 + \widehat{c}_2 \int_{s+m}^{s+m+1} \|p(\cdot, t)\|_{H^0}^2 dt \\ &\leq C_1^2 \left[e^{-2\omega_*(m-1)} + \widehat{c}_2 \int_{s+m}^{s+m+1} e^{-2\omega_*(t-s-1)} dt \right] \\ &\quad \times \|p(\cdot, s + 1)\|_{H^0}^2 \\ &\leq C_* e^{-2\omega_* m} \|p(\cdot, s + 1)\|_{H^0}^2 \quad \forall m = 1, 2, \dots \end{aligned}$$

Here $C_* = C_*(\mathcal{P}_q) > 0$ is a constant that does not depend on m . Then (3.18) follows. This completes the proof of Lemma 3.2. □

Note that (3.8) can be derived by the following way. Similarly (3.19)–(3.20), one can derive (3.21). By Gronwall’s inequality, inequality (3.21) implies (3.8).

Let $0 \leq s < T$, let $\mathcal{Q} \triangleq D \times (s, T)$, and let $\gamma \geq 1$. Introduce linear normed spaces $\mathcal{W}_\gamma(s, T)$ of functions $u : [s, T] \rightarrow W_\gamma^2(D)$ that belong to $L_\gamma([s, T], \vec{B}_1, \ell_1, W_\gamma^2(D))$ and such that $\frac{\partial u}{\partial t}$ belong to $L_\gamma([s, T], \vec{B}_1, \ell_1, L_\gamma(D))$, with finite norm

$$\|u\|_{\mathcal{W}_\gamma(s, T)} = \left(\int_s^T \|u(\cdot, t)\|_{W_\gamma^2(D)}^\gamma dt \right)^{1/\gamma} + \left(\int_s^T \left\| \frac{\partial u}{\partial t}(\cdot, t) \right\|_{L_\gamma(D)}^\gamma dt \right)^{1/\gamma}.$$

It is easy to see that $\mathcal{W}_\gamma(s, T) \subset C([s, T]; L_\gamma(D))$, and this embedding is continuous. Moreover, $\mathcal{W}_\gamma(s, T) = W_\gamma^{2,1}(\mathcal{Q})$, meaning the natural bijection such that the norms are equivalent.

The space $W_\gamma^l(D)$ with non-integer l will be used below. It is a Banach space consisting of the elements of $W_\gamma^{[l]}(D)$ with finite norm

$$\|u\|_{W_\gamma^l(D)} \triangleq \|u\|_{W_\gamma^{[l]}(D)} + \sum_{j:|j|=|l|} \left(\int_D dx \int_D |D_x^j u(x) - D_y^j u(y)|^\gamma \frac{dy}{|x-y|^{n+\gamma(l-[l])}} \right)^{1/\gamma}.$$

Here $[l]$ is the integer part of l , $j = (j_1, \dots, j_n)$, where $j_k \geq 0$ are integers, $|j| = \sum_k j_k$,

$$D_x^j u(x) = \frac{\partial^{|j|} u}{\partial x_1^{j_1} \dots \partial x_n^{j_n}}.$$

(See, e.g., Ladyzhenskaya *et al.* (1968), p. 70, and Adams (1975), p. 214).

Consider the boundary value problem in \mathcal{Q}

$$\begin{cases} \frac{\partial u}{\partial t}(x, t) + \mathcal{A}(t)u(x, t) + q(x, t)u(x, t) = -\varphi(x, t), \\ u(x, t)|_{x \in \partial D} = 0, \\ u(x, T) \equiv \Phi(x). \end{cases} \tag{3.23}$$

Here $q : \mathcal{Q} \rightarrow \mathbf{R}$, $\varphi : \mathcal{Q} \rightarrow \mathbf{R}$ and $\Phi : D \rightarrow \mathbf{R}$ are some measurable functions, the function $q(x, t)$ is bounded.

Let $\theta \in (s, T)$, and let $\mathcal{Q}_\theta \triangleq D \times (s, \theta)$.

Lemma 3.3. *Let $0 \leq s < \theta < T$ and $\gamma \geq 2$. Assume that $(f, \beta) \in \Theta(\mathcal{P}_0)$, $\varphi \in X_2^{-1}(s, T)$, $\Phi \in H^0$, $T - s \leq d$, and $T - \theta \geq d_0$, where $d > 0$ and $d_0 > 0$ are given. Then*

- (i) *There exists the unique solution $u \in C^0(s, T) \cap X_2^1(s, T)$ of problem (3.23), and there exists a constant $C = C(\mathcal{P}_q, d) > 0$ such that*

$$\|u\|_{C^0(s,T)} + \|u\|_{X_2^1(s,T)} \leq C \left(\|\Phi\|_{H^0} + \|\varphi\|_{X_2^{-1}(s,T)} \right) \tag{3.24}$$

for all $(f, \beta) \in \Theta(\mathcal{P}_0)$.

- (ii) *Let $\rho \in H^0$ be arbitrary, and let p be the solution of (3.7), where $\xi \equiv 0$. Then*

$$(u(\cdot, T), p(\cdot, T))_{H^0} - (u(\cdot, s), p(\cdot, s))_{H^0} = - \int_s^T (\varphi(\cdot, t), p(\cdot, t))_{H^0} dt.$$

- (iii) *If $\varphi \in L_2(\mathcal{Q})$ and $\Phi \in H^1$, then $u \in C^1(s, T) \cap X_2^2(s, T)$, and there exists a constant $C = C(\mathcal{P}_{|q|}, d) > 0$ such that*

$$\|u\|_{C^1(s,T)} + \|u\|_{X_2^2(s,T)} \leq C \left(\|\Phi\|_{H^1} + \|\varphi\|_{L_2(\mathcal{Q})} \right) \tag{3.25}$$

for all $(f, \beta) \in \Theta(\mathcal{P}_0)$.

(iv) If $\varphi \in L_\gamma(\mathcal{Q})$ and $\Phi \in W_\gamma^{2-2/\gamma}(D) \cap H^1$, then the solution u of problem (3.23) belongs to $\mathcal{W}_\gamma(s, T)$, and there exists a constant $C = C(\mathcal{P}_{|q|}, d, \gamma) > 0$ such that

$$\|u\|_{\mathcal{W}_\gamma(s, T)} \leq C \left(\|\Phi\|_{W_\gamma^{2-2/\gamma}(D)} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right) \tag{3.26}$$

for all $(f, \beta) \in \Theta(\mathcal{P}_0)$.

(v) If $\varphi \in L_\gamma(\mathcal{Q})$, then the solution u is such that $u|_{\mathcal{Q}_\theta} \in \mathcal{W}_\gamma(s, \theta)$, and there exists a constant $C = C(\mathcal{P}_{|q|}, d, \gamma) > 0$ such that

$$\|u|_{\mathcal{Q}_\theta}\|_{\mathcal{W}_\gamma(s, \theta)} \leq C \left(\|\Phi\|_{H^0} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right) \tag{3.27}$$

for all $(f, \beta) \in \Theta(\mathcal{P}_0)$.

(vi) If $\gamma > n + 2$ and $\varphi \in L_\gamma(\mathcal{Q})$, then $u(x, t)|_{\mathcal{Q}_\theta}$ and $\frac{\partial u}{\partial x_k}(x, t)|_{\mathcal{Q}_\theta}$, $k = 1, \dots, n$, are continuous and belong to Hölder class $\mathcal{H}^{1+\alpha, (1+\alpha)/2}(\mathcal{Q}_\theta)$ for $\alpha = 1 - (n + 2)/\gamma$. Moreover, there exists a constant $C = C(\mathcal{P}_{|q|}, d, d_0, \gamma) > 0$ such that

$$\langle\langle u|_{\mathcal{Q}_\theta} \rangle\rangle_{\mathcal{Q}_\theta}^{(1+\alpha)} \leq C \left(\|\Phi\|_{H^0} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right)$$

for all $(f, \beta) \in \Theta(\mathcal{P}_0)$.

Remark 3.1. Under the assumptions of statement (iv) in Lemma 3.3, $u \in W_\gamma^{2,1}(\mathcal{Q})$, and $\|u\|_{W_\gamma^{2,1}(\mathcal{Q})} \leq \text{const} \|u\|_{\mathcal{W}_\gamma(s, T)}$, because there is a natural bijection between $W_\gamma^{2,1}(\mathcal{Q})$ and $\mathcal{W}_\gamma(s, T)$ such that the norms are equivalent. Under assumptions of statement (v), $u|_{\mathcal{Q}_\theta} \in W_\gamma^{2,1}(\mathcal{Q}_\theta)$, and $\|u|_{\mathcal{Q}_\theta}\|_{W_\gamma^{2,1}(\mathcal{Q}_\theta)} \leq \text{const} \|u\|_{\mathcal{W}_\gamma(s, \theta)}$.

Proof of Lemma 3.3. Statement (i) follows from inequality (3.14) from Ladyzhenskaya (1985). Statement (ii) follows from the fact that the parabolic equations in (3.7) and (3.23) are adjoint, and from the equations for $\partial u/\partial t$ and $\partial p/\partial t$. Statement (iii) follows from Theorem 1.2 from Dokuchaev (1997). (Note that statement (iii) can be also derived from Theorem 6.1 and Remark 6.3 from Ladyzhenskaya *et al.* (1968)) (pp. 178-180). More precisely, this statement follows from the inequality (6.25) from Ladyzhenskaya *et al.* (1968), p. 180, and from the inequality (6.29) from Ladyzhenskaya (1985). In fact, Theorem 6.1 from Ladyzhenskaya *et al.* (1968) deals with a special case of (f, q) , but it is not really important).

Statement (iv) is a special case of Theorem 9.1, Chapter IV, from Ladyzhenskaya *et al.* (1968). Formally, this theorem requires that $\Phi \in W_\gamma^{2-2/\gamma}(D)$ and $\Phi|_{\partial D} = 0$. However, these conditions can be easily replaced by our condition $\Phi \in W_\gamma^{2-2/\gamma}(D) \cap H^1$. Let us show this. Let $\Phi \in W_\gamma^{2-2/\gamma}(D) \cap H^1$. Clearly, there exists a sequence $\{\Phi_i\}_{i=1}^{+\infty} \subset C^2(D)$ such that $\Phi_i|_{\partial D} = 0$ ($\forall i$), and $\Phi_i \rightarrow \Phi$ in both spaces $W_\gamma^{2-2/\gamma}(D)$ and H^1 as $i \rightarrow \infty$. Let u_i be the solution of problem (3.23) with $\Phi = \Phi_i$. By Theorem 9.1, Chapter IV, from Ladyzhenskaya *et al.* (1968), the constant C in (3.26) does not depend on $\Phi = \Phi_i$. Therefore, the sequence $\{u_i\}_{i=1}^{+\infty}$ is a Cauchy sequence in $\mathcal{W}_\gamma(s, T)$ and has a limit in this space. By statement (iii), $u_i \rightarrow u$ in $\mathcal{C}^1(s, T)$, where $u \in \mathcal{C}^1(s, T) \cap X_2^1(s, T)$ is the solution of (3.23) given Φ . Hence $u \in \mathcal{W}_\gamma(s, T)$ and (3.26) is satisfied. This completes the proof of statement (iv).

Let us prove statement (v). Consider the following sequences:

$$\begin{aligned} h_1 &= 2, & h_m &\triangleq h_{m-1} \frac{n+2}{n}, \\ \chi_1 &= 2, & \chi_m &= 2 - \frac{2}{h_m}, \end{aligned} \quad m = 2, 3, \dots$$

It is easy to see that

$$\begin{aligned} \chi_m &= 2 - \frac{n}{h_{m-1}} + \frac{n}{h_m}, \quad \chi_m > 0, \quad h_{m+1} > h_m, \\ h_m &\rightarrow \infty \quad \text{as } m \rightarrow +\infty. \end{aligned}$$

Clearly, there exists $N = N(n)$ such that $h_N \geq \gamma$ and $h_m < \gamma$ for all $m < N$. Let $s_m \triangleq T - (m - 1)(T - \theta)/N$, $m = 1, \dots, N + 1$. It is easy to see that

$$\theta = s_{N+1} < \dots < s_{m+1} < s_m < \dots < s_1 = T.$$

Let us prove that there exists a set $\{t_m\}_{m=1}^N \subset [\theta, T]$ such that

$$\begin{aligned} t_m &\in (s_{m+1}, s_m], \quad u(\cdot, t_m) \in W_{h_m}^2(D) \cap H^1, \\ \|u(\cdot, t_m)\|_{W_{h_m}^2(D)} &\leq C \left(\|\Phi\|_{H^0} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right), \end{aligned} \quad (3.28)$$

where $C = C(\mathcal{P}_{|q|}, \gamma, d)$, $m = 1, \dots, N$. Note that we allow that $\{t_m\}_{m=1}^N$ can depend on (Φ, φ) .

First, let us prove that (3.28) is satisfied for $m = 1$ for some t_1 . Clearly, $H^2 \subset W_2^2(D) = W_{h_1}^1(D)$, and this embedding is continuous. Therefore, it suffices to prove that there exists $t_1 \in (s_2, s_1] = (s_2, T]$ such that

$$u(\cdot, t_1) \in H^2, \quad \|u(\cdot, t_1)\|_{H^2} \leq C \left(\|\Phi\|_{H^0} + \|\varphi\|_{L_2(\mathcal{Q})} \right), \quad (3.29)$$

where $C = C(\mathcal{P}_{|q|}, d, d_0)$.

Let $h \triangleq (s_2 + s_1)/2 = (s_2 + T)/2$. By statement (i), it follows that $u \in C^0(s, T) \cap X_2^1(s, T)$, and

$$\int_s^T \|u(\cdot, t)\|_{H^1}^2 dt \leq C_1 \left(\|\Phi\|_{H^0}^2 + \|\varphi\|_{L_2(\mathcal{Q})}^2 \right),$$

where $C_1 = C_1(\mathcal{P}_q, d) > 0$. Hence

$$\inf_{r \in [h, T]} \|u(\cdot, r)\|_{H^1}^2 \leq \frac{1}{T-h} \int_h^T \|u(\cdot, t)\|_{H^1}^2 dt \leq \frac{C_1}{T-h} \left(\|\Phi\|_{H^0}^2 + \|\varphi\|_{L_2(\mathcal{Q})}^2 \right).$$

By statement (iii), if $r \in [h, T]$ is such that $u(\cdot, r) \in H^1$, then $u \in \mathcal{W}_2(s, r)$, and

$$\int_s^h \|u(\cdot, t)\|_{H^2}^2 dt \leq \int_s^r \|u(\cdot, t)\|_{H^2}^2 dt \leq C_2 \left(\|u(\cdot, r)\|_{H^1}^2 + \|\varphi\|_{L_2(\mathcal{Q})}^2 \right),$$

where $C_2 = C_2(\mathcal{P}_{|q|}, d, d_0) > 0$. Hence

$$\begin{aligned} \inf_{r \in [\widehat{s}_2, h]} \|u(\cdot, r)\|_{H^2}^2 &\leq \frac{1}{h - \widehat{s}_2} \int_{\widehat{s}_2}^h \|u(\cdot, t)\|_{H^2}^2 dt \\ &\leq C_3 \left(\inf_{r \in [h, T]} \|u(\cdot, r)\|_{H^1}^2 + \|\varphi\|_{L_2(\mathcal{Q})}^2 \right) \leq C_4 \left(\|\Phi\|_{H^0}^2 + \|\varphi\|_{L_2(\mathcal{Q})}^2 \right), \end{aligned}$$

where $\widehat{s}_2 \triangleq (s_2 + h)/2$, and where $C_i = C_i(\mathcal{P}_{|q|}, d, d_0) > 0, i = 3, 4$. Thus, there exists $t_1 \in (s_2, s_1]$ such that (3.29) is satisfied for $m = 1$. Hence (3.28) is satisfied for $m = 1$.

Let us show that if there exists t_k such that (3.28) is satisfied for $m = k$ with $k \in \{2, \dots, N - 1\}$, then there exists t_{k+1} such that (3.28) is satisfied with $m = k + 1$.

Let us now assume that there exists $t_k \in (s_{k+1}, s_k]$ such that (3.28) holds.

By the direct embedding theorem, if $\chi \triangleq \psi - n/g + n/h > 0$ and $h > g$, then $W_g^\psi(D) \subset W_h^\chi(D)$, and the embedding is continuous (see, e.g., Theorem 7.58 from Adams (1975), p. 218; the case of bounded domain is covered by Theorem 4.26, on page 84 of the cited book; see also related comments before Theorem 7.58 and Remark 7.49 there). We have that

$$W_{h_{m-1}}^2(D) \subset W_{h_m}^{\chi_m}(D), \quad m = 2, 3, \dots \tag{3.30}$$

and the embedding is continuous. Thus, $W_{h_k}^2(D) \subset W_{h_{k+1}}^{\chi_{k+1}}(D)$, and $u(\cdot, t_k) \in W_{h_{k+1}}^{\chi_{k+1}}(D)$. Moreover, $\|u(\cdot, t)\|_{W_{h_{k+1}}^{\chi_{k+1}}(D)} \leq C \|u(\cdot, t)\|_{W_{h_k}^2(D)}$ for any t such that $u(\cdot, t) \in W_{h_k}^2(D)$, where $C = C(n, D, k, \gamma) > 0$ is a constant.

Let $R_m \triangleq D \times (s, t_m)$ and $\mathcal{Q}_{s_m} \triangleq D \times (s, s_m), m = 1, \dots, N + 1$. By statement (iv),

$$\|u|_{R_k}\|_{\mathcal{W}_{h_k}(s, t_k)} \leq C \left(\|u(\cdot, t_k)\|_{W_{h_k}^{\chi_k}(D)} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right),$$

where $C = C(\mathcal{P}_{|q|}, d, d_0, h_k) > 0$ is a constant. By this estimate and (3.30), we have

$$\begin{aligned} \|u|_{\mathcal{Q}_{s_{k+1}}}\|_{\mathcal{W}_{h_{k+1}}(s, s_{k+1})} &\leq C_1 \left(\inf_{r \in [s_{k+1}, t_k]} \|u(\cdot, r)\|_{W_{h_{k+1}}^{\chi_{k+1}}(D)} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right) \\ &\leq C_2 \left(\inf_{r \in [s_{k+1}, t_k]} \|u(\cdot, r)\|_{W_{h_k}^2(D)} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right) \\ &\leq C_3 \left(\frac{1}{t_k - s_{k+1}} \int_{s_{k+1}}^{t_k} \|u(\cdot, t)\|_{W_{h_k}^2(D)} dt + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right) \\ &\leq C_4 \left(\left[\int_{s_{k+1}}^{t_k} \|u(\cdot, t)\|_{W_{h_k}^{h_k}(D)} dt \right]^{1/h_k} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right) \\ &\leq C_5 \left(\|u\|_{\mathcal{W}_{h_k}(s, t_k)} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right) \\ &\leq C_6 \left(\|u(\cdot, t_k)\|_{W_{h_k}^{\chi_k}(D)} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right) \\ &\leq C_7 \left(\|\Phi\|_{H^0} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right), \end{aligned} \tag{3.31}$$

where $C_i = C_i(\mathcal{P}_{|q|}, d, d_0, h_k, \gamma) > 0$ are constants, $i = 1, \dots, 7$.

Further, we have

$$\begin{aligned} & \inf_{r \in [\widehat{s}_{k+2}, s_{k+1}]} \|u(\cdot, r)\|_{W_{h_{k+1}}^2(D)} \\ & \leq C_1 \frac{1}{\widehat{s}_{k+1} - s_{k+2}} \left(\int_{\widehat{s}_{k+2}}^{s_{k+1}} \|u(\cdot, t)\|_{W_{h_{k+1}}^2(D)}^{h_{k+1}} dt \right)^{1/h_{k+1}} \\ & \leq C_2 \|u\|_{\mathcal{Q}_{s_{k+1}}} \| \mathcal{W}_{h_{k+1}}(s, s_{k+1}) \cdot \end{aligned} \tag{3.32}$$

where $\widehat{s}_{k+2} \triangleq (s_{k+2} + s_{k+1})/2$, and where $C_i = C_i(n, D, d, d_0) > 0$. By statement (iii),

$$u(\cdot, t) \in H^1 \quad \forall t \leq t_1. \tag{3.33}$$

By (3.31)–(3.33), it follows that there exists $t_{k+1} \in (s_{k+2}, s_{k+1}]$ such that (3.28) is satisfied for $m = k + 1$. Therefore, we have proved that (3.28) is satisfied for all $m = 1, \dots, N$.

Further, we have that $W_{h_1}^2(D) = W_{h_1}^{\chi_1}(D)$ and $W_{h_m}^2(D) \subset W_{h_{m-1}}^2(D) \subset W_{h_m}^{\chi_m}(D)$, $m = 2, 3, \dots, N + 1$, and the embedding is continuous. By statement (iv), (3.28) implies that $u|_{R_m} \in \mathcal{W}_{h_m}(s, t_m)$, and

$$\begin{aligned} \|u|_{R_m}\|_{\mathcal{W}_{h_m}(s, t_m)} & \leq C_1 \left(\|u(\cdot, t_m)\|_{W_{h_m}^{\chi_m}(D)} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right) \\ & \leq C_2 \left(\|u(\cdot, t_m)\|_{W_{h_m}^2(D)} + \|\varphi\|_{L_\gamma(\mathcal{Q})} \right), \quad m = 1, \dots, N, \end{aligned}$$

where $C_i = C_i(\mathcal{P}_q, d, d_0, h_k, \gamma) > 0$ are constants, $i = 1, 2$. Remind that $\mathcal{Q}_\theta = \mathcal{Q}_{s_{N+1}} \subset R_N$ and $h_N > \gamma$. Thus, statement (v) follows from this estimate with $m = N$. This completes the proof of statement (v).

Let us prove statement (vi). Note that $u|_{\mathcal{Q}_\theta} \in \mathcal{W}_\gamma(s, \theta)$, and there is the natural bijection between $W_\gamma^{2,1}(\mathcal{Q}_\theta)$ and $\mathcal{W}_\gamma(s, \theta)$ such that the norms are equivalent. Then statement (vi) follows from (v) and from continuity of embedding of $W_\gamma^{2,1}(\mathcal{Q}_\theta)$ to the Hölder class $\mathcal{H}^{1+\alpha, (1+\alpha)/2}(\mathcal{Q}_\theta)$ with $\gamma > n+2$ (see, e.g., Lemma 3.3 of Chapter II from Ladyzhenskaya *et al.* (1968)). This completes the proof of Lemma 3.3. \square

Proof of Theorem 2.1. Let $L_{s,t}^* : H^0 \rightarrow H^0$ be the operator such that $p(\cdot, t) = L_{s,t}^* \rho$, where p is the solution of (3.7) with $\xi \equiv 0$, and where $\rho \in H^0$, $0 \leq s \leq t$. By Lemma 3.2, this operator is continuous, and $\|p\|_{Z^1(s, +\infty)} \leq C \|\rho\|_{H^0}$ for $p = L_{s,t}^* \rho$, where $C = C(\mathcal{P}_q)$ is a constant.

Given $\varphi \in Y^{-1}(0, +\infty)$ and $s \geq 0$, let $v(s) \in H^0$ be defined such that

$$(v(s), \rho)_{H^0} = \int_s^{+\infty} (\varphi(\cdot, t), L_{s,t}^* \rho)_{H^0} dt \quad \forall \rho \in H^0. \tag{3.34}$$

Note that $v(s) \in H^0$ is well defined for all $s \geq 0$. This can be seen from the following. Let $B_{H^0} \triangleq \{\rho \in H^0 : \|\rho\|_{H^0} \leq 1\}$. By (3.18), it follows that

$$\begin{aligned} & \sup_{\rho \in B_{H^0}} \int_s^{+\infty} (\varphi(\cdot, t), L_{s,t}^* \rho)_{H^0} dt \\ & \leq \|\varphi\|_{Y^{-1}(s, +\infty)} \sup_{\rho \in B_{H^0}} \|L_{s,t}^* \rho\|_{Z^1(s, +\infty)} \leq c \|\varphi\|_{Y^{-1}(0, +\infty)}, \end{aligned}$$

where $c = c(\mathcal{P}_q)$ is a constant. Therefore,

$$\sup_{s \geq 0} \|v(s)\|_{H^0} \leq c \|\varphi\|_{Y^{-1}(0, +\infty)}. \tag{3.35}$$

Let us show that the function $v = v(\cdot, s)$ is the unique solution of problem (2.4), and v has all desired properties.

For $s \geq 0$, set

$$B_s \triangleq \{\xi \in Y^0(s, +\infty) : \xi(\cdot, t) = 0 \text{ if } t \geq s + 1, \quad \|\xi\|_{X_2^{-1}(s, s+1)} \leq 1\}.$$

We have

$$\|v\|_{Y^1(0, +\infty)}^2 = \sup_{s=0,1,2,\dots} \sup_{\xi \in B_s} \int_s^{s+1} (v(\cdot, t), \xi(\cdot, t))_{H^0} dt.$$

Further, for $\xi \in B_s$, we have

$$\begin{aligned} \int_s^{s+1} (v(\cdot, t), \xi(\cdot, t))_{H^0} dt &= \int_s^{+\infty} (v(\cdot, t), \xi(\cdot, t))_{H^0} dt \\ &= \int_s^{+\infty} dt \int_t^{+\infty} (\varphi(\cdot, t), L_{t,r}^* \xi(\cdot, t))_{H^0} dr \\ &= \int_s^{+\infty} dr \int_s^r (\varphi(\cdot, r), L_{t,r}^* \xi(\cdot, t))_{H^0} dt = \int_s^{+\infty} (\varphi(\cdot, r), p_\xi^{(s)}(\cdot, r))_{H^0} dr, \end{aligned}$$

where

$$p_\xi^{(s)}(\cdot, r) \triangleq \int_s^r L_{t,r}^* \xi(\cdot, t) dt$$

is the solution of (3.7) with this ξ and $\rho = 0$. By Lemma 3.2, it follows that

$$\begin{aligned} \|v\|_{Y^1(0, +\infty)}^2 &= \sup_{s=0,1,2,\dots} \sup_{\xi \in B_s} \int_s^{+\infty} (\varphi(\cdot, r), p_\xi^{(s)}(\cdot, r))_{H^0} dr \\ &\leq \sup_{s=0,1,2,\dots} \sup_{\xi \in B_s} \|\varphi\|_{Y^{-1}(s, +\infty)} \|p_\xi^{(s)}\|_{Z^1(s, +\infty)} \\ &\leq c \|\varphi\|_{Y^{-1}(s, +\infty)}, \end{aligned} \tag{3.36}$$

where $c = c(\mathcal{P}_q)$ is a constant. By this estimate and (3.35), it follows that estimate (2.5) holds for v .

By (3.35), $v \in X_\infty^0(0, +\infty)$. Let us show that $v \in \mathcal{C}^0(0, +\infty)$.

Set

$$\varphi_m(x, t) = \begin{cases} \varphi(x, t) & t \leq m \\ 0 & t > m \end{cases}, \quad m = 0, 1, 2, \dots$$

Denote by $v_m(\cdot, s)$ elements of H^0 defined by (3.34) for $\varphi = \varphi_m$.

By (3.36), $v_m \in Y^1(0, +\infty)$. Further, $v_m(x, s) = 0$ for all $s \geq m$ for a.e. x . By Lemma 3.3(ii), it follows that $v_m(x, s)$ is the solution of the boundary value problem in $D \times (0, m)$

$$\begin{cases} \frac{\partial v_m}{\partial s}(x, s) + \mathcal{A}(s)v_m(x, s) + q(x, s)v_m(x, s) = -\varphi(x, s) \\ v_m(x, s)|_{x \in \partial D} = 0 \\ v_m(x, m) = 0. \end{cases} \tag{3.37}$$

Clearly, $v_m \in C^0(0, +\infty)$, since $v_m|_{D \times (0, m)} \in C^0(0, m)$, and $v_m(\cdot, s) = 0$ for $s \geq m$. For any $\rho \in H^0$ and $s \geq 0$, we have that

$$\begin{aligned} (v(s) - v_m(\cdot, s), \rho)_{H^0} &= \int_m^{+\infty} (\varphi(\cdot, t), L_{s,t}^* \rho)_{H^0} dt \\ &\leq \|\varphi\|_{Y^{-1}(0, +\infty)} \sum_{k=m}^{+\infty} \left(\int_k^{k+1} \|L_{s,t}^* \rho\|_{H^1}^2 dt \right)^{1/2} \rightarrow 0 \quad \text{as } m \rightarrow +\infty, \end{aligned}$$

since

$$\sum_{k=m}^{+\infty} \left(\int_k^{k+1} \|L_{s,t}^* \rho\|_{H^1}^2 dt \right)^{1/2} < +\infty.$$

Hence $v_m(\cdot, s) \rightarrow v(s)$ weakly in H^0 for all $s \geq 0$.

Let us show that $v_m(\cdot, s) \rightarrow v(\cdot, s)$ in H^0 uniformly in s from any finite interval. Parabolic equations in (3.37) and (3.7) are adjoint. This means that

$$v_m(x, s) = \int_s^m dt \int_D g(y, x, t, s) \varphi(y, t) dy, \quad s \leq m. \tag{3.38}$$

Here $g(x, y, t, s)$ is the Green’s function for problem (3.7) such that (3.12) holds. By semi-group properties of the solution of problem (3.7), we have that $g(\cdot, y, t, s) = L_{s+1,t}^* \rho_y(\cdot, s)$ for any $y \in D$ and $t > s + 1$, where $\rho_y(\cdot, s) \triangleq g(\cdot, y, s + 1, s)$. Similarly to (3.13), we have that $\|\rho_y(\cdot, s)\|_{L_\infty(D)} \leq c$ for all $y \in D, s > 0$, where $c = c(\mathcal{P}_q)$ is a constant.

Therefore, $\|\rho_y(\cdot, s)\|_{H^0} \leq c_*$ for all $y \in D, s \in (s_1, s_2)$, where $(s_1, s_2) \subset [0, +\infty)$ is an arbitrary finite interval, $c_* = c_*(\mathcal{P}_q, s_1, s_2)$ is a constant that does not depend on $y \in D$.

Let $\varphi \in Y^0(0, +\infty)$, and let $k = 1, 2, \dots$. By (3.38) and (3.37), we have that

$$v_{m+k}(y, s) - v_m(y, s) = \int_m^{m+k} dt \int_D \varphi(x, t) g(x, y, t, s) dx.$$

Hence

$$\begin{aligned} \|v_{m+k}(\cdot, s) - v_m(\cdot, s)\|_{H^0}^2 &= \int_D \left[\int_m^{m+k} dt \int_D \varphi(x, t) g(x, y, t, s) dx \right]^2 dy \\ &= \int_D \left[\int_m^{m+k} (\varphi(\cdot, t), L_{s+1,t}^* \rho_y(\cdot, s))_{H^0} dt \right]^2 dy \\ &\leq \int_D \left[\int_m^{m+k} \|\varphi(\cdot, t)\|_{H^0} \|L_{s+1,t}^* \rho_y(\cdot, s)\|_{H^0} dt \right]^2 dy \\ &= \int_D \left[\sum_{i=m}^{m+k} \int_i^{i+1} \|\varphi(\cdot, t)\|_{H^0} \|L_{s+1,t}^* \rho_y(\cdot, s)\|_{H^0} dt \right]^2 dy \\ &\leq \int_D \left[\sum_{i=m}^{m+k} \left\{ \int_i^{i+1} \|\varphi(\cdot, t)\|_{H^0}^2 dt \right\}^{1/2} \left\{ \int_i^{i+1} \|L_{s+1,t}^* \rho_y(\cdot, s)\|_{H^0}^2 dt \right\}^{1/2} \right]^2 dy \\ &\leq \|\varphi\|_{Y^0(0,+\infty)}^2 \int_D \left[\sum_{i=m}^{m+k} \left\{ \int_i^{i+1} \|L_{s+1,t}^* \rho_y(\cdot, s)\|_{H^0}^2 dt \right\}^{1/2} \right]^2 dy. \end{aligned}$$

By (3.10),

$$\sup_{y \in D} \|L_{s+1,t}^* \rho_y(\cdot, s)\|_{H^0} \leq C_1 \sup_{y \in D} \|\rho_y(\cdot, s)\|_{H^0} e^{-\omega_*(t-s)},$$

where $C_1 = C_1(\mathcal{P}_q) > 0$ and $\omega_* = -\ln v - \max q(x, t) > 0$. Hence

$$\begin{aligned} \|v_{m+k}(\cdot, s) - v_m(\cdot, s)\|_{H^0} &\leq \|\varphi\|_{Y^0(0,+\infty)} \sum_{i=m}^{m+k} \left(\int_i^{i+1} C_1^2 \right. \\ &\quad \left. \times \sup_{y \in D} \|\rho_y(\cdot, s)\|_{H^0}^2 e^{-2\omega_*(t-s)} dt \right)^{1/2} \rightarrow 0 \end{aligned}$$

as $m \rightarrow +\infty$ uniformly in k and $s \in [s_1, s_2]$, where $[s_1, s_2] \subset [0, +\infty)$ is an arbitrary finite interval. Hence $\{v_m|_{D \times [s_1, s_2]}\}_{m=1}^{+\infty}$ is a Cauchy sequence in $\mathcal{C}^0(s_1, s_2)$, and it converges in this space. Remind that $v_m(\cdot, s) \rightarrow v(s)$ weakly in H^0 for all $s \geq 0$. Hence $v_m|_{D \times [s_1, s_2]} \rightarrow v|_{D \times [s_1, s_2]}$ in $\mathcal{C}^0(s_1, s_2)$, and $v \in \mathcal{C}^0(0, +\infty)$ for any $\varphi \in Y^0(0, +\infty)$. The set $Y^0(0, +\infty)$ is dense in $Y^{-1}(0, +\infty)$, the space $\mathcal{C}^0(0, +\infty)$ is complete, and (3.35) holds, i.e., $\sup_{s \geq 0} \|v(\cdot, s)\|_{H^0} \leq \text{const} \|\varphi\|_{Y^{-1}(0,+\infty)}$. It follows that $v \in \mathcal{C}^0(0, +\infty)$ for any $\varphi \in Y^{-1}(0, +\infty)$.

Let us show that $v(x, s)$ satisfies (2.4) in the desired sense. For an arbitrary $\zeta(x) \in C^\infty(D)$, such that $\text{supp } \zeta \subseteq \text{int} D$, for any $\theta > t > 0$, we have

$$\begin{aligned} (\zeta, v(\cdot, \theta) - v(\cdot, t))_{H^0} &= \lim_{m \rightarrow +\infty} (\zeta, v_m(\cdot, \theta) - v_m(\cdot, t))_{H^0} \\ &= \lim_{m \rightarrow +\infty} (\zeta, \int_t^\theta (\mathcal{A}(r)v_m(x, r) + q(x, r)v_m(x, r) - \varphi_m(x, r)) dr)_{H^0} \end{aligned}$$

$$\begin{aligned}
 &= \lim_{m \rightarrow +\infty} \left\{ \int_t^\theta (A^*(r)\zeta(x) + q(x, r)\zeta(x), v_m(x, r))_{H^0} dr \right. \\
 &\quad \left. - (\zeta, \int_t^\theta \varphi_m(x, r) dr)_{H^0} \right\} \\
 &= \int_t^\theta (A^*(r)\zeta(x) + q(x, r)\zeta(x), v(x, r))_{H^0} dr \\
 &\quad - (\zeta, \int_t^\theta \varphi(x, r) dr)_{H^0}. \tag{3.39}
 \end{aligned}$$

Thus, v satisfies (2.4) in the desired sense, i.e., as a generalized solution.

Let us prove uniqueness of the solution of (2.4). Let $\tilde{v}(x, t)$ be another solution from $X_\infty^0(0, +\infty) \cap X_{2,loc}^1(0, +\infty)$. Let $\rho \in H^0$ be arbitrary, and let $p(\cdot, t) \triangleq L_{s,t}^* \rho$, where $t \geq s$. By Lemma 3.3(ii),

$$\begin{aligned}
 &(\tilde{v}(\cdot, T), p(\cdot, T))_{H^0} - (\tilde{v}(\cdot, s), p(\cdot, s))_{H^0} = - \int_s^T (\varphi(\cdot, t), p(\cdot, t))_{H^0} dt \\
 &\quad \forall s, T : 0 \leq s \leq T.
 \end{aligned}$$

Remind that $p \in Z^1(s, +\infty)$. It follows that

$$(\tilde{v}(\cdot, s), \rho)_{H^0} = \int_s^\infty (\varphi(\cdot, t), p(\cdot, t))_{H^0} dt.$$

Since ρ was arbitrary, we have that $\tilde{v}(\cdot, s) = v(\cdot, s)$ in H^0 (see (3.34)). This completes the proof of statements (i)-(ii) of Theorem 2.1.

Let us prove statement (iii). For $\varphi \in Y^0(0, +\infty)$, the solution of problem (3.37) can be presented as

$$v_m(x, s) = \mathbf{E}\xi_m(x, s), \tag{3.40}$$

where

$$\xi_m(x, s) \triangleq \int_s^{\tau_m^{x,s}} \varphi(y^{x,s}(t), t) \exp \left\{ \int_s^t q(y^{x,s}(r), r) dr \right\} dt, \quad \tau_m^{x,s} \triangleq \tau^{x,s} \wedge m.$$

The equality (3.40) is satisfied for all $s \geq 0$ for a.e. x . For $\varphi|_{D \times (0,m)} \in L_{n+1}(D \times (0, m))$, it follows from the generalized Itô's formula from Krylov (1985), §II.10. If $\varphi \in Y^0(0, +\infty)$, then the generalized Itô's formula from Dokuchaev (1994) can be used.

Let us prove (2.6) for $v(s) = v(\cdot, s)$ defined by (3.34). We have proved already that $v_m(\cdot, s) \rightarrow v(\cdot, s)$ in H^0 and, therefore, in $L_1(D)$, as $m \rightarrow +\infty$ for any given $s \geq 0$. By linearity of (2.4), it suffices to consider the case of $\varphi(x, t) \geq 0$. Then $\xi_m(x, s)$ is non-decreasing in m (in the sense of non-negativity in $L_1(D)$). Then (2.6) follows for $\varphi \in Y^0(0, +\infty)$ and for $v(x, s)$ defined by (3.34) for all $s \geq 0$ for a.e. x .

Let us prove statement (iv). Let $Q_s^* \triangleq D \times (s, s + 1/2)$, $Q_s \triangleq D \times (s, s + 1)$. By Lemma 3.3 (v), we have that

$$\begin{aligned} \|v|_{Q_s^*}\|_{W_y^{2,1}(Q_s^*)} &\leq C_1 \|v|_{Q_s^*}\|_{W_\gamma(s,s+1/2)} \\ &\leq C (\|\varphi\|_{L_\gamma(Q_s)} + \|v(\cdot, s + 1)\|_{H^0}) \leq C \sup_{s \geq 0} \|\varphi\|_{L_\gamma(Q_s)} \end{aligned}$$

for all $s \geq 0$, where $C_i = C_i(\mathcal{P}_q, \gamma) > 0$ are constants. Then statement (iv) follows.

Let us prove statement (iv). By Lemma 3.3 (vi), we have that

$$\langle\langle \mu|_{Q_s^*} \rangle\rangle_{Q_s^*}^{(1+\alpha)} \leq C (\|\varphi\|_{L_\gamma(Q_s)} + \|v(\cdot, s + 1)\|_{H^0}) \leq C \sup_{s \geq 0} \|\varphi\|_{L_\gamma(Q_s)}$$

for all $s \geq 0$, where $C = C(\mathcal{P}_q, \gamma) > 0$ is a constant. Then statement (v) follows. This completes the proof of Theorem 2.1. □

Proof of Theorem 2.2. Consider first $\varphi \in L_2(Q_0)$. Instead of (2.9), consider boundary value problem (2.4) when

$$f(x, t) \equiv f(x, t + 1), \quad \beta(x, t) \equiv \beta(x, t + 1), \quad q(x, t) \equiv q(x, t + 1), \tag{3.41}$$

and when the parabolic equation is replaced by

$$\frac{\partial v}{\partial t}(x, t) + \mathcal{A}(t)v(x, t) + \{q(x, t) + \ln |\mu|\}v(x, t) = -\widehat{\varphi}(x, t),$$

where $\widehat{\varphi}$ is such that

$$\widehat{\varphi}(x, t) \equiv \varphi(x, t)e^{-t \ln |\mu|}, \quad t \in [0, 1], \quad \widehat{\varphi}(x, t) \equiv \frac{\mu}{|\mu|} \widehat{\varphi}(x, t + 1).$$

We have that

$$v(x, s) = \mathbf{E} \int_s^{\tau^{x,s}} \widehat{\varphi}(y^{x,s}(t), t) \exp\left(\int_s^t \{q(y^{x,s}(r), r) + \ln |\mu|\} dr\right) dt,$$

and this equality holds for all $s \geq 0$ for a.e. $x \in D$. By (3.41), the probability distribution of the vector $y^{x,s}(t)$ coincides with that of vector $y^{x,s+k}(t+k)$ for all $k = 1, 2, \dots$. Then $\frac{\mu}{|\mu|} v(x, 0) = v(x, 1)$.

Set $V(x, t) \triangleq v(x, t)e^{t \ln |\mu|}$. We have that

$$\begin{aligned} &\frac{\partial V}{\partial t}(x, t) + \mathcal{A}(t)V(x, t) + q(x, t)V(x, t) \\ &= \frac{\partial v}{\partial t}(x, t)e^{t \ln |\mu|} + \ln |\mu|V(x, t) + \mathcal{A}(t)V(x, t) + q(x, t)V(x, t) \\ &= -[\mathcal{A}(t)v(x, t) + \{q(x, t) + \ln |\mu|\}v(x, t) + \widehat{\varphi}(x, t)]e^{t \ln |\mu|} \\ &\quad + \ln |\mu|V(x, t) + \mathcal{A}(t)V(x, t) + q(x, t)V(x, t) = \varphi(x, t). \end{aligned}$$

Clearly, $V(x, 0) \equiv v(x, 0)$, and $V(x, 1) \equiv |\mu|v(x, 1)$. Hence $\mu V(x, 0) \equiv V(x, 1)$, and V is the solution of (2.9). Inequality (2.10) is satisfied with a constant c defined by the estimate for v from Theorem 2.1 (i), and this c does not depend on $\varphi \in L_2(Q_0)$.

Therefore, statements (i)-(ii) are proved for all $\varphi \in L_2(Q_0)$.

Let $\varphi \in X_2^{-1}(0, 1)$. Clearly, $L_2(Q_0)$ is dense $X_2^{-1}(0, 1)$, and there exists a sequence $\{\varphi_i\}_{i=1}^{+\infty} \subset L_2(Q_0)$ such that $\varphi_i \rightarrow \varphi$ in $X_2^{-1}(0, 1)$ as $i \rightarrow \infty$. Let V_i be the solution of problem (2.9) with $\varphi = \varphi_i$. By statement (i) that is proved already for $\varphi_i \in L_2(Q_0)$, the sequence $\{V_i\}_{i=1}^{+\infty}$ is a Cauchy sequence in $X_2^1(0, 1)$ and in $C^0(0, 1)$. Hence this sequence has the limit $V \in X_2^1(0, 1) \cap C^0(0, 1)$. It is easy to see that this V is a solution of problem (2.9). Uniqueness of V follows from (2.10). Therefore, statements (i)-(ii) hold for all $\varphi \in X_2^{-1}(0, 1)$.

Statement (iii) follows from Theorem 2.1(v) applied for v . This completes the proof of Theorem 2.2. □

Proof of Theorem 2.3. Let e_1 and e_2 be the indicator functions of the random events $\{\tau_1 \geq \tau_2\}$ and $\{\tau_2 > \tau_1\}$ respectively.

Let \mathcal{F}_t be the filtration generated by $w(t)$ and a .

The random variables e_i are measurable with respect to the σ -algebras $\mathcal{F}_{\tilde{\tau}}, \mathcal{F}_{\tau_i}$, $i = 1, 2$, associated with the Markov times (with respect to the filtration \mathcal{F}_t) $\tilde{\tau}$ and τ_i (see, e.g., Gihman and Skorohod (1975), Chapter 4, §2).

Set

$$\zeta_i(t) \triangleq v_i(y_i(t), t), \quad \xi_i(t) \triangleq \zeta_i(t)e^{\lambda(t-\tilde{\tau})}, \quad t \in [\tilde{\tau}, \tau_i].$$

Clearly, $1 \in L_\gamma(D)$ for all $\gamma > 1$. By Theorem 2.1 (iv)-(v), it follows that $v_i(x, t)$ and $\frac{\partial v_i}{\partial x_k}(x, t)$ are continuous and bounded, and the norms $\|\frac{\partial v_i}{\partial t}\|_{L_\gamma(Q_s)}$, $\|\frac{\partial^2 v_i}{\partial x_k \partial x_m}\|_{L_\gamma(Q_s)}$ are bounded in $s \geq 0$ for any $\gamma > 1$, where $Q_s \triangleq D \times (s, s + 1)$, $k, m = 1, \dots, n$. Therefore, we can apply to $\zeta_i(t)$ the generalized Itô's formula given by Theorem II.10.1 from Krylov (1980), p. 122. By this Itô's formula and (2.13), we obtain

$$\begin{aligned} d\zeta_i(t) &= \left[\frac{\partial v_i}{\partial t}(y_i(t), t) + \mathcal{A}_i(t)v_i(y_i(t), t) \right] dt + \frac{\partial v_i}{\partial x}(y_i(t), t)\beta_i(y_i(t), t)dw(t) \\ &= -[\lambda v_i(y_i(t), t) + 1]dt + \frac{\partial v_i}{\partial x}(y_i(t), t)\beta_i(y_i(t), t)dw(t) \\ &= -[\lambda \zeta_i(t) + 1]dt + \frac{\partial v_i}{\partial x}(y_i(t), t)\beta_i(y_i(t), t)dw(t), \end{aligned}$$

and

$$\begin{aligned} d\xi_i(t) &= e^{\lambda(t-\tilde{\tau})}d\zeta_i(t) + \lambda e^{\lambda(t-\tilde{\tau})}\zeta_i(t)dt \\ &= e^{\lambda[t-\tilde{\tau}]} \frac{\partial v_i}{\partial x}(y_i(t), t)\beta_i(y_i(t), t)dw(t) - e^{\lambda(t-\tilde{\tau})}dt. \end{aligned}$$

Hence

$$\begin{aligned}
 \mathbf{E}\{e_1\{v_1[y_1(\tilde{\tau}), \tilde{\tau}] - v_1[y_2(\tilde{\tau}), \tilde{\tau}]\}\} &= \mathbf{E}\{e_1\{v_1[y_1(\tau_2), \tau_2] - v_1[y_2(\tau_2), \tau_2]\}\} \\
 &= -\mathbf{E}\{e_1\{v_1[y_1(\tau_1), \tau_1] - v_1[y_1(\tau_2), \tau_2]\}\} \\
 &= -\mathbf{E}\{e_1\{\xi_1(\tau_1) - \xi_1(\tilde{\tau})\}\} \\
 &= \mathbf{E}\left\{e_1 \int_{\tilde{\tau}}^{\tau_1} e^{\lambda(t-\tilde{\tau})} dt\right\} \\
 &= \frac{1}{\lambda} \mathbf{E}\{e_1\{e^{\lambda(\tau_1-\tilde{\tau})} - 1\}\} \\
 &= \frac{1}{\lambda} \mathbf{E}\{e_1\{e^{\lambda(\tau_1-\tau_2)} - 1\}\}. \tag{3.42}
 \end{aligned}$$

Hence

$$\frac{1}{\lambda} \mathbf{E}\{e_1\{e^{\lambda(\tau_1-\tau_2)} - 1\}\} \leq \sup_{(x,t) \in Q} \left| \frac{dv_1}{dx}(x,t) \right| \mathbf{E}\{e_1|y_1(\tilde{\tau}) - y_2(\tilde{\tau})|\}. \tag{3.43}$$

If we replaced the indices 1, 2 in (3.42) by 2,1, we get similarly that

$$\frac{1}{\lambda} \mathbf{E}\{e_2\{e^{\lambda(\tau_2-\tau_1)} - 1\}\} \leq \sup_{(x,t) \in Q} \left| \frac{dv_2}{dx}(x,t) \right| \mathbf{E}\{e_2|y_1(\tilde{\tau}) - y_2(\tilde{\tau})|\}. \tag{3.44}$$

Clearly,

$$\mathbf{E}[e^{\lambda|\tau_1-\tau_2|} - 1] = \mathbf{E}\{e_1\{e^{\lambda(\tau_1-\tau_2)} - 1\}\} + \mathbf{E}\{e_2\{e^{\lambda(\tau_2-\tau_1)} - 1\}\}. \tag{3.45}$$

Now (2.14) follows from (3.42)–(3.45). □

Remark 3.2. In fact, the condition in (2.2) that $\partial f/\partial x$ is locally bounded can be lifted. Without this condition, equation (2.1) has an unique weak solution for any given (s, a) . More precisely, there exists a set $(\Omega, \mathcal{F}, \mathbf{P}, w(\cdot), y^{a,s}(\cdot))$ such that equation (2.1) holds and $w(\cdot)$ does not depend on a ; the distribution of $y^{a,s}(\cdot)$ is uniquely defined (see, e.g., Chapter II from Krylov (1980), Section 3 of Chapter 3 from Gihman and Skorohod (1975), and Theorems 4.1 and 4.3-4.4 from Dokuchaev (1997)). In this case, the formulations of the results need to be adjusted as the following. Lemma 2.1 holds for any $y^{a,s}(t)$ such as described here. Theorem 2.1 (iii) holds for $y^{x,s}(t)$ defined in the conditional probability space as $y^{a,s}(t)$ given $a = x$, where a is such that it has the probability density function in H^0 . Theorem 2.1 (i)-(ii), (iv)-(v) and Theorem 2.3 hold in their present form. Remind that Theorem 2.3 requires that (2.11) is satisfied for $y_i(t)$ with the same $w(\cdot)$ for $i = 1, 2$.

The author wishes to thank R. McEachin, A. Rodkina, T. Salisbury, and W. Zhang for useful discussion. As it was mentioned earlier, T. Salisbury suggested to the author the way how to simplify the proof of Lemma 2.1. The author also wishes to thank the anonymous referee for their insightful comments which greatly strengthened the paper.

References

- Adams, R.A.: Sobolev Spaces. Academic press, 1975
- Aronson, D.J.: Bounds for the fundamental solution of a parabolic equation. *Bull. of Amer. Math. Soc.* **73** (6), 890–896 (1967)
- Aronson, D.J.: Non-negative solutions of linear parabolic equations. *Ann. Scu. Norm. Sup. di Pisa.* **22** (4), 607–694 (1968)
- Bokalo, N.M.: Unique solvability of boundary value problems for semi-linear parabolic equations in unbounded domains without conditions at infinity. *Siberian Mathematical Journal* **34** (4), 33–40 (1993)
- Dokuchaev, N.G.: On moment's of first exit for homogeneous processes of diffusion type. *Theory of Probability and its Applications* **31** (3), 497–498 (1987)
- Dokuchaev, N.G.: The integral estimations for ordinary differential equations with a discontinuity on a domain boundary. *Diff. Eqs.* **28** (11), 1540–1549 (1992)
- Dokuchaev, N.G.: Distribution of Ito processes: estimates for density functions and for conditional expectations of integral functionals. *Theory of Probability and Its Applications* **39** (4), 662–670 (1994)
- Dokuchaev, N.G.: Cordes conditions and some alternatives for parabolic equations and discontinuous diffusion. *Diff. Eqs.* **33** (4), 433–442 (1997)
- Gihman, I.I., Skorohod, A.V.: *The Theory of Stochastic Processes. Vol. 2.* New York: Springer-Verlag, 1975
- Krylov, N.V.: *Controlled Diffusion Processes.* Berlin-Heidelberg, New York: Springer-Verlag, 1980
- Ladyzenskaia, O.A., Solonnikov, V.A., Ural'ceva, N.N.: *Linear and Quasi-Linear Equations of Parabolic Type.* Providence, RI: American Mathematical Society, 1968
- Ladyzhenskaia, O.A.: *The Boundary Value Problems of Mathematical Physics.* New York: Springer-Verlag, 1985
- Nakao, M.: Periodic solution of some nonlinear degenerate parabolic equations. *J. Math. Anal. Appl.* **104** (2), 554–557 (1984)
- Oleinik, O.A., Yosifian, G.A.: The analogue of Saint-Venant's principle and the uniqueness of solutions to boundary value problems in unbounded domains for parabolic equations. *Uspekhi Matematicheskikh Nauk* **31** (6), 143–166 (1976)
- Revuz, D., Yor, M.: *Continuous Martingales and Brownian Motion.* New York: Springer-Verlag, 1999
- Shelukhin, V.V.: Variation principle for non-local in time problems for linear evolutionary equations. *Siberian Math. J.* **34** (2), 191–207 (1993)
- Yosida, K.: *Functional Analysis.* Berlin-Gottingen-Heidelberg: Springer-Verlag, 1965