

Richard F. Bass\* · Dahae You

# A Fatou theorem for $\alpha$ -harmonic functions in Lipschitz domains

Received: 16 July 2004 / Revised version: 5 January 2005 /  
Published online: 15 March 2005 – © Springer-Verlag 2005

**Abstract.** We study  $\alpha$ -harmonic functions in Lipschitz domains. We prove a Fatou theorem when the boundary function is bounded and  $L^p$ -Hölder continuous of order  $\beta$  with  $\beta p > 1$ .

---

## 1. Introduction

Let  $D$  be a bounded Lipschitz domain in  $\mathbb{R}^d$  and let  $\alpha \in (0, 2)$ . Our purpose in this paper is to establish a Fatou theorem for bounded functions which are  $\alpha$ -harmonic in  $D$ . An  $\alpha$ -harmonic function is one which is harmonic with respect to a symmetric stable process of index  $\alpha$ , or equivalently, with respect to the operator  $-(-\Delta)^{\alpha/2}$ . There has been a great deal of recent work on the properties of  $\alpha$ -harmonic functions; see [5]–[15], [18], [20], and [23].

To be more precise, let  $f$  be a bounded function on  $D^c$ , let  $(X_t, \mathbb{P}^x)$  be a symmetric stable process of index  $\alpha \in (0, 2)$ , let  $\tau_D = \inf\{t : X_t \notin D\}$ , and let

$$u_f(x) = \mathbb{E}^x f(X_{\tau_D}). \quad (1.1)$$

Then  $u_f$  is what is known as a regular  $\alpha$ -harmonic function. It is not the case that  $u_f$  must have nontangential limits a.e. with respect to surface measure on the boundary of  $D$ . This was shown in Section 5 of [3] in the case of a half space; the example can be easily modified to hold in bounded Lipschitz domains. Therefore we assume in addition that  $f$  is  $L^p$ -Hölder continuous. That is, we let  $\Lambda_\beta^{p,\infty}(\mathbb{R}^d)$  be the collection of functions in  $L^p(\mathbb{R}^d)$  such that the norm

$$\|f\|_p + \sup_{|t|>0} \frac{\|f(\cdot + t) - f(\cdot)\|_p}{|t|^\beta}$$

is finite (the notation agrees with that of [21]). Let  $B(x, \zeta)$  be the open ball with radius  $\zeta$  and center  $x$ . We let  $\Lambda_{loc}^b(\beta, p, D^c)$  be the collection of bounded functions

---

R. F. Bass, D. You: Department of Mathematics, University of Connecticut, Storrs, CT 06269, USA e-mail: bass@math.uconn.edu; you@math.uconn.edu

*Mathematics Subject Classifications (2000):* Primary 31B25; Secondary 60J50

*Key words or phrases:* Stable processes – Fatou theorem –  $\alpha$ -harmonic – Nontangential convergence – Maximal function – Hitting probabilities

\*Research supported by NSF Grant DMS0244737

$f$  such that for each  $x \in \partial D$  there exists  $\zeta$  depending on  $x$  such that  $f$  agrees on  $D^c \cap B(x, \zeta)$  with a bounded function in  $\Lambda_\beta^{p, \infty}(\mathbb{R}^d)$ .

Next we explain what nontangential convergence means. Define the truncated cone  $C_{a, \theta} = \{(\tilde{z}, z_d) : 0 < z_d < a, |\tilde{z}| < \theta z_d\}$ , where  $\tilde{z} \in \mathbb{R}^{d-1}$ . Choose  $a < a'$  and  $\theta < \theta'$  and for each  $x \in \partial D$  a rotation  $R_x$  such that  $x + R_x(C_{a', \theta'}) \subset D$ . Then let  $C(x) = x + R_x(C_{a, \theta})$ . We need to choose  $R_x$  in a measurable way, but since each rotation is associated with a  $d \times d$  orthogonal matrix, this is routine.

We then say that  $u_f$  converges nontangentially at  $x \in \partial D$  if

$$\lim_{z \in C(x), z \rightarrow x} u_f(x)$$

exists.

Our main theorem is

**Theorem 1.1.** *Suppose  $D$  is a bounded Lipschitz domain,  $\beta \in (0, 1)$ ,  $p \in (1, \infty]$ , and  $\beta p > 1$ . Suppose  $f$  is a function in  $\Lambda_{loc}^b(\beta, p, D^c)$ . Then  $u_f$  defined by (1.1) converges nontangentially at all points of  $\partial D$  except for a set of zero surface measure.*

It is routine to extend this result to unbounded Lipschitz domains. The example in Section 5 of [3] shows that our condition  $\beta p > 1$  is sharp. We discuss the identification of the limit in the remark at the end of the paper.

Theorem 1.1 is in sharp contrast to the Fatou theorem for functions that are harmonic in  $D$  (with respect to the Laplacian). The Fatou theorem for harmonic functions says that if  $D$  is a Lipschitz domain and  $f \in L^p(\partial D)$ , then  $u_f$  converges nontangentially a.e. with respect to surface measure on  $\partial D$ . See [1], Section III.4, for details and some applications. While most results that are true for harmonic functions are also true for  $\alpha$ -harmonic functions, often with weaker assumptions, the Fatou theorem is an exception in that extra regularity is needed.

Our method is to first obtain an estimate on the Poisson kernel. Unlike the half space case, our estimate is not at all sharp, and for  $\alpha \in [1, 2)$  is not even locally integrable. We therefore have to derive some hitting probability estimates for symmetric stable processes in Lipschitz domains. Nevertheless, the theorem we obtain is as sharp as the one we obtained in [3] in the half space case, where an explicit formula for the Poisson kernel was available. We point out that our results complement those of [18]: see the remark following the proof of Proposition 3.5.

The paper is organized as follows. In Section 2 we compute an upper bound of the Poisson kernel for Lipschitz domains. Section 3 contains some estimates for  $\alpha$ -harmonic functions in Lipschitz domains. In Section 4 we introduce a maximal function similar to the one in [3] and establish an estimate for it, while in Section 5 we prove Theorem 1.1.

We use the letter  $c$ , with or without subscripts, to denote positive finite constants whose exact value is unimportant. Let  $B(x_0, r) = \{x \in \mathbb{R}^d : |x - x_0| < r\}$  be the open ball centered at  $x_0$  with radius  $r$ . Given a Borel subset  $D$  of  $\mathbb{R}^d$ , let  $|D|$  denote the Lebesgue measure of  $D$  and let  $\delta_D(x)$  be the distance between  $x$  and  $\partial D$ , where  $\partial D$  is the boundary of  $D$ . We sometimes write points  $z = (z_1, \dots, z_d) \in \mathbb{R}^d$  as  $z = (\tilde{z}, z_d)$  with  $\tilde{z} \in \mathbb{R}^{d-1}$ . Given a Borel set  $A$ , we use  $A - x = \{z - x : z \in A\}$ ,

$aA = \{az : z \in A\}$ , and  $\Psi(A) = \{\Psi(z) : z \in A\}$  when  $\Psi$  is a map from  $\mathbb{R}^d$  to  $\mathbb{R}^d$ . The paths of  $X_t$  are right continuous with left limits; we use  $X_{t-} = \lim_{s \uparrow t, s < t} X_s$  and  $\Delta X_t = X_t - X_{t-}$ . If  $A$  is a Borel set, we define

$$\tau_A = \inf\{t : X_t \notin A\}, \quad T_A = \inf\{t : X_t \in A\}$$

for the first exit and first entrance of  $A$ , respectively. A Lipschitz function  $\Gamma$  with Lipschitz constant  $\lambda$  is a map  $\Gamma : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$  satisfying

$$|\Gamma(\tilde{x}) - \Gamma(\tilde{y})| \leq \lambda|\tilde{x} - \tilde{y}|, \quad \tilde{x}, \tilde{y} \in \mathbb{R}^{d-1}.$$

A Lipschitz domain is one where for each  $x \in \partial D$  there is  $r_x > 0$  and a coordinate system depending on  $x$  such that  $D \cap B(x, r_x)$  agrees with the intersection of  $B(x, r_x)$  with the region above the graph of a Lipschitz function.

### 2. Poisson kernel

The Green function  $G_D$  for  $X_t$  is

$$G_D(x, y) = c(\alpha, d) \left[ \frac{1}{|x - y|^{d-\alpha}} - \int_{D^c} \frac{1}{|y - z|^{d-\alpha}} \omega^x(dz, D) \right]$$

where  $\omega^x(\cdot, D)$  is the  $\alpha$ -harmonic measure on  $D^c$  given by

$$\omega^x(A, D) = \mathbb{P}^x(X_{\tau_D} \in A), \quad A \subset D^c.$$

It is known (see [13], [17]) that the distribution of  $X_{\tau_D}$  under  $\mathbb{P}^x$  has a density with respect to  $d$ -dimensional Lebesgue measure on  $D^c$  given by

$$P_D(x, z) = c(\alpha, d) \int_D \frac{G_D(x, y)}{|y - z|^{d+\alpha}} dy, \quad x \in D, \quad z \in \overline{D}^c, \quad (2.1)$$

when  $D$  is a domain satisfying a uniform exterior cone condition. In particular, in this case  $\mathbb{P}^x(X_{\tau_D} \in \partial D) = 0$ .  $P_D$  is called the Poisson kernel for  $D$ .

In the case of a ball,

$$P_{B(0,r)}(x, y) = c_1 \left[ \frac{r^2 - |x|^2}{|y|^2 - r^2} \right]^{\alpha/2} |x - y|^{-d}; \quad (2.2)$$

see [19], pp. 121–122. We deduce from this that

$$\mathbb{P}^x(|X_{\tau_{B(x,r)}} - x| > K) \leq c_2(r/K)^\alpha. \quad (2.3)$$

Another consequence of (2.2) is that  $u_f$  is  $C^\infty$  in  $D$  if  $f$  is bounded, although we do not use this fact.

The following proposition gives an upper bound for the Poisson kernel in a Lipschitz domain.

**Proposition 2.1.** *Let  $d \geq 2$ . Let  $D$  be either a bounded Lipschitz domain in  $\mathbb{R}^d$  or else the region above the graph of a Lipschitz function with Lipschitz constant  $\lambda$ . There exists a constant  $c_1$  depending only on  $\alpha, \lambda$ , and  $d$  such that*

$$P_D(x, z) \leq c_1 \frac{1}{|x - z|^{d-\alpha} \delta_D(z)^\alpha}, \quad x \in D, z \in \overline{D}^c.$$

*Proof.* Let  $r = |x - z|$ . We write  $D = A_1 \cup A_2 \cup A_3$ , where

$$\begin{aligned} A_1 &= D \cap B(x, r/2), \\ A_2 &= D \cap [B(x, 6r) \setminus B(x, r/2)], \\ A_3 &= D \cap B(x, 6r)^c. \end{aligned}$$

Then by (2.1)

$$P_D(x, z) \leq c_2 \int_D \frac{1}{|x - y|^{d-\alpha}} \frac{1}{|y - z|^{d+\alpha}} dy = c_2 \left[ \int_{A_1} + \int_{A_2} + \int_{A_3} \right].$$

For  $y \in A_1$  we have  $|y - z| \geq r/2$ , and so the integral over  $A_1$  is bounded by

$$c_3 r^{-d-\alpha} \int_{B(x, r/2)} \frac{1}{|x - y|^{d-\alpha}} dy = c_4 r^{-d}. \tag{2.4}$$

For  $y \in A_2$  we have  $|x - y| \geq r/2$  and  $|y - z| \geq \delta_D(z)$ , and so the integral over  $A_2$  is bounded by

$$r^{-d+\alpha} \int_{\{|y-z| \geq \delta_D(z)\}} \frac{1}{|y - z|^{d+\alpha}} dy \leq c_5 r^{-d+\alpha} \delta_D(z)^{-\alpha}. \tag{2.5}$$

For  $y \in A_3$  we have  $|y - z| \geq |x - y|/2$ , and so the integral over  $A_3$  is bounded by

$$c_6 \int_{B(x, 6r)^c} \frac{1}{|x - y|^{2d}} dy \leq c_7 r^{-d}. \tag{2.6}$$

If we combine (2.4)–(2.6) and note that  $r \geq \delta_D(z)$ , we obtain our result. □

### 3. Estimates

Our estimate for the Poisson kernel given in the previous section is satisfactory when  $\alpha \in (0, 1)$  because the kernel is locally integrable. However, when  $\alpha \in [1, 2)$ , more information is needed.

Let us suppose throughout this section that  $\Gamma$  is a bounded Lipschitz function with Lipschitz constant  $\lambda$  and that  $D$  is the region above  $\Gamma$ , that is,  $D = \{z = (\tilde{z}, z_d) : z_d > \Gamma(\tilde{z})\}$ . Define for  $z \in \mathbb{R}^d$

$$v(z) = z_d - \Gamma(\tilde{z}).$$

Note  $v(z)$  and  $\delta_D(z)$  are comparable. Let

$$W(b) = \{z : 0 < v(z) < b\}.$$

Let  $\rho \in (0, \frac{1}{4})$  be chosen later, let

$$W_i = W(\rho^i), \quad \tilde{W}_i = W(2\rho^i),$$

and let

$$T_i = T_{W_i}, \quad \tilde{T}_i = T_{\tilde{W}_i},$$

where  $T_A = \inf\{t : X_t \in A\}$  for any Borel set  $A$ .

Because we are in the region above a Lipschitz function, the exterior cone condition holds. If  $z \in D$  and  $\gamma < 1$ , the distribution of  $\mathbb{P}^z(X_{T_{B(z, \gamma \delta_D(z))^c}} \in dy)$  is given by (2.2). Because of the exterior cone condition, we see that there exists  $a \in (0, 1)$ , depending on  $\gamma$ , such that

$$\mathbb{P}^z(X_{T_{B(z, \gamma \delta_D(z))^c}} \in D^c) \geq a. \tag{3.1}$$

**Lemma 3.1.** *There exists a constant  $c_1$  not depending on  $\rho$  such that if  $i > j$  and  $z \in D \setminus W_j$ , then*

$$\mathbb{P}^z(X_{T_j} \in \tilde{W}_i) \leq c_1 \rho^{i-j}.$$

*Proof.* Set  $n(x, h) = |h|^{-d-\alpha}$ ; although  $n$  does not depend on  $x$ , we include  $x$  in the notation to conform to [2], which we will use later on. We begin by claiming that if  $x \in D \setminus W_j$ , then

$$\int_{\tilde{W}_i-x} n(x, h) dh \leq c_2 \rho^i v(x)^{-1-\alpha}. \tag{3.2}$$

To see this, note that the distance from  $x$  to  $\tilde{W}_i$  is at least  $c_3(v(x) - 2\rho^j) \geq c_4 v(x)$  since  $x \notin W_j$  and  $j < i$ . So

$$\int_{(\tilde{W}_i-x) \cap B(x, v(x))} n(x, h) dh \leq c_5 \frac{\rho^i v(x)^{d-1}}{v(x)^{d+\alpha}} \tag{3.3}$$

and for  $k \geq 0$

$$\int_{(\tilde{W}_i-x) \cap (B(x, 2^{k+1}v(x)) \setminus B(x, 2^k v(x)))} n(x, h) dh \leq c_6 \frac{\rho^i (2^{k+1}v(x))^{d-1}}{(2^k v(x))^{d+\alpha}}. \tag{3.4}$$

Summing (3.4) over  $k$  from 0 to  $\infty$  and adding (3.3) yields (3.2).

Because  $D$  satisfies a uniform exterior cone condition, there exists a cone  $V_x$  with vertex at  $x$  and with axis in the  $(0, \dots, 0, -1)$  direction and a constant  $c_7$  such that  $B(x, c_7 \delta_D(x))^c \cap V_x \subset D^c$ ; we may take the aperture of  $V_x$  to be independent of  $x$ . Then

$$\int_{D^c-x} n(x, h) dh \geq c_8 \int_{B(x, c_7 \delta_D(x))^c \cap V_x} \frac{1}{|h|^{d+\alpha}} dh \geq c_9 v(x)^{-\alpha}. \tag{3.5}$$

Combining (3.2) and (3.5) and using the fact that  $v(x) \geq \rho^j$  we obtain

$$\begin{aligned} \int_{\tilde{W}_i-x} n(x, h) dh &\leq c_2 \rho^i v(x)^{-1-\alpha} \\ &\leq c_{10} \frac{\rho^i}{v(x)} \int_{D^c-x} n(x, h) dh \\ &\leq c_{10} \rho^{i-j} \int_{D^c-x} n(x, h) dh. \end{aligned} \tag{3.6}$$

If there is a jump from  $D \setminus W_j$  to  $\tilde{W}_i$  before or at time  $T_j \wedge \tau_D$ , then

$$\sum_{t \leq T_j \wedge \tilde{T}_i \wedge \tau_D} 1_{(\Delta X_t \neq 0)} 1_{(X_t \in \tilde{W}_i)} \geq 1.$$

Therefore

$$\mathbb{P}^z(X_{T_j} \in \tilde{W}_i) \leq \mathbb{E}^z \sum_{t \leq T_j \wedge \tilde{T}_i \wedge \tau_D} 1_{(\Delta X_t \neq 0)} 1_{(X_t \in \tilde{W}_i)}. \tag{3.7}$$

Using the Lévy system formula (see [2], Proposition 2.3) and (3.6), if  $t_0 > 0$  we have

$$\begin{aligned} & \mathbb{E}^z \sum_{t \leq T_j \wedge \tilde{T}_i \wedge \tau_D \wedge t_0} 1_{(\Delta X_t \neq 0)} 1_{(X_t \in \tilde{W}_i)} \\ &= \mathbb{E}^z \int_0^{T_j \wedge \tilde{T}_i \wedge \tau_D \wedge t_0} \int_{\tilde{W}_i - X_s} n(X_s, h) dh ds \\ &\leq c_{10} \rho^{i-j} \mathbb{E}^z \int_0^{T_j \wedge \tilde{T}_i \wedge \tau_D \wedge t_0} \int_{D^c - X_s} n(X_s, h) dh ds \\ &= c_{10} \rho^{i-j} \mathbb{E}^z \sum_{t \leq T_j \wedge \tilde{T}_i \wedge \tau_D \wedge t_0} 1_{(\Delta X_t \neq 0)} 1_{(X_t \in D^c)} \\ &\leq c_{10} \rho^{i-j}. \end{aligned}$$

The last inequality follows because  $\sum_{t \leq \tau_D} 1_{(\Delta X_t \neq 0)} 1_{(X_t \in D^c)}$  is at most 1. If we now let  $t_0 \uparrow \infty$ , use monotone convergence, and combine with (3.7), we obtain our result.  $\square$

We next prove

**Proposition 3.2.** *There exist  $\rho \in (0, \frac{1}{4})$  and  $\sigma \in (\frac{1}{2}, 1)$  such that if  $v(x_0) \geq 2$ , then*

$$\mathbb{P}^{x_0}(T_i < \tau_D) \leq 2\sigma^i. \tag{3.8}$$

*Proof.* Because we are in a Lipschitz domain with Lipschitz constant  $\lambda$ , there exists  $\gamma < 1$  independent of  $i$  such that if  $z \in W_i \setminus \tilde{W}_{i+1}$ , then  $B(z, \gamma \delta_D(z)) \subset W_i \setminus W_{i+1}$ . Let  $a$  be chosen as in (3.1). We will choose suitable  $\rho \in (0, \frac{1}{4})$  and  $\sigma \in (\frac{1}{2}, 1)$  later. Let  $W_0 = D \setminus W_1$ . We want to show that (3.8) holds. We will prove this by induction. Since  $\sigma > \frac{1}{2}$ , the case  $i = 1$  is obvious. We now suppose that (3.8) holds for all  $j \leq i$  and we will prove it holds for  $i + 1$ .

If  $T_{i+1} < \tau_D$ , then either

- (a) the process  $X_t$  hits  $W_i \setminus \tilde{W}_{i+1}$  and then hits  $W_{i+1}$ , all before time  $\tau_D$ , or else
- (b) there is a largest  $j < i$  such that  $X_t$  hits  $W_j$  but only hits  $W_{j+1}$  when it jumps into  $\tilde{W}_{i+1}$ .

We first estimate the probability of (a). We need to bound

$$\mathbb{P}^{x_0}(T_i < \tau_D, X_{T_i} \in W_i \setminus \tilde{W}_{i+1}, T_{i+1} < \tau_D).$$

Using the strong Markov property at time  $T_i$ , this is bounded by

$$\mathbb{E}^{x_0} [\mathbb{P}^{X_{T_i}} (T_{i+1} < \tau_D); X_{T_i} \in W_i \setminus \tilde{W}_{i+1}, T_i < \tau_D]. \tag{3.9}$$

Using (3.1) we conclude that

$$\mathbb{P}^z (T_{i+1} < \tau_D) \leq \mathbb{P}^z (X_{T_{B(z, \gamma \delta_D(z))}^c} \in D) \leq 1 - a.$$

Therefore, using the induction hypothesis, (3.9) is bounded by

$$\begin{aligned} (1 - a) \mathbb{P}^{x_0} (X_{T_i} \in W_i \setminus \tilde{W}_{i+1}, T_i < \tau_D) \\ \leq (1 - a) \mathbb{P}^{x_0} (T_i < \tau_D) \\ \leq 2(1 - a) \sigma^i. \end{aligned}$$

So the probability of (a) is bounded by

$$2(1 - a) \sigma^i. \tag{3.10}$$

Next we bound the probability of the event in (b). When  $j = 0$  we need to estimate

$$\mathbb{P}^{x_0} (X_{T_1} \in \tilde{W}_{i+1}),$$

and by Lemma 3.1 this is bounded by

$$c_2 \rho^i. \tag{3.11}$$

Now suppose  $1 \leq j < i$ . We need to bound

$$\mathbb{P}^{x_0} (T_j < \tau_D, X_{T_j} \in W_j \setminus W_{j+1}, X_{T_{j+1}} \in \tilde{W}_{i+1}).$$

By the strong Markov property, Lemma 3.1, and the induction hypothesis, this is bounded by

$$\begin{aligned} \mathbb{E}^{x_0} [\mathbb{P}^{X_{T_j}} (X_{T_{j+1}} \in \tilde{W}_{i+1}); T_j < \tau_D, X_{T_j} \in W_j \setminus W_{j+1}] \\ \leq c_3 \rho^{i-j} \mathbb{P}^{x_0} (T_j < \tau_D) \\ \leq 2c_3 \rho^{i-j} \sigma^j. \end{aligned}$$

So if  $1 \leq j < i$ , the probability of (b) is bounded by

$$2c_3 \rho^{i-j} \sigma^j. \tag{3.12}$$

If we add the estimates (3.10), (3.11), and (3.12), we have

$$\begin{aligned} \mathbb{P}^{x_0} (T_{i+1} < \tau_D) &\leq c_2 \rho^i + 2 \sum_{j=1}^{i-1} c_3 \rho^{i-j} \sigma^j + 2(1 - a) \sigma^i \\ &= c_2 \rho^i + 2c_3 \rho^i \frac{\sigma}{\rho} \frac{(\sigma/\rho)^{i-1} - 1}{(\sigma/\rho) - 1} + 2(1 - a) \sigma^i. \end{aligned} \tag{3.13}$$

Since  $\sigma/\rho > 2$ , this is bounded by

$$c_2\rho^i + c_4\rho^i\left(\frac{\sigma}{\rho}\right)^{i-1} + 2(1-a)\sigma^i \leq 2\sigma^i(c_2(\rho/\sigma)^i + c_5(\rho/\sigma) + (1-a)).$$

If we choose  $\rho$  small enough so that  $(c_2 + c_5)(\rho/\sigma) \leq a/2$  and  $\sigma \in (1 - a/2, 1)$ , then the left hand side of (3.13) is bounded by

$$2\sigma^i(1 - a/2) \leq 2\sigma^{i+1}.$$

This proves the induction hypothesis, and hence the proposition. □

The main estimate we need is the following. Let

$$S_\varepsilon = \{z : -\varepsilon < v(z) < 0\}.$$

**Proposition 3.3.** *There exist  $c_1, b > 0$  such that if  $v(x_0) \in [2, 3]$  and  $\varepsilon > 0$ , then*

$$\mathbb{P}^{x_0}(X_{\tau_D} \in S_\varepsilon) \leq c_1\varepsilon^b.$$

*Proof.* Let  $\rho, \sigma$  be as in Proposition 3.2. Given  $\varepsilon$  there exists  $i$  such that  $\rho^{i+1} \leq \varepsilon < \rho^i$ . Let  $D^* = \{z : -\rho^i < v(z)\}$ . If  $X_{\tau_D} \in S_\varepsilon$ , then  $T_{S_\varepsilon} < \tau_{D^*}$ . By the above proposition applied to  $D^*$ ,

$$\mathbb{P}^{x_0}(T_{S_\varepsilon} < \tau_{D^*}) \leq 2\sigma^i.$$

If we choose  $b = \log \sigma / \log \rho$  and  $c_1 = 2\rho^{-b}$ , then

$$2\sigma^i = \frac{2}{\rho^b}(\rho^{i+1})^b \leq c_1\varepsilon^b,$$

which completes the proof. □

Since for  $r > 0$  the process  $r^{-\alpha} X_{rt}$  has the same law under  $\mathbb{P}^0$  as  $X_t$  and  $rD$  is again the region above the graph of a Lipschitz function with Lipschitz constant  $\lambda$ , a scaling argument yields

**Corollary 3.4.** *There exist  $c_1, b > 0$  such that if  $x_0 \in D$ , then*

$$\mathbb{P}^{x_0}(X_{\tau_D} \in S_{\varepsilon v(x_0)}) \leq c_1\varepsilon^b. \tag{3.14}$$

The constant in Corollary 3.4 is not necessarily the same as the one in Proposition 3.3.

Although dominated convergence and the Harnack inequality give us something like (3.14) for  $x_0$  such that  $v(x_0) \in [2, 3]$ , the uniformity over  $v(x_0)$  requires the extra work above.

In the following, again the uniformity is where the work comes in.

**Proposition 3.5.** *Let  $\varepsilon > 0$ . There exists  $\eta$  such that if  $v(x_0) < \eta$ , then*

$$\mathbb{P}^{x_0}(\tau_{B(x_0,1)} < \tau_D) < \varepsilon.$$



*Proof.* Let  $z \in \partial D$ . We will show that there exists  $\rho \in (0, \frac{1}{4})$  and  $\sigma \in (\frac{1}{2}, 1)$  such that if  $x \in D \cap B(z, \rho^i)$ , then

$$h(x) = \mathbb{P}^x(\tau_{B(z, 1/2)} < \tau_D) \leq 2\sigma^i.$$

This is true for  $i = 1$ . We suppose it is true for all  $j \leq i$  and prove it for  $i + 1$ . Let  $U_i = B(z, \rho^i)$ .

If  $x \in U_{i+1}$ , then since  $h$  is regular  $\alpha$ -harmonic in  $B(z, 1/2) \cap D$  and 0 in  $D^c$ ,

$$\begin{aligned} h(x) &= \mathbb{E}^x[h(X_{\tau_{U_i} \wedge \tau_D})] \\ &\leq \mathbb{E}^x[h(X_{\tau_{U_i}}); \tau_{U_i} < \tau_D, X_{\tau_{U_i}} \in U_{i-1}] \\ &\quad + \sum_{j=2}^{i-1} \mathbb{E}^x[h(X_{\tau_{U_i}}); X_{\tau_{U_i}} \in U_{i-j} \setminus U_{i-j+1}] \\ &\quad + \mathbb{E}^x[h(X_{\tau_{U_i}}); X_{\tau_{U_i}} \notin U_1]. \end{aligned}$$

Using (3.1) and the induction hypothesis, there exists  $a > 0$  not depending on  $i$  such that the first term on the right is bounded by  $2(1 - a)\sigma^{i-1}$ . Using the induction hypothesis and (2.3), the second term is bounded by

$$2c_1 \sum_{j=2}^{i-1} \sigma^{i-j} \rho^{j\alpha}.$$

Finally, using (2.3), the third term is bounded by  $c_2\rho^{i\alpha}$ . We will take  $\rho$  small enough so that  $\rho^\alpha/\sigma < \frac{1}{2}$ . Adding the three terms together, we have the bound

$$\begin{aligned} 2(1 - a)\sigma^{i-1} + 2c_1 \sum_{j=2}^{i-1} \sigma^{i-j} \rho^{j\alpha} + c_2\rho^{i\alpha} \\ \leq 2(1 - a)\sigma^{i-1} + c_3\sigma^i(\rho^\alpha/\sigma) + c_2\rho^{i\alpha} \\ \leq 2\sigma^{i-1}[(1 - a) + c_3(\rho^\alpha/\sigma) + c_4(\rho^\alpha/\sigma)]. \end{aligned}$$

We choose  $\rho$  small so that  $\rho^\alpha/\sigma < \frac{1}{2}$  and  $2(c_3 + c_4)\rho < a/2$  and then choose  $\sigma \in (\sqrt{1 - a/2}, 1)$ . So the above is bounded by

$$2\sigma^{i-1}[(1 - a) + (a/2)] \leq 2\sigma^{i-1}(1 - a/2) \leq 2\sigma^{i+1},$$

and the induction step is established. The result follows easily from this. □

*Remark.* The above proposition actually shows that  $\mathbb{P}^{x_0}(\tau_{B(z, 1/2)} < \tau_D)$  goes to 0 as some power of  $v(x_0)$ . An argument using the boundary Harnack principle shows that the same is true for the Green function  $G_D(x_0, y)$  if  $y$  is fixed. This complements the estimates in [18] on the behavior of the Green function.

**Proposition 3.6.** *Let  $\varepsilon > 0$ . There exist  $\eta, K > 0$  such that if  $v(x_0) \leq \eta$ , then*

$$\mathbb{P}^{x_0}(|X_{\tau_D} - x_0| > K) < \varepsilon.$$

*Proof.* By Proposition 3.5, there exists  $\eta$  such that

$$\mathbb{P}^{x_0}(\tau_{B(x_0,1)} < \tau_D) < \varepsilon/2 \tag{3.15}$$

if  $v(x_0) \leq \eta$ . Provided  $K > 1$ , notice that it is not possible that  $|X_{\tau_D} - x_0| > K$  on the event where  $\tau_{B(x_0,1)} > \tau_D$ . So we have

$$\begin{aligned} \mathbb{P}^{x_0}(|X_{\tau_D} - x_0| > K) &= \mathbb{P}^{x_0}(|X_{\tau_D} - x_0| > K, \tau_{B(x_0,1)} < \tau_D) \\ &\quad + \mathbb{P}^{x_0}(|X_{\tau_D} - x_0| > K, \tau_{B(x_0,1)} = \tau_D). \end{aligned}$$

The first term on the right is less than  $\varepsilon/2$  by (3.15). Using (2.3) let us choose  $K > 1$  large so that

$$\mathbb{P}^{x_0}(|X_{\tau_{B(x_0,1)}} - x_0| > K) < \varepsilon/2.$$

So the second term on the right is also less than  $\varepsilon/2$ . □

A scaling argument shows that if  $v(x_0) \leq 1$ , then  $\mathbb{P}^{x_0}(|X_{\tau_D} - x_0| > K/\eta) < \varepsilon$ . Let  $M = K/\eta$ . Using scaling again, we have

**Corollary 3.7.** *Let  $\varepsilon > 0$ . There exists  $M > 0$  such that if  $x_0 \in D$ , then*

$$\mathbb{P}^{x_0}(|X_{\tau_D} - x_0| > Mv(x_0)) < \varepsilon.$$

#### 4. Maximal functions

Let  $D$  be the region above a bounded Lipschitz function  $\Gamma : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$  with Lipschitz constant  $\lambda$ . Set  $\lambda_0 = 6(\lambda + 1)$ . For  $i$  an integer and  $\tilde{x} \in \mathbb{R}^{d-1}$ , let

$$\begin{aligned} A_i(\tilde{x}) &= \{(\tilde{t}, t_d) \in \mathbb{R}^d : |t_k - x_k| \leq 2^{-i-1}, k = 1, \dots, d-1, \\ &\quad |t_d - (\Gamma(\tilde{x}) - 3\lambda_0 2^{-i-1})| \leq 2^{-i-1}\lambda_0\}, \end{aligned} \tag{4.1}$$

and let  $\tilde{A}_i(\tilde{x})$  be the  $(d-1)$ -dimensional cube with center  $\tilde{x}$  and side length  $2^{-i}$ . Note that  $|A_i(\tilde{x})| = 2^{-id}\lambda_0$ . Set

$$B_i = \{\tilde{x} \in \mathbb{R}^{d-1} : \text{each coordinate of } \tilde{x} \text{ is a multiple of } 2^{-i}\}.$$

Let us define for  $i \geq 0$  and  $L \geq 0$

$$\begin{aligned} F_i(\tilde{x}) &= \frac{1}{|A_i(\tilde{x})|} \int_{A_i(\tilde{x})} |f(y)| dy, \quad G_{iL}(\tilde{x}) = \sup_{\tilde{y} \in B_i, |\tilde{y}| \leq L 2^{-i}} F_i(\tilde{x} + \tilde{y}). \\ M_L f(\tilde{x}) &= \sup_{i \geq 0} G_{iL}(\tilde{x}). \end{aligned}$$

We use  $\|\cdot\|_1$  to denote the  $L^1$  norm of a function with respect to  $(d-1)$ -dimensional Lebesgue measure.

The following proposition's proof is similar to the analogous one in [3], but we take this opportunity to correct some errors in that paper. Note also that  $G_{iL}$  is defined differently here.

**Proposition 4.1.** *Suppose  $f \in \Lambda_\beta^{p,\infty}(\mathbb{R}^d)$ ,  $\beta p > 1$ ,  $p > 1$ , and the support of  $f$  is contained in  $B(0, J)$  for some  $J \geq 1$ . Then*

$$\|M_L f(\tilde{x})\|_1 < \kappa \|f\|_{\Lambda_\beta^{p,\infty}(\mathbb{R}^d)}. \tag{4.2}$$

The constant  $\kappa$  depends on  $d, p, L, J, \lambda$ , and  $\beta$ . In particular,  $M_L f(\tilde{x})$  is finite a.e.

*Proof.* Since

$$\begin{aligned} |G_{iL}(\tilde{x}) - G_{i+1,L}(\tilde{x})| &\leq \sup_{\tilde{y}, \tilde{z} \in \mathcal{B}_{i+1}, |\tilde{y}| \vee |\tilde{z}| \leq 2L2^{-i}} |F_i(\tilde{x} + \tilde{y}) - F_{i+1}(\tilde{x} + \tilde{z})| \\ &\leq \sum_{\tilde{y}, \tilde{z} \in \mathcal{B}_{i+1}, |\tilde{y}| \vee |\tilde{z}| \leq 2L2^{-i}} |F_i(\tilde{x} + \tilde{y}) - F_{i+1}(\tilde{x} + \tilde{z})|, \end{aligned}$$

we have

$$\begin{aligned} \|G_{iL}(\tilde{x}) - G_{i+1,L}(\tilde{x})\|_1 &\leq \sum_{\tilde{y}, \tilde{z} \in \mathcal{B}_{i+1}, |\tilde{y}| \vee |\tilde{z}| \leq 2L2^{-i}} \|F_i(\tilde{x} + \tilde{y}) - F_{i+1}(\tilde{x} + \tilde{z})\|_1 \\ &\leq c_1 L^{d-1} \sup_{\tilde{w} \in \mathcal{B}_{i+1}, |\tilde{w}| \leq 4L2^{-i}} \|F_i(\tilde{x}) - F_{i+1}(\tilde{x} + \tilde{w})\|_1. \end{aligned} \tag{4.3}$$

Let  $A_{ij}(\tilde{x})$ ,  $j = 1, \dots, 2^d$ , be a subdivision of  $A_i(\tilde{x})$  into  $2^d$  equal rectangular solids, each of which is congruent to  $A_{i+1}(\tilde{x})$ ; to be more precise, if  $A_i(\tilde{x}) = \prod_{i=1}^d [a_i, b_i]$ , then each  $A_{ij}(\tilde{x})$  is of the form  $\prod_{i=1}^d [d_i, e_i]$ , where for each  $i$  either  $d_i = a_i$  and  $e_i = (a_i + b_i)/2$  or else  $d_i = (a_i + b_i)/2$  and  $e_i = b_i$ .

Fix a  $\tilde{w} \in \mathcal{B}_{i+1}$  with  $|\tilde{w}| \leq 4L2^{-i}$ . Let  $T_{ij} : \mathbb{R}^d \rightarrow \mathbb{R}^d$  be the linear map defined by

$$T_{ij}(\tilde{x}, x_d) = (\tilde{x} + \tilde{t}_{ij}, x_d + t_{ij}^d),$$

where  $t_{ij} = (\tilde{t}_{ij}, t_{ij}^d)$  are points such that  $A_{ij}(\tilde{x}) = T_{ij}(A_{i+1}(\tilde{x} + \tilde{w}))$ ; we can find a constant  $c_2$  not depending on  $i$  such that  $|t_{ij}| \leq c_2 2^{-i}$  for  $j = 1, \dots, 2^d$ .

Then

$$\begin{aligned} &F_i(\tilde{x}) - F_{i+1}(\tilde{x} + \tilde{w}) \\ &= \frac{1}{|A_i(\tilde{x})|} \int_{A_i(\tilde{x})} |f(y)| dy - \frac{1}{|A_{i+1}(\tilde{x} + \tilde{w})|} \int_{A_{i+1}(\tilde{x} + \tilde{w})} |f(y)| dy \\ &= \sum_{j=1}^{2^d} \frac{1}{|A_i(\tilde{x})|} \left[ \int_{A_{ij}(\tilde{x})} |f(y)| dy - \int_{A_{i+1}(\tilde{x} + \tilde{w})} |f(y)| dy \right] \\ &= \sum_{j=1}^{2^d} 2^{id} \lambda_0^{-1} \left[ \int_{T_{ij}(A_{i+1}(\tilde{x} + \tilde{w}))} |f(y)| dy - \int_{A_{i+1}(\tilde{x} + \tilde{w})} |f(y)| dy \right] \\ &= \sum_{j=1}^{2^d} 2^{id} \lambda_0^{-1} \int_{A_{i+1}(\tilde{x} + \tilde{w})} (|f(T_{ij}(y))| - |f(y)|) dy. \end{aligned} \tag{4.4}$$

Integrating (4.4) gives

$$\begin{aligned} & \int_{\mathbb{R}^{d-1}} |F_i(\tilde{x} + \tilde{w}) - F_{i+1}(\tilde{x})| d\tilde{x} \\ & \leq \sum_{j=1}^{2^d} 2^{id} \lambda_0^{-1} \int_{\mathbb{R}^{d-1}} \int_{A_{i+1}(\tilde{x} + \tilde{w})} |f(y) - f(T_{ij}(y))| dy d\tilde{x} \\ & = \sum_{j=1}^{2^d} 2^{id} \lambda_0^{-1} \int_{\mathbb{R}^{d-1}} \int_{\tilde{A}_{i+1}(\tilde{x} + \tilde{w})} \int_{\Gamma(\tilde{x} + \tilde{w}) - 2^{-i} \lambda_0}^{\Gamma(\tilde{x} + \tilde{w}) - 2^{-i-1} \lambda_0} |f(y) - f(T_{ij}(y))| dy_d d\tilde{y} d\tilde{x}. \end{aligned}$$

Let  $\tilde{z} = \tilde{y} - (\tilde{x} + \tilde{w})$ ,  $z_d = y_d$ . Then by Fubini's theorem the last expression can be written as

$$\begin{aligned} & \sum_{j=1}^{2^d} 2^{id} \lambda_0^{-1} \int_{\tilde{A}_{i+1}(\tilde{0})} \int_{\mathbb{R}^{d-1}} \int_{\Gamma(\tilde{x} + \tilde{w}) - 2^{-i} \lambda_0}^{\Gamma(\tilde{x} + \tilde{w}) - 2^{-i-1} \lambda_0} |f(\tilde{z} + \tilde{x} + \tilde{w}, z_d) \\ & \quad - f(\tilde{z} + \tilde{x} + \tilde{w} + \tilde{t}_{ij}, z_d + t_{ij}^d)| dz_d d\tilde{x} d\tilde{z} \\ & \leq \sum_{j=1}^{2^d} 2^{id} \lambda_0^{-1} \int_{\tilde{A}_{i+1}(\tilde{0})} \left( \int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}} |f(\tilde{z} + \tilde{x} + \tilde{w}, z_d) \right. \\ & \quad \left. - f(\tilde{z} + \tilde{x} + \tilde{w} + \tilde{t}_{ij}, z_d + t_{ij}^d)|^p dz_d d\tilde{x} \right)^{1/p} \\ & \quad \times \left( \int_{\mathbb{R}^{d-1}} \int_{\mathbb{R}} \chi_{B(0, J) \cap (\mathbb{R}^{d-1} \times [-2^{-i}, -2^{-i-1}])} dz_d d\tilde{x} \right)^{1/q} d\tilde{z} \\ & \leq \sum_{j=1}^{2^d} 2^{id} \lambda_0^{-1} \int_{\tilde{A}_{i+1}(\tilde{0})} |\tilde{t}_{ij}, t_{ij}^d|^\beta \\ & \quad \times \|f\|_{\Lambda_\beta^{p, \infty}(\mathbb{R}^d)} J^{(d-1)/q} 2^{-i/q} d\tilde{z}, \end{aligned} \tag{4.5}$$

where we use Hölder's inequality with  $p^{-1} + q^{-1} = 1$ . From (4.5) we have

$$\int_{\mathbb{R}^{d-1}} |F_i(\tilde{x}) - F_{i+1}(\tilde{x} + \tilde{w})| d\tilde{x} \leq c_3 \sum_{j=1}^{2^d} 2^i |t_{ij}|^\beta \|f\|_{\Lambda_\beta^{p, \infty}(\mathbb{R}^d)} 2^{-i/q}. \tag{4.6}$$

Since  $|t_{ij}| \leq c_4 L 2^{-i}$ , for  $j = 1, \dots, 2^d$ ,  $\tilde{w} \in \mathcal{B}_{i+1}$ , and  $|\tilde{w}| \leq 4L 2^{-i}$ , from (4.3) and (4.6) we have

$$\begin{aligned} \|G_{iL}(\tilde{x}) - G_{i+1, L}(\tilde{x})\|_1 & \leq c_5 2^{i(1-1/q)} (2^{-i})^\beta \|f\|_{\Lambda_\beta^{p, \infty}(\mathbb{R}^d)} \\ & \leq c_5 2^{((1/p) - \beta)i} \|f\|_{\Lambda_\beta^{p, \infty}(\mathbb{R}^d)}. \end{aligned} \tag{4.7}$$

To prove  $\|M_L f(\tilde{x})\|_1 < \kappa \|f\|_{\Lambda_\beta^{p, \infty}(\mathbb{R}^d)}$ , note that

$$\sup_i |G_{iL}(\tilde{x})| \leq |G_{0L}(\tilde{x})| + \sum_{i=1}^{\infty} |G_{iL}(\tilde{x}) - G_{i-1, L}(\tilde{x})|.$$

Clearly  $\|G_{0L}\|_1 < c_6\|f\|_p$  because  $f$  is in  $L^p$  and has compact support. Hence by (4.7)

$$\begin{aligned} \|M_L f(\tilde{x})\|_1 &= \|\sup_i G_{iL}(\tilde{x})\|_1 \leq \|G_{0L}(\tilde{x})\|_1 + \sum_{i=1}^{\infty} \|G_{iL}(\tilde{x}) - G_{i-1,L}(\tilde{x})\|_1 \\ &\leq c_6\|f\|_p + c_7\|f\|_{\Lambda_{\beta}^{p,\infty}(\mathbb{R}^d)} \sum_{i=1}^{\infty} 2^{((1/p)-\beta)i} \\ &< c_8\|f\|_{\Lambda_{\beta}^{p,\infty}(\mathbb{R}^d)}, \end{aligned} \tag{4.8}$$

if  $\beta p > 1$ . □

### 5. The Fatou theorem

In this section we prove Theorem 1.1.

**Proposition 5.1.** *Suppose  $D$  is the region above the graph of a Lipschitz function with Lipschitz constant  $\lambda$  and  $f$  is bounded, has compact support, and is in  $\Lambda_{\beta}^{p,\infty}(\mathbb{R}^d)$  with  $\beta p > 1$ . Then  $u_f$  converges nontangentially at almost every point on the boundary.*

*Proof.* Let  $\varepsilon > 0$ . By Corollary 3.4 there exists  $a_1$  such that for  $x \in D$

$$\mathbb{P}^x(X_{\tau_D} \in S_{a_1 v(x)}) < \varepsilon.$$

By Corollary 3.7 there exists  $a_2$  such that for  $x \in D$

$$\mathbb{P}^x(X_{\tau_D} \in B(x, a_2 \delta_D(x))^c) < \varepsilon.$$

Recall the definition of  $A_i(\tilde{x})$  given in (4.1). We claim that if  $v(x)$  is sufficiently small, there exist positive integers  $b_1, b_2$ , and  $L$  independent of  $v(x)$  and an integer  $i_0 \geq b_1 + 1$  depending on  $x$  such that

$$\left( D^c \cap B(x, a_2 \delta_D(x)) \right) \setminus S_{a_1 v(x)} \subset \bigcup_{i=i_0-b_1}^{i_0+b_2} \left( \bigcup_{\{\tilde{t}_j \in B_i, |\tilde{t}_j| \leq L2^{-i}\}} A_i(\tilde{x} + \tilde{t}_j) \right). \tag{5.1}$$

To see this, first choose  $i_0$  such that  $\lambda_0 2^{-i_0} \leq \delta_D(x) < \lambda_0 2^{-i_0+1}$ . We can then choose  $b_1$  and  $b_2$  independently of  $i_0$  such that if  $y$  is an element of the left hand side of (5.1), then  $y \in \bigcup_{i=i_0-b_1}^{i_0+b_2} A_i(\tilde{y})$ . Now if  $y$  is an element of the left hand side of (5.1), then the distance between  $\tilde{x}$  and  $\tilde{y}$  is less than  $a_2 \delta_D(x) \leq c_1 2^{-i_0}$ ; so if we take  $L$  larger than  $c_1 2^{b_2+1}$ , then  $y$  will be an element of the right hand side of (5.1). This proves (5.1).

Choose  $\beta' \in (1/p, \beta)$ . Let  $\kappa$  be the constant in (4.2) when we replace  $\beta$  by  $\beta'$ ; recall that  $\kappa$  depends on  $L$ . By [3], Lemma 4.1, we can choose  $h \in \Lambda_{\beta'}^{p,\infty}(\mathbb{R}^d)$  such that  $\|h\|_{\infty} \leq 2\|f\|_{\infty}$ ,

$$\|h\|_{\Lambda_{\beta'}^{p,\infty}(\mathbb{R}^d)} < \varepsilon^2 / (\kappa(b_1 + b_2 + 1)L^{d-1}),$$

and  $g = f - h$  is continuous with compact support. For any function  $k$  on  $D$ , define

$$\Omega k(x) = \limsup_{z \in C(x), z \rightarrow x} u_k(z) - \liminf_{z \in C(x), z \rightarrow x} u_k(z)$$

for  $x \in \partial D$ . We want to show that if  $x_0 \in \partial D$ , then

$$\Omega f(x_0) \leq c_2 \varepsilon \|f\|_\infty + c_3 Mh(x_0). \tag{5.2}$$

If  $z \in A_i(\tilde{x} + t_j)$ , then  $P_D(x, z) \leq c_4 \delta_D(z)^{-d}$  by Proposition 2.1. Suppose  $x \in D$  with  $\tilde{x} = \tilde{x}_0$ . Writing  $h^+$  for the positive part of  $h$  and defining  $u_{h^+}$  analogously to (1.1), we have

$$\begin{aligned} u_{h^+}(x) &\leq \mathbb{E}^x[h^+(X_{\tau_D}); X_{\tau_D} \in S_{a_1 v(x)}] + \mathbb{E}^x[h^+(X_{\tau_D}); X_{\tau_D} \notin B(x, a_2 \delta_D(x))] \\ &\quad + \sum_{i=i_0-b_1}^{i_0+b_2} \sum_{\{\tilde{t}_j \in \mathcal{B}_i, |\tilde{t}_j| \leq L2^{-i}\}} \mathbb{E}^x[h^+(X_{\tau_D}); X_{\tau_D} \in A_i(\tilde{x} + \tilde{t}_j)] \\ &\leq 2\|h\|_\infty \varepsilon + \sum_{i=i_0-b_1}^{i_0+b_2} \sum_{\{\tilde{t}_j \in \mathcal{B}_i, |\tilde{t}_j| \leq L2^{-i}\}} \int_{A_i(\tilde{x} + \tilde{t}_j)} h^+(z) P_D(x, z) dz \\ &\leq 2\|h^+\|_\infty \varepsilon + c_5 \sum_{i=i_0-b_1}^{i_0+b_2} \sum_{\{\tilde{t}_j \in \mathcal{B}_i, |\tilde{t}_j| \leq L2^{-i}\}} F_i(\tilde{x} + \tilde{t}_j) \\ &\leq 2\|h^+\|_\infty \varepsilon + c_6 \sum_{i=i_0-b_1}^{i_0+b_2} L^{d-1} G_{iL}(\tilde{x}) \\ &\leq 2\|h^+\|_\infty \varepsilon + c_6(b_1 + b_2 + 1)L^{d-1} M_L h^+(\tilde{x}), \end{aligned}$$

where we define  $F_i$  and  $G_{iL}$  in terms of  $h^+$  instead of  $f$ . By the Harnack inequality for nonnegative  $\alpha$ -harmonic functions,

$$\begin{aligned} \limsup_{y \in C(x_0), y \rightarrow x_0} u_{h^+}(y) &\leq c_7 \|h^+\|_\infty \varepsilon + c_7(b_1 + b_2 + 1)L^{d-1} M_L h^+(\tilde{x}_0) \\ &\leq c_8 \|f\|_\infty \varepsilon + c_8(b_1 + b_2 + 1)L^{d-1} M_L h(\tilde{x}_0). \end{aligned}$$

We have a similar estimate when  $h^+$  is replaced by  $h^-$ . Since  $g$  is continuous with compact support,

$$\begin{aligned} \Omega f(x_0) &\leq \Omega g(x_0) + \Omega h(x_0) = \Omega h(x_0) \\ &\leq 2c_8 \|f\|_\infty \varepsilon + 2c_8(b_1 + b_2 + 1)L^{d-1} M_L h(\tilde{x}_0) \end{aligned}$$

as desired.

By Proposition 4.1,

$$\begin{aligned} |\{x_0 \in \partial D : (b_1 + b_2 + 1)L^{d-1} M_L h(\tilde{x}_0) > \varepsilon\}| &\leq \frac{(b_1 + b_2 + 1)L^{d-1}}{\varepsilon} \|M_L h\|_1 \\ &\leq \frac{\kappa(b_1 + b_2 + 1)L^{d-1}}{\varepsilon} \|h\|_{\Lambda_{\beta'}^{p, \infty}(\mathbb{R}^d)} \\ &\leq c_9 \varepsilon. \end{aligned}$$

Recall that  $(d - 1)$ -dimensional Lebesgue measure is comparable to surface measure. Therefore, except for a set of surface measure at most  $c_{10}\varepsilon$  we have

$$\Omega f(x_0) \leq \Omega h(x_0) \leq (2c_8\|f\|_\infty + 2c_8)\varepsilon. \tag{5.3}$$

Since  $\varepsilon$  is arbitrary, this proves the proposition. □

To handle bounded Lipschitz domains we first need some lemmas.

**Lemma 5.2.** *Let  $D$  be a bounded Lipschitz domain,  $z \in \partial D$ , and  $M > 0$ . If  $x_n \in D, n = 1, 2, \dots$ , and  $x_n \rightarrow z$ , then  $\mathbb{P}^{x_n}(X_{\tau_D} \in B(z, M) \cap D^c) \rightarrow 1$ .*

*Proof.*  $z \in \partial D$  is regular for  $D^c$  (the proof that every point in  $\partial D$  is regular for the complement of the domain is the same as that for Brownian motion; see [1], Proposition II.1.13), and so  $\mathbb{P}^{x_n}(\tau_D \leq t) \rightarrow 1$  for all  $t > 0$ ; see [1], Corollary II.1.11. Let  $\varepsilon > 0$  and choose  $t$  such that  $\mathbb{P}^0(\sup_{s \leq t} |X_s| \geq M/2) < \varepsilon$ ; this is possible by the right continuity of the paths of  $X_t$ . We have  $|x_n - z| < M/2$  and  $\mathbb{P}^{x_n}(\tau_D \leq t) \geq 1 - \varepsilon$  for sufficiently large  $n$ . So we have

$$\begin{aligned} \mathbb{P}^{x_n}(X_{\tau_D} \in B(z, M) \cap D^c) &\geq \mathbb{P}^{x_n}(\tau_D \leq t, \sup_{s \leq t} |X_s - x_n| < M/2) \\ &\geq \mathbb{P}^{x_n}(\tau_D \leq t) - \mathbb{P}^0(\sup_{s \leq t} |X_s| \geq M/2) \\ &\geq 1 - \varepsilon - \varepsilon. \end{aligned}$$

□

**Lemma 5.3.** *Let  $D$  be the region above the graph of a Lipschitz function with Lipschitz constant  $\lambda, z \in \partial D$ , and  $M > 0$ , and let  $f$  be a bounded function. If  $x_n \in D, x_n \rightarrow z$ , and either  $\lim_{x_n \rightarrow z} \mathbb{E}^{x_n} f(X_{\tau_D})$  or  $\lim_{x_n \rightarrow z} \mathbb{E}^{x_n} f(X_{\tau_{D \cap B(z, M)}})$  exists, then they both exist and coincide.*

*Proof.* Letting  $B_M = D \cap B(z, M)$ , we have

$$\begin{aligned} \mathbb{E}^{x_n} f(X_{\tau_D}) &= \mathbb{E}^{x_n}[f(X_{\tau_D}); \tau_{B_M} < \tau_D] + \mathbb{E}^{x_n}[f(X_{\tau_D}); \tau_{B_M} = \tau_D] \\ &= \mathbb{E}^{x_n}[f(X_{\tau_D}); \tau_{B_M} < \tau_D] + \mathbb{E}^{x_n}[f(X_{\tau_{B_M}}); \tau_{B_M} = \tau_D] \\ &= \mathbb{E}^{x_n}[f(X_{\tau_D}); \tau_{B_M} < \tau_D] + \mathbb{E}^{x_n} f(X_{\tau_{B_M}}) \\ &\quad - \mathbb{E}^{x_n}[f(X_{\tau_{B_M}}); \tau_{B_M} < \tau_D]. \end{aligned} \tag{5.4}$$

Note that the first and third terms of the last line are both bounded by

$$\|f\|_\infty \mathbb{P}^{x_n}(\tau_{B_M} < \tau_D).$$

Since

$$\mathbb{P}^{x_n}(\tau_{B_M} < \tau_D) = 1 - \mathbb{P}^{x_n}(\tau_{B_M} = \tau_D) \leq 1 - \mathbb{P}^{x_n}(X_{\tau_D} \in B(z, M) \cap D^c) \rightarrow 0$$

as  $x_n \rightarrow z$  by Lemma 5.2, the first and the last terms in the last line of (5.4) go to 0 as  $x_n \rightarrow z$ , which completes the proof. □

We now prove Theorem 1.1.

*Proof.* First we will consider the case when  $D$  is the region above the graph of a Lipschitz function  $\Gamma$  with Lipschitz constant  $\lambda$ . Suppose that  $f$  is bounded,  $f \in \Lambda_\beta^{p,\infty}(\mathbb{R}^d)$ , and  $\beta p > 1$ . Let  $M > 0$ . We will show nontangential convergence for  $\tilde{x}$  in  $B(0, M/2) \cap \partial D$ . Since  $M$  is arbitrary, the theorem in this case will follow. Let  $\varphi \in C^\infty$  be a cut-off function such that  $\varphi = 1$  on  $B(0, 2M)$  and  $0$  on  $B(0, 3M)^c$ . Since  $f$  is bounded,  $f\varphi$  is bounded and supported on  $B(0, 3M)$ . Since  $\varphi$  is smooth, then  $f\varphi \in \Lambda_\beta^{p,\infty}(\mathbb{R}^d)$ , so by Proposition 5.1 the Fatou theorem holds for this function. Note that  $f = f\varphi + f(1 - \varphi)$  and so  $u_f(x) = u_{f\varphi}(x) + u_{f(1-\varphi)}(x)$ . Thus it is enough to show for  $z \in B(0, M/2) \cap \partial D$  that  $\lim_{x_n \rightarrow z} u_{f(1-\varphi)}(x_n) = 0$  if  $x_n \in D$ . Since the support of  $f(1 - \varphi)$  is contained in  $B(0, 2M)^c \subset B(z, 3M/2)^c$ , we have

$$u_{f(1-\varphi)}(x_n) = \mathbb{E}^{x_n} f(1 - \varphi)(X_{\tau_D}) \leq \|f(1 - \varphi)\|_\infty \mathbb{P}^{x_n}(X_{\tau_D} \in B(z, M)^c \cap D^c).$$

By Lemma 5.2 the right hand side goes to 0; this completes the proof in the first case.

Next we consider a bounded Lipschitz domain  $D$ . For every  $z \in \partial D$  there exist  $M > 0$  and a Lipschitz function  $\Gamma$  with Lipschitz constant  $\lambda$  (which can depend on  $x$ ) such that in some coordinate system  $D \cap B(z, M) = D_\Gamma \cap B(z, M)$ , where  $D_\Gamma$  is the region above the graph of  $\Gamma$ . Suppose  $f$  is bounded and in  $\Lambda_\beta^{p,\infty}(\mathbb{R}^d)$  with  $\beta p > 1$ . By the paragraph above, nontangential limits exist a.e. in  $D_\Gamma$ . By Lemma 5.3 the nontangential limits for  $D_\Gamma$  and for  $D_\Gamma \cap B(z, M)$  in  $B(z, M)$  are the same. By the same reasoning as in the proof of Lemma 5.3 the nontangential limits for  $D$  and for  $D \cap B(z, M)$  in  $B(z, M)$  are the same. Hence the nontangential limits for  $D$  are the same as those for  $D_\Gamma$  in  $B(z, M)$ .

Finally, suppose  $D$  is a bounded Lipschitz domain and  $f \in \Lambda_{loc}^b(\beta, p, D^c)$ . If  $z \in \partial D$ , there exists  $\zeta$  (depending on  $z$ ) such that  $f$  agrees on  $B(z, \zeta) \cap D^c$  with a bounded function,  $\tilde{f}$ , say, that is in  $\Lambda_\beta^{p,\infty}(\mathbb{R}^d)$ . Let  $\varphi$  be a  $C^\infty$  cutoff function that is 1 on  $B(z, \zeta/2)$  and 0 on  $B(z, \zeta)^c$ . By the preceding paragraph, nontangential limits exists a.e. for  $u_{f\varphi} = u_{\tilde{f}\varphi}$  in  $B(z, \zeta/4) \cap D$ . By the argument in the first paragraph of this proof, nontangential limits also exist there for  $u_{f(1-\varphi)}$ . Therefore every point of  $\partial D$  has a neighborhood in which nontangential convergence in  $D$  holds; this proves the theorem.  $\square$

*Remark.* A function  $f$  that is in  $\Lambda_\beta^{p,\infty}(\mathbb{R}^d)$  need not be continuous unless  $\beta p > d$ , and so in general its value on  $\partial D$  is not defined in a pointwise manner. However if  $\beta' p > 1$ , one can define a bounded trace operator  $\mathcal{T} : \Lambda_{\beta'}^{p,\infty} \rightarrow L^p(\partial D)$  which does give a definition of the restriction of  $f$  to  $\partial D$ ; moreover the definition of  $\mathcal{T}(f)$  is independent of the precise value of  $\beta'$ . A proof of this may be derived from the results in [22], Chapter VI. If  $\beta p > 1$ , choose  $\beta' \in (1/p, \beta)$ . As in the proof of Proposition 5.1, given  $\varepsilon$ , a function  $f \in \Lambda_\beta^{p,\infty}(\mathbb{R}^d)$  can be written as the sum of a continuous function with compact support  $g$  plus a function  $h$ , where

$$\|h\|_{\Lambda_{\beta'}^{p,\infty}(\mathbb{R}^d)} < \varepsilon.$$

Since  $g$  is continuous, it is easy to see that the nontangential limit of  $u_g$  agrees with  $\mathcal{T}(g)$  at every point of the boundary. Let  $\delta > 0$ . The proof of Proposition 5.1



(see (5.3)) shows that the nontangential limit of  $u_h$  is less than  $\delta$  except for a set of surface measure less than  $\delta$  provided we take  $\varepsilon$  small enough. Since  $\mathcal{T}$  is a bounded operator, then  $\mathcal{T}(h)$  will be less than  $\delta$  except for a set of surface measure less than  $\delta$  provided we take  $\varepsilon$  small enough. Therefore the nontangential limit of  $u_f$  differs from  $\mathcal{T}$  by at most  $2\delta$ , except for a set of surface measure at most  $2\delta$ . Since  $\delta$  is arbitrary, this shows that the nontangential limit of  $u_f$  agrees with  $\mathcal{T}(f)$  almost everywhere.

*Acknowledgements.* We would like to thank Zhenqing Chen and James Norris for helpful advice regarding this paper.

## References

1. Bass, R.F.: Probabilistic Techniques in Analysis. Springer Verlag, Berlin, 1995
2. Bass, R.F., Levin, D.A.: Harnack inequalities for jump processes. *Potential Anal.* **17**, 375–388 (2002)
3. Bass, R.F., You, D.: A Fatou theorem for  $\alpha$ -harmonic functions. *Bull. Sci. Math.* **127**, 635–648 (2003)
4. Blumenthal, R.M., Gettoor, R.K.: Markov Processes and Their Potential Theory. Academic Press, New York, 1968
5. Bogdan, K.: The boundary Harnack principle for the fractional Laplacian. *Studia Math.* **123**, 43–80 (1997)
6. Bogdan, K.: Representation of  $\alpha$ -harmonic functions in Lipschitz domains. *Hiroshima Math. J.* **29**, 227–243 (1999)
7. Bogdan, K., Byczkowski, T.: On the Schrödinger operator based on the fractional Laplacian. *Bull. Polish Acad. Sci. Math.* **49**, 291–301 (2001)
8. Bogdan, K., Byczkowski, T.: Potential theory of Schrödinger operator based on fractional Laplacian. *Probab. Math. Statist.* **20**, 293–335 (2000)
9. Bogdan, K., Byczkowski, T.: Probabilistic proof of boundary Harnack principle for  $\alpha$ -harmonic functions. *Potential Anal.* **11**, 135–156 (1999)
10. Bogdan, K., Byczkowski, T.: Potential theory for the  $\alpha$ -stable Schrödinger operator on bounded Lipschitz domains. *Studia Math.* **133**, 53–92 (1999)
11. Bogdan, K., Kulczycki, T., Nowak, A.: Gradient estimates for harmonic and  $q$ -harmonic functions of symmetric stable processes. *Illinois J. Math.* **46**, 541–556 (2002)
12. Bogdan, K., Stós, A., Sztonyk, P.: Potential theory for Lévy stable processes. *Bull. Polish Acad. Sci. Math.* **50**, 361–372 (2002)
13. Chen, Z.-Q.: Multidimensional symmetric stable processes. *Korean J. Comput. Appl. Math.* **6**, 227–266 (1999)
14. Chen, Z.-Q., Song, R.: Martin boundary and integral representation for harmonic functions of symmetric stable processes. *J. Funct. Anal.* **159**, 267–294 (1998)
15. Chen, Z.-Q., Song, R.: Estimates on Green functions and Poisson kernels for symmetric stable processes. *Math. Ann.* **312**, 465–501 (1998)
16. Hunt, R.A., Wheeden, R.L.: On the boundary values of harmonic functions. *Trans. Am. Math. Soc.* **132**, 307–322 (1968)
17. Ikeda, N., Watanabe, S.: On some relations between the harmonic measure and the Lévy measure for a certain class of Markov processes. *J. Math. Kyoto Univ.* **2**, 79–95 (1962)
18. Jakubowski, T.: The estimates for the Green function in Lipschitz domains for the symmetric stable processes. *Probab. Math. Statist.* **22**, 419–441 (2002)
19. Landkof, N.S.: Foundations of Modern Potential Theory. Springer, New York, 1972

20. Song, R., Wu, J.-M.: Boundary Harnack principle for symmetric stable processes. *J. Funct. Anal.* **168**, 403–427 (1999)
21. Stein, E.M.: *Singular Integrals and Differentiability Properties of Functions*. Princeton Univ. Press, Princeton, 1970
22. Jonsson, A., Wallin, H.: Function spaces on subsets of  $R^n$ . *Math. Rep.* **2** (1), 1984
23. Wu, J.-M.: Harmonic measures for symmetric stable processes. *Studia Math.* **149**, 281–293 (2002)