

Geometric ergodicity and the spectral gap of non-reversible Markov chains

I. Kontoyiannis · S. P. Meyn

Received: 29 June 2009 / Revised: 14 May 2011 / Published online: 3 June 2011
© Springer-Verlag 2011

Abstract We argue that the spectral theory of non-reversible Markov chains may often be more effectively cast within the framework of the naturally associated weighted- L_∞ space L_∞^V , instead of the usual Hilbert space $L_2 = L_2(\pi)$, where π is the invariant measure of the chain. This observation is, in part, based on the following results. A discrete-time Markov chain with values in a general state space is geometrically ergodic if and only if its transition kernel admits a spectral gap in L_∞^V . If the chain is reversible, the same equivalence holds with L_2 in place of L_∞^V . In the absence of reversibility it fails: There are (necessarily non-reversible, geometrically ergodic) chains that admit a spectral gap in L_∞^V but not in L_2 . Moreover, if a chain admits a spectral gap in L_2 , then for any $h \in L_2$ there exists a Lyapunov function $V_h \in L_1$ such that V_h dominates h and the chain admits a spectral gap in $L_\infty^{V_h}$. The relationship between the size of the spectral gap in L_∞^V or L_2 , and the rate at which the chain converges to equilibrium is also briefly discussed.

I. Kontoyiannis was supported in part by a Sloan Foundation Research Fellowship, and by a Marie Curie International Outgoing Fellowship, PIOF-GA-2009-235837. S.P. Meyn was supported in part by the National Science Foundation ECS-0523620, and by AFOSR grant FA9550-09-1-0190. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

I. Kontoyiannis (✉)
Department of Informatics, Athens University of Economics and Business,
Patisision 76, Athens 10434, Greece
e-mail: yiannis@aueb.gr

S. P. Meyn
Department of Electrical and Computer Engineering and the Coordinated Sciences Laboratory,
University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA
e-mail: meyn@uiuc.edu

Keywords Markov chain · Geometric ergodicity · Spectral theory · Stochastic Lyapunov function · Reversibility · Spectral gap

Mathematics Subject Classification (2000) 60J05 · 60J10 · 37A30 · 37A25

1 Introduction and main results

There is increasing interest in spectral theory and rates of convergence for Markov chains. Research is motivated by elegant mathematics as well as a range of applications. In particular, one of the most effective general methodologies used to establish bounds on the convergence rate of a geometrically ergodic chain is via an analysis of the spectrum of the chain's transition kernel. See, e.g., [2, 3, 6, 8, 10, 12, 19–21, 23, 25, 26, 29, 31–33], and the relevant references therein.

The word *spectrum* naturally invites techniques grounded in a Hilbert space framework. The majority of quantitative results on rates of convergence are obtained using such methods, within the Hilbert space $L_2 = L_2(\pi)$, where π denotes the stationary distribution of the Markov chain in question. Indeed, most successful studies have been carried out for Markov chains that are reversible, in which case a key to analysis is the fact that the transition kernel, viewed as a linear operator on L_2 , is self-adjoint. See, e.g., [4, 11, 13, 22], as well as the relevant references above. In this paper we argue that, in the *absence* of reversibility, the Hilbert space framework may not be the appropriate setting for spectral analysis.

To be specific, let $X = \{X(n) : n \geq 0\}$ denote a discrete-time Markov chain with values on a general state space \mathbf{X} . We assume that \mathbf{X} is equipped with a countably generated sigma-algebra \mathcal{B} . The distribution of X is described by its initial state $X(0) = x_0 \in \mathbf{X}$ and the transition semigroup $\{P^n : n \geq 0\}$, where, for each n ,

$$P^n(x, A) := \Pr\{X(n) \in A \mid X(0) = x\}, \quad x \in \mathbf{X}, \quad A \in \mathcal{B}.$$

For simplicity we write P for the one-step kernel P^1 . Recall that each P^n , like any (not necessarily probabilistic) kernel $Q(x, dy)$ acts on functions $F : \mathbf{X} \rightarrow \mathbb{C}$ and signed measures ν on $(\mathbf{X}, \mathcal{B})$, via,

$$QF(\cdot) = \int_{\mathbf{X}} Q(\cdot, dy)F(y) \quad \text{and} \quad \nu Q(\cdot) = \int_{\mathbf{X}} \nu(dx)Q(x, \cdot),$$

whenever the integrals exist. Throughout the paper, we assume that the chain $X = \{X(n)\}$ is *ψ -irreducible and aperiodic*; cf. [24, 27]. This means that there is a σ -finite measure ψ on $(\mathbf{X}, \mathcal{B})$ such that, for any $A \in \mathcal{B}$ with $\psi(A) > 0$, and any $x \in \mathbf{X}$,

$$P^n(x, A) > 0, \quad \text{for all } n \text{ sufficiently large.}$$

Moreover, we assume that ψ is *maximal* in the sense that any other such ψ' is absolutely continuous with respect to ψ .

1.1 Geometric ergodicity

The natural class of chains to consider in the present context is that of *geometrically ergodic* chains, namely, chains with the property that there exists an invariant measure π on (X, \mathcal{B}) and functions $\rho : X \rightarrow (0, 1)$ and $C : X \rightarrow [1, \infty)$, such that,

$$\|P^n(x, \cdot) - \pi\|_{TV} \leq C(x)\rho(x)^n, \quad \text{for all } n \geq 0, \pi\text{-a.e. } x \in X,$$

where $\|\mu\|_{TV} := \sup_{A \in \mathcal{B}} |\mu(A)|$ denotes the total variation norm on signed measures. Under ψ -irreducibility and aperiodicity this is equivalent [24,29] to the seemingly stronger requirement that there is a single constant $\rho \in (0, 1)$, a constant $B < \infty$ and a π -a.e. finite function $V : X \rightarrow [1, \infty]$, such that,

$$\|P^n(x, \cdot) - \pi\|_V \leq B V(x)\rho^n, \quad \text{for all } n \geq 0, \pi\text{-a.e. } x \in X, \tag{1}$$

where $\|\mu\|_V := \sup\{|\int F d\mu| : F \in L_\infty^V\}$ denotes the V -norm on signed measures, and where L_∞^V denotes the weighted- L_∞ space consisting of all measurable functions $F : X \rightarrow \mathbb{C}$ with,

$$\|F\|_V := \sup_{x \in X} \frac{|F(x)|}{V(x)} < \infty. \tag{2}$$

Another equivalent and operationally simpler definition of geometric ergodicity for a ψ -irreducible, aperiodic chain $X = \{X(n)\}$, is that it satisfies the following drift criterion [24]:

$$\left. \begin{aligned} &\text{There is a function } V : X \rightarrow [1, \infty], \text{ a small set } C \subset X, \\ &\text{and constants } \delta > 0, b < \infty, \text{ such that:} \\ &PV \leq (1 - \delta)V + b\mathbb{1}_C. \end{aligned} \right\} \tag{V4}$$

We then say that the chain is *geometrically ergodic with Lyapunov function V* . In (V4) it is always assumed that the Lyapunov function V is finite for at least one x (and then it is necessarily finite ψ -a.e.). Also, recall that a set $C \in \mathcal{B}$ is *small* if there exist $n \geq 1, \epsilon > 0$ and a probability measure ν on (X, \mathcal{B}) such that, $P^n(x, A) \geq \epsilon \mathbb{1}_C(x)\nu(A)$, for all $x \in X, A \in \mathcal{B}$.

Our first result relates geometric ergodicity to the spectral properties of the kernel P . Its proof, given at the end of Sect. 3, is based on ideas from [20]. See Sect. 2 for more precise definitions.

Proposition 1.1 *A ψ -irreducible and aperiodic Markov chain $X = \{X(n)\}$ is geometrically ergodic with Lyapunov function V if and only if P admits a spectral gap in L_∞^V .*

1.2 Reversibility

Recall that the chain $X = \{X(n)\}$ is called *reversible* if there is a probability measure π on (X, \mathcal{B}) satisfying the detailed balance conditions,

$$\pi(dx)P(x, dy) = \pi(dy)P(y, dx).$$

This is equivalent to saying that the linear operator P is self-adjoint on the space $L_2 = L_2(\pi)$ of (measurable) functions $F : X \rightarrow \mathbb{C}$ that are square-integrable under π , endowed with the inner product $(F, G) = \int FG^* d\pi$, where ‘*’ denotes the complex conjugate operation.

The following result is the natural analog of Proposition 1.1 for reversible chains. Its proof, given in Sect. 3, is partly based on results in [29].

Proposition 1.2 *A reversible, ψ -irreducible and aperiodic Markov chain $X = \{X(n)\}$ is geometrically ergodic if and only if P admits a spectral gap in L_2 .*

1.3 Spectral theory

The main question addressed in this paper is whether the reversibility assumption of Proposition 1.2 can be relaxed. In other words, whether the space L_2 can be used to characterize geometric ergodicity like L_∞^V was in Proposition 1.1. One direction is true without reversibility: A spectral gap in L_2 implies that the chain is “geometrically ergodic in L_2 ” [29,30], and this implies the existence of a Lyapunov function V satisfying (V4) [30]. Therefore, the chain is geometrically ergodic in the sense of [20], where it is also shown that it must admit a central gap in L_∞^V . A direct, explicit construction of a Lyapunov function V_h is given in our first main result stated next, where quantitative information about V_h is also obtained. It is proved in Sect. 3.

Theorem 1.3 *Suppose that a ψ -irreducible, aperiodic chain $X = \{X(n)\}$ admits a spectral gap in L_2 . Then, for any $h \in L_2$, there is π -integrable function V_h , such that the chain is geometrically ergodic with Lyapunov function V_h and $h \in L_\infty^{V_h}$.*

But the other direction may not hold in the absence of reversibility. Based on earlier counterexamples constructed by Häggström [17, 18] and Bradley [1], in Sect. 3 we prove the following:

Theorem 1.4 *There exists a ψ -irreducible, aperiodic Markov chain $X = \{X(n)\}$ which is geometrically ergodic but does not admit a spectral gap in L_2 .*

The results above, which can be summarized as saying that, “exponentially fast convergence to equilibrium is not equivalent to the existence of a spectral gap in L_2 ,” admit the following operationally positive interpretation: In practice, even in cases where a chain is known to not admit a spectral gap in L_2 , it may still be geometrically ergodic.

Related results on the connections between reversibility, geometric convergence rates, and the spectral gap δ_2 , can be found in the recent work [34,35].

1.4 Convergence rates

The existence of a spectral gap is intimately connected to the exponential convergence rate for a ψ -irreducible, aperiodic Markov chain. For example, if the chain is reversible, we have the following well-known, quantitative bound. See Sect. 2 for detailed definitions; the result follows from the results in [29], combined with Lemma 2.2 given in Sect. 2.

Proposition 1.5 *Suppose that a reversible chain $X = \{X(n)\}$ is ψ -irreducible, aperiodic, and has initial distribution μ . If the chain X admits a spectral gap $\delta_2 > 0$ in L_2 , then,*

$$\|\mu P^n - \pi\|_{TV} \leq \frac{1}{2} \|\mu - \pi\|_2 (1 - \delta_2)^n, \quad n \geq 1,$$

where the L_2 -norm on signed measures ν is defined as the $L_2(\pi)$ -norm of the density $d\nu/d\pi$ if it exists, and is set equal to infinity otherwise.

In the absence of reversibility, the size of the spectral gap in L_∞^V precisely determines the exponential convergence rate of any geometrically ergodic chain. The result of the following proposition is stated in Lemma 2.3 in Sect. 2.

Proposition 1.6 *Suppose that the chain $X = \{X(n)\}$ is ψ -irreducible and aperiodic. If it admits a spectral gap $\delta_V > 0$ in L_∞^V , then, for π -a.e. x ,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \|P^n(x, \cdot) - \pi\|_V = \log(1 - \delta_V).$$

In fact, the convergence is uniform in that:

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\sup_{x \in X, \|F\|_V=1} \frac{|P^n F(x) - \int F d\pi|}{V(x)} \right) = \log(1 - \delta_V). \tag{3}$$

Although Proposition 1.5 gives a finite- n bound for reversible chains, the corresponding result of Proposition 1.6 in the general case is only asymptotic. This is necessarily so, because, in the absence of the Hilbert space setting L_2 , there is no guarantee that the eigenfunctions of the transition kernel will be orthogonal. Indeed, it is possible that the first n , say, eigenfunctions are “almost” aligned, which could induce a polynomial pre-factor in the corresponding bound, the degree of which can be made arbitrarily large by making n arbitrarily large.

We close this introduction with a few more words on this issue of obtaining accurate, non-asymptotic estimates for the mixing time of a non-necessarily-reversible chain, in connection with the introduction of the Banach space L_∞^V for the analysis of such chains. Although, as noted earlier, there are several spectral convergence bounds for reversible chains, these are also known to *not* hold in the absence of reversibility. On the other hand, similar results do exist for non-reversible chains (typically utilizing the spectrum of some reversible “version” of the transition kernel, see, e.g., [2, 15, 16]),

although the resulting bounds can be possibly quite loose. In such cases, knowing the spectral gap $(1 - \delta_2)$, or even knowing the entire L_2 -spectrum explicitly may be of little or even no use; a well-known such example is that of the the move-to-front chain, see, e.g., [7, 14, 28]. In these cases, when precise estimates are sought, it is common that different methods are employed; see, e.g., the analysis of the MCMC sampler in [9].

Section 2 contains precise definitions regarding the spectrum and the spectral gap of the kernel P acting either on L_2 or the weighted- L_∞ space L_∞^V . Simple properties of the spectrum are also stated and proved. Section 3 contains the proofs of the first four results stated above.

2 Spectra and geometric ergodicity

We begin by giving precise definitions for the spectrum and spectral gap of the transition kernel P , viewed as a linear operator. The spectrum depends on the domain of P , for which we consider two possibilities:

- (i) The Hilbert space $L_2 = L_2(\pi)$, equipped with the norm $\|F\|_2 = [\int |F|^2 d\pi]^{1/2}$.
- (ii) The Banach space L_∞^V , with norm $\|\cdot\|_V$ defined in (2).

In either case, the spectrum is defined as the set of nonzero $\lambda \in \mathbb{C}$ for which the inverse $(I\lambda - P)^{-1}$ does not exist as a bounded linear operator on the domain of P . The transition kernel admits a *spectral gap* if there exists $\varepsilon_0 > 0$ such that $\mathcal{S} \cap \{z : |z| \geq 1 - \varepsilon_0\}$ is finite, and contains only poles of finite multiplicity; see [20, Sect. 4] for more details. The spectrum is denoted \mathcal{S}_2 when P is viewed as a linear operator on L_2 , and it is denoted \mathcal{S}_V when P is viewed as a linear operator on L_∞^V .

The induced operator norm of a linear operator $\widehat{P} : L_\infty^V \rightarrow L_\infty^V$ is defined as usual via,

$$\|\widehat{P}\|_V := \sup \frac{\|\widehat{P}F\|_V}{\|F\|_V},$$

where the supremum is over all $F \in L_\infty^V$ satisfying $\|F\|_V \neq 0$. An analogous definition gives the induced operator norm $\|\widehat{P}\|_2$ of a linear operator \widehat{P} acting on L_2 .

For a ψ -irreducible, aperiodic chain $X = \{X(n)\}$, geometric ergodicity expressed in the form (1) implies that P^n converges to a rank-one operator, at a geometric rate: For some constants $B < \infty, \rho \in (0, 1)$,

$$\|P^n - \mathbf{1} \otimes \pi\|_V \leq B \rho^n, \quad n \geq 0, \tag{4}$$

where the outer product $\mathbf{1} \otimes \pi$ denotes the kernel $\mathbf{1} \otimes \pi(x, dy) = \pi(dy)$. It follows that the inverse $[I\lambda - P + \mathbf{1} \otimes \pi]^{-1}$ exists as a bounded linear operator on L_∞^V , whenever $\lambda > \rho$. This in turn implies that P has a single isolated pole at $\lambda = 1$ in the set $\{\lambda \in \mathbb{C} : \lambda > \rho\}$, so that P admits a spectral gap.

In Lemma 2.1 we clarify the location of poles when the chain admits a spectral gap in L_2 or L_∞^V .

Lemma 2.1 *If a ψ -irreducible, aperiodic Markov chain admits a spectral gap in L_∞^V or L_2 , then the only pole on the unit circle in \mathbb{C} is $\lambda = 1$, and this pole has multiplicity one.*

Proof We present the proof for L_∞^V ; the proof in L_2 is identical.

We first note that the existence of a spectral gap implies ergodicity: There is a left eigenmeasure μ corresponding to the eigenvalue 1, satisfying $\mu P = \mu$ and $|\mu|(V) = \|\mu\|_V < \infty$. On letting $\pi(\cdot) = |\mu(\cdot)|/|\mu(X)|$ we conclude that π is super-invariant: $\pi P \geq \pi$. Since $\pi(X) = 1$ we must have invariance. The ergodic theorem for positive recurrent Markov chains implies that $E[G(X(n)) | X(0) = x] \rightarrow \int G d\pi$, as $n \rightarrow \infty$, whenever $G \in L_1(\pi)$.

Ergodicity rules out the existence of multiple eigenfunctions corresponding to $\lambda = 1$. Hence, if this pole has multiplicity greater than one, then there is a generalized eigenfunction $h \in L_\infty^V$ satisfying,

$$Ph = h + 1.$$

Iterating gives $P^n h(x) = E[h(X(n)) | X(0) = x] = h(x) + n$ for $n \geq 1$. This rules out ergodicity, and proves that $\lambda = 1$ has multiplicity one.

We now show that if $\lambda \in \mathcal{S}_V$ with $|\lambda| = 1$, then $\lambda = 1$. To see this, let $h \in L_\infty^V$ denote an eigenfunction, $Ph = \lambda h$. Iterating, we obtain,

$$E[h(X(n)) | X(0) = x] = h(x)\lambda^n.$$

Then, letting $n \rightarrow \infty$, the right-hand-side converges to $\int h d\pi$ for a.e. x ., so that $\lambda = 1$ and $h(x) = \int h d\pi$, π -a.e. □

Therefore, for a ψ -irreducible, aperiodic chain, the existence of a spectral gap in L_2 is equivalent to the existence of a single eigenvalue $\lambda = 1$ on the unit circle, which has multiplicity one. The spectral gap δ_2 is then defined as,

$$\delta_2 = 1 - \sup\{|\lambda| : \lambda \in \mathcal{S}_2, \lambda \neq 1\},$$

and similarly for δ_V .

Next we state two well-known, alternative expressions for the L_2 -spectral gap δ_2 of a reversible chain. See, e.g., [29, Theorem 2.1] and [5, Proposition VIII.1.11].

Lemma 2.2 *Suppose X is a ψ -irreducible, aperiodic, reversible Markov chain. Then, its L_2 -spectral gap δ_2 admits the alternative characterizations,*

$$\begin{aligned} \delta_2 &= 1 - \sup \left\{ \frac{\|vP\|_2}{\|v\|_2} : \text{signed measures } v \text{ with } v(X) = 0, \|v\|_2 \neq 0 \right\} \\ &= 1 - \lim_{n \rightarrow \infty} (\|P^n - \mathbf{1} \otimes \pi\|_2)^{1/n}, \end{aligned}$$

where the limit is the usual spectral radius of the semigroup $\{\widehat{P}^n\}$ generated by the kernel $\widehat{P} = P - \mathbf{1} \otimes \pi$, acting on functions in $L_2(\pi)$.

A similar result holds for δ_V , even in the absence of reversibility; see, e.g., [21].

Lemma 2.3 *Suppose X is a ψ -irreducible, aperiodic Markov chain. Then, its L^∞_V -spectral gap δ_V admits the following alternative characterization in terms of the spectral radius,*

$$\delta_V = 1 - \lim_{n \rightarrow \infty} (\|P^n - \mathbf{1} \otimes \pi\|_V)^{1/n}.$$

3 Proofs

First we prove Theorem 1.3. The following notation will be useful throughout this section.

For a Markov chain $X = \{X(n)\}$, the first hitting time and first return time to a set $C \in \mathcal{B}$ are defined, respectively, by,

$$\begin{aligned} \sigma_C &:= \min\{n \geq 0 : X(n) \in C\}; \\ \tau_C &:= \min\{n \geq 1 : X(n) \in C\}. \end{aligned} \tag{5}$$

Conditional on $X(0) = x$, the expectation operator corresponding to the measure defining the distribution of the process $X = \{X(n)\}$ is denoted $E_x(\cdot)$, so that, for example, $P^n F(x) = E[F(X(n)) | X(0) = x] = E_x[F(X(n))]$. For an arbitrary signed measure μ on (X, \mathcal{B}) , we write $\mu(F)$ for $\int F d\mu$, for any function $F : X \rightarrow \mathbb{C}$ for which the integral exists.

Proof of Theorem 1.3 Since $\pi(h^2) < \infty$, and the chain is ψ -irreducible, it follows that there exists an increasing sequence of h^2 -regular sets providing a π -a.e. covering of X [24, Theorem 14.2.5]. That is, there is a sequence of sets $\{S_r : r \in \mathbb{Z}_+\}$ such that $\pi(S_r) \rightarrow 1$ as $r \rightarrow \infty$, $S_r \subset S_{r+1}$ for each r , and the following bounds hold,

$$\begin{aligned} V_r(x) &:= E_x \left[\sum_{n=0}^{\tau_{S_r}} h^2(X(n)) \right] < \infty, \quad \text{for } \pi\text{-a.e. } x \\ \sup_{x \in S_r} V_r(x) &< \infty. \end{aligned}$$

Since the chain admits a spectral gap in L_2 , combining Theorem 2.1 of [29] with Lemma 2.2 and the results of [30], we have that it is geometrically ergodic. Hence, from [24, Theorem 15.4.2] it follows that there exists a sequence of *Kendall sets* providing a π -a.e. covering of X . That is, there is a sequence of sets $\{K_r : r \in \mathbb{Z}_+\}$ and positive constants $\{\theta_r : r \in \mathbb{Z}_+\}$ satisfying $\pi(K_r) \rightarrow 1$ as $r \rightarrow \infty$, $K_r \subset K_{r+1}$ for each r , and the following bounds hold,

$$\begin{aligned} U_r(x) &:= E_x [\exp(\theta_r \tau_{K_r})] < \infty, \quad \text{for } \pi\text{-a.e. } x \\ \sup_{x \in K_r} U_r(x) &< \infty. \end{aligned}$$

We also define another collection of sets,

$$C_{r,m} := \{x \in X : U_r(x) + V_r(x) \leq m\}.$$

For each $r \geq 1$, these sets are non-decreasing in m , and $\pi(C_{r,m}) \rightarrow 1$ as $m \rightarrow \infty$. Moreover, whenever $C_{r,m} \in \mathcal{B}^+$, this set is both an h^2 -regular set and a Kendall set. This follows by combining Theorems 14.2.1 and 15.2.1 of [24]. Fix r_0 and m_0 so that $\pi(C_{r_0,m_0}) > 0$. We henceforth denote C_{r_0,m_0} by C , and let $\theta > 0$ denote a value satisfying the bound,

$$\mathbb{E}_x [\exp(\theta \tau_C)] < \infty, \quad \text{for } \pi\text{-a.e. } x,$$

where the expectation is uniformly bounded over the Kendall set C .

The candidate Lyapunov function can now be defined as,

$$V_h(x) := \mathbb{E}_x \left[\sum_{n=0}^{\sigma_C} (1 + |h(X(n))|) \exp\left(\frac{1}{2}\theta n\right) \right]. \tag{6}$$

We first obtain a bound on this function. Writing,

$$V_h(x) = \mathbb{E}_x \left[\sum_{n=0}^{\sigma_C} \exp\left(\frac{1}{2}\theta n\right) \right] + \sum_{n=0}^{\infty} \mathbb{E}_x [|h(X(n))| \exp\left(\frac{1}{2}\theta n\right) \mathbb{I}\{n \leq \sigma_C\}],$$

we see that the first term is finite π -a.e. by construction. The square of the second term is bounded above, using the Cauchy-Schwartz inequality, by,

$$\begin{aligned} & \mathbb{E}_x \left[\sum_{n=0}^{\infty} |h(X(n))|^2 \mathbb{I}\{n \leq \sigma_C\} \right] \mathbb{E}_x \left[\sum_{n=0}^{\infty} \exp(\theta n) \mathbb{I}\{n \leq \sigma_C\} \right] \\ &= U_{r_0}(x) \mathbb{E}_x \left[\sum_{t=0}^{\sigma_C} \exp(\theta t) \right], \end{aligned}$$

so that V_h is finite π -a.e., and we also easily see that $|h| \leq V_h$ so that $h \in L^\infty_{V_h}$.

Next we show that V_h satisfies (V4): First apply $e^{\frac{1}{2}\theta} P$ to the function V_h to obtain,

$$e^{\frac{1}{2}\theta} P V_h(x) = \mathbb{E}_x \left[\sum_{n=1}^{\tau_C} (1 + |h(X(n+1))|) \exp\left(\frac{1}{2}\theta(t+1)\right) \right] \tag{7}$$

We have $\tau_C = \sigma_C$ when $X(0) \in C^c$. This gives,

$$e^{\frac{1}{2}\theta} P V_h(x) = V_h(x) - (1 + |h(x)|), \quad x \in C^c.$$

If $X(0) = x \in C$, then the previous arguments imply that the right-hand-side of (7) is finite, and in fact uniformly bounded over $x \in C$. Combining these results, we conclude that there exists a constant b_0 such that,

$$PV_h \leq e^{-\frac{1}{2}\theta} V_h + b_0 \mathbb{1}_C$$

Regular sets are necessarily small [24, Theorem 11.3.11] so that this is a version of the drift inequality (V4).

Finally note that, by the fact that (V4) implies the weaker drift condition (V3) of [24], the function V_h is π -integrable by [24, Theorem 14.0.1]. \square

Theorem 1.3 states that (V4) holds for a Lyapunov function V_h with $h \in L_\infty^{V_h}$. If this could be strengthened to show that for every geometrically ergodic chain and any $h \in L_2$, the chain was geometrically ergodic with a Lyapunov function V_h that had $h^2 \in L_\infty^{V_h}$, then the central limit theorem would hold for the partial sums of $h(X(n))$ [24, Theorem 17.0.1]. But this is not generally possible:

Proposition 3.1 *There exists a geometrically ergodic Markov chain on a countable state space X and a function $G \in L_2$ with mean $\pi(G) = 0$, for which the central limit theorem fails in that the normalized partial sums,*

$$\frac{1}{\sqrt{n}} \sum_{i=0}^{n-1} G(X(i)), \quad n \geq 1, \tag{8}$$

converge neither to a normal distribution nor to a point mass.

The result of the proposition appears in [17, Theorem 1.3], and an earlier counterexample in [1] yields the same conclusion. Based on these counterexamples we now show that geometric ergodicity does not imply a spectral gap in the Hilbert space setting.

Proof of Theorem 1.4 Suppose that the Markov chain $X = \{X(n)\}$ constructed in Proposition 3.1 does admit a spectral gap in L_2 . Then its autocorrelation function decays geometrically fast, for any $h \in L_2$: Assuming without loss of generality that $\pi(h) = 0$, and letting $R_h(n) = \pi(hP^n h)$, for all n , we have the bound,

$$|R(n)| \leq \sqrt{\pi(h^2)\pi((P^n h)^2)}, \quad n \geq 1.$$

Applying Theorem 1.3, we conclude that the right-hand-side decays geometrically fast as $n \rightarrow \infty$. Consequently, the sequence of normalized sums,

$$S_n := \frac{1}{\sqrt{n}} \sum_{i=0}^{n-1} h(X(i)), \quad n \geq 1,$$

is uniformly bounded in L_2 , i.e.,

$$\limsup_{n \rightarrow \infty} \mathbf{E}_\pi[S_n^2] \leq \sum_{n=-\infty}^{\infty} |R(n)|,$$

where $\mathbf{E}_\pi[\cdot]$ denotes the expectation operator corresponding to the stationary version of the chain. However, this is impossible for the choice of the function $h = G$ as in Proposition 3.1: In [17, p. 81] it is shown that the corresponding normalized sums in (8) fail to define a tight sequence of probability distributions. This is a consequence of [17, Lemma 3.2].

This contradiction establishes the claim that the Markov chain of Proposition 3.1 cannot admit a spectral gap in L_2 . □

Finally we prove Propositions 1.1 and 1.2.

Proof of Proposition 1.1 The equivalence stated in the proposition is obtained on combining Lemma 2.1 with [20, Proposition 4.6]. To explain this, we introduce new terminology: The transition kernel is called *V-uniform* if $\lambda = 1$ is the only pole on the unit circle in \mathbb{C} , and this pole has multiplicity one. Proposition 4.6 of [20] states that geometric ergodicity with respect to a Lyapunov function V is equivalent to V -uniformity of the kernel P . Consequently, the direct part of the proposition holds, since V -uniformity of P implies that it admits a spectral gap in L_∞^V .

Conversely, if the chain admits a spectral gap in L_∞^V , then Lemma 2.1 states that P is V -uniform. Applying Proposition 4.6 of [20] once more, we conclude that the chain is geometrically ergodic with the same Lyapunov function V . □

Proof of Proposition 1.2 The forward direction of the statement of the proposition is contained in [29] and [30].

The converse again follows from Lemma 2.1 and a minor modification of the arguments used in [20, Proposition 4.6]. If the chain admits a spectral gap in L_2 , then the lemma states that $\lambda = 1$ has multiplicity one, and that this is the only pole on the unit circle in \mathbb{C} . It follows that for some $\rho < 1$, the inverse $[zI - (P - \mathbf{1} \otimes \pi)]^{-1}$ exists as a bounded linear operator on L_2 , whenever $|z| \geq \rho$. Denote $b_\rho = \sup \{ \| [zI - (P - \mathbf{1} \otimes \pi)]^{-1} \|_2 : |z| = \rho \}$, where $\| \cdot \|_2$ is the induced operator norm on L_2 .

Following the proof of [20, Theorem 4.1], we conclude that finiteness of b_ρ implies a form of geometric ergodicity: For any $g \in L_2$,

$$\frac{1}{2\pi} \int_0^{2\pi} e^{in\phi} [\rho e^{in\phi} I - (P - \mathbf{1} \otimes \pi)]^{-1} g = \rho^{-n-1} (P^n g - \pi(g)).$$

Therefore, the L_2 -norm of the left-hand-side is bounded by $b_\rho \|g\|_2$. This gives,

$$\|P^n g - \pi(g)\|_2 \leq b_\rho \|g\|_2 \rho^{n+1}, \quad n \geq 1.$$

It follows from [24, Theorem 15.4.3] that the Markov chain is geometrically ergodic. □

Acknowledgments We wish to thank David Aldous, Yuval Peres, Jeff Rosenthal and David Wilson for their useful bibliographical comments.

References

1. Bradley, R.C. Jr.: Information regularity and the central limit question. *Rocky Mountain J. Math.* **13**(1), 77–97 (1983)
2. Brémaud, P.: Markov chains: Gibbs fields, Monte Carlo simulation, and queues, volume 31 of Texts in Applied Mathematics. Springer-Verlag, New York (1999)
3. Brooks, S.P., Roberts, G.O.: Convergence assessment techniques for Markov chain Monte Carlo. *Statistics and Computing* **8**, 319–335 (1998)
4. Chen, Mu-Fa.: Eigenvalues, inequalities, and ergodic theory, Probability and its Applications (New York). Springer-Verlag London Ltd., London (2005)
5. Conway, J.B.: A course in functional analysis 2nd edn. Springer-Verlag, New York (1990)
6. Dellnitz, M., Junge, O.: On the approximation of complicated dynamical behavior. *SIAM J. Numer. Anal.* **36**(2), 491–515 (1999)
7. Diaconis, P.: Mathematical developments from the analysis of riffle shuffling. In: Groups, combinatorics & geometry (Durham, 2001), pp. 73–97. World Sci. Publ., River Edge NJ, (2003)
8. Diaconis, P.: The Markov chain Monte Carlo revolution. *Bull. Amer. Math. Soc. (N.S.)* **46**(2), 179–205 (2009)
9. Diaconis, P., Holmes, S., Neal, R.M.: Analysis of a nonreversible Markov chain sampler. *Ann. Appl. Probab.* **10**(3), 726–752 (2000)
10. Diaconis, P., Khare, K., Saloff-Coste, L.: Gibbs sampling, exponential families and orthogonal polynomials. *Statist. Sci.*, **23**(2), pp. 151–200 (2008). (With comments and a rejoinder by the authors)
11. Diaconis, P., Saloff-Coste, L.: Comparison theorems for reversible Markov chains. *Ann. Appl. Probab.* **3**(3), 696–730 (1993)
12. Diaconis, P., Saloff-Coste, L.: What do we know about the Metropolis algorithm? *J. Comput. System Sci.*, **57**(1), pp. 20–36 (1998). 27th Annual ACM Symposium on the Theory of Computing (STOC'95) (Las Vegas, NV)
13. Diaconis, P., Stroock, D.: Geometric bounds for eigenvalues of Markov chains. *Ann. Appl. Probab.* **1**(1), 36–61 (1991)
14. Dobrow, R.P., Fill, J.A.: The move-to-front rule for self-organizing lists with Markov dependent requests. In: Discrete probability and algorithms (Minneapolis, MN, 1993), volume 72 of IMA Vol. Math. Appl., pp. 57–80. Springer, New York, (1995)
15. Evans, W., Kenyon, C., Peres, Y., Schulman, L.J.: Broadcasting on trees and the Ising model. *Ann. Appl. Probab.* **10**(2), 410–433 (2000)
16. Fill, J.A.: Eigenvalue bounds on convergence to stationarity for nonreversible Markov chains, with an application to the exclusion process. *Ann. Appl. Probab.* **1**(1), 62–87 (1991)
17. Häggström, O.: On the central limit theorem for geometrically ergodic Markov chains. *Probab. Theory Relat. Fields* **132**(1), 74–82 (2005)
18. Häggström, O.: Acknowledgement of priority concerning “On the central limit theorem for geometrically ergodic Markov chains” [*Probab. Theory Relat. Fields* **132**(1), 74–82 (2005)]. *Probab. Theory Relat. Fields* **135**(3):470 (2006)
19. Huisinga, W., Meyn, S.P., Schutte, C.: Phase transitions and metastability in Markovian and molecular systems. *Ann. Appl. Probab.* (2001). (in press)
20. Kontoyiannis, I., Meyn, S.P.: Spectral theory and limit theorems for geometrically ergodic Markov processes. *Ann. Appl. Probab.* **13**, 304–362 (2003)
21. Kontoyiannis, I., Meyn, S.P.: Large deviation asymptotics and the spectral theory of multiplicatively regular Markov processes. *Electron. J. Probab.* **10**(3), 61–123 (2005)
22. Lawler, G.F., Sokal, A.D.: Bounds on the L^2 spectrum for Markov chains and Markov processes: a generalization of Cheeger’s inequality. *Trans. Amer. Math. Soc.* **309**(2), 557–580 (1988)
23. Lebeau, G., Diaconis, P.: Métropolis: le jour où l’étoile probabilité entra dans le champ gravitationnel de la galaxie microlocale. In: Séminaire: Équations aux Dérivées Partielles. 2006–2007, Sémin. Équ. Dériv. Partielles, pages Exp. No. XIV, 13. École Polytech., Palaiseau (2007)

24. Meyn, S.P., Tweedie, R.L.: Markov Chains and Stochastic Stability 2nd edn. Cambridge University Press, London (2009). Published in the Cambridge Mathematical Library. 1993 edition online: <http://black.csl.uiuc.edu/~meyn/pages/book.html>
25. Meyn, S.P., Tweedie, R.L.: Computable bounds for geometric convergence rates of Markov chains. *Ann. Appl. Probab.* **4**(4), 981–1011 (1994)
26. Montenegro, R., Tetali, P.: Mathematical aspects of mixing times in Markov chains. *Found. Trends Theor. Comput. Sci.* **1**(3), 1–121 (2006)
27. Nummelin, E.: General Irreducible Markov Chains and Nonnegative Operators. Cambridge University Press, Cambridge (1984)
28. Phatarfod, R.M., Pryde, A.J., Dyte, D.: On the move-to-front scheme with Markov dependent requests. *J. Appl. Probab.* **34**(3), 790–794 (1997)
29. Roberts, G.O., Rosenthal, J.S.: Geometric ergodicity and hybrid Markov chains. *Electron. Comm. Probab.* **2**(2), 13–25 (electronic), (1997)
30. Roberts, G.O., Tweedie, R.L.: Geometric L^2 and L^1 convergence are equivalent for reversible Markov chains. *J. Appl. Probab.* **38A**, 37–41 (2001). Probability, statistics and seismology
31. Rosenthal, J.S.: Correction: Minorization conditions and convergence rates for Markov chain Monte Carlo. *J. Amer. Statist. Assoc.* **90**(431), 1136 (1995)
32. Rosenthal, J.S.: Minorization conditions and convergence rates for Markov chain Monte Carlo. *J. Amer. Statist. Assoc.* **90**(430), 558–566 (1995)
33. Winkler, G.: Image Analysis, Random Fields and Dynamic Monte Carlo Methods: A Mathematical Introduction. Springer-Verlag, Berlin (1995)
34. Wübker, A.: L^2 -spectral gaps for time discrete reversible Markov chains. Preprint (2009)
35. Wübker, A., Kabluchko, Z.: L^2 -spectral gaps, weak-reversible and very weak-reversible Markov chains. Preprint (2009)