


Foliar $\delta^{15}\text{N}$ values characterize soil N cycling and reflect nitrate or ammonium preference of plants along a temperate grassland gradient

Ansgar Kahmen · Wolfgang Wanek · Nina Buchmann

Published online: 3 September 2008
© Springer-Verlag 2008

Erratum to: Oecologia (2008) 156:861–870
DOI 10.1007/s00442-008-1028-8

Unfortunately, the publisher printed Fig. 2 with errors. The correct Fig. 2 is given here.



The online version of the original article can be found under doi:10.1007/s00442-008-1028-8.

A. Kahmen · N. Buchmann
Max Planck Institute for Biogeochemistry,
Hans-Knöll-Strasse 10, 07745 Jena, Germany

W. Wanek
Department of Chemical Ecology and Ecosystem Research,
University of Vienna, Althanstrasse 14, 1090 Vienna, Austria

A. Kahmen (✉)
Center for Stable Isotope Biogeochemistry,
Department of Integrative Biology, University of California,
Berkeley, CA 94720-3140, USA
e-mail: akahmen@berkeley.edu;
ansgar.kahmen@ipw.agrl.ethz.ch

Present Address:

N. Buchmann
Institute of Plant Sciences, ETH Zurich,
Universitätsstrasse 2, 8092 Zurich, Switzerland

Fig. 2 Simple linear regression analyses between $\delta^{15}\text{N}$ of bulk soil and leaf $\delta^{15}\text{N}$ for different forb and grass species as well as the relationship between daily net mineralization (N_{\min}) and foliar $\Delta\delta^{15}\text{N}$ ($\delta^{15}\text{N}$ plant – $\delta^{15}\text{N}$ soil) for different forb and grass species in 18 temperate grasslands. d^{-1} Day $^{-1}$