

Erratum to: Absence of snow cover reduces understory plant cover and alters plant community composition in boreal forests

Juergen Kreyling · Mahsa Haei · Hjalmar Laudon

Published online: 21 April 2013
© Springer-Verlag Berlin Heidelberg 2013

Erratum to: Oecologia (2012) 168:577–587
DOI 10.1007/s00442-011-2092-z

Unfortunately, Table 3 caption was incorrectly published in the original article. The correct caption with table is given below:

Table 3 Fine root biomass (mean \pm SE in mg cm^{-3}) at 20 cm soil depth increments as influenced by the snow manipulations

| | 0–20 cm | 20–40 cm | 40–60 cm | 0–60 cm |
|--------------|--------------------|-----------------|-----------------|--------------------|
| Snow removal | 2.82 ± 0.34 a | 0.55 ± 0.38 | 0.17 ± 0.08 | 3.53 ± 0.80 a |
| Control | 4.58 ± 0.79 ab | 0.54 ± 0.17 | 0.36 ± 0.03 | 5.49 ± 0.54 ab |
| Reference | 6.31 ± 1.12 b | 0.85 ± 0.27 | 0.37 ± 0.16 | 7.52 ± 1.39 b |
| Insulation | 4.85 ± 0.79 ab | 0.44 ± 0.22 | 0.30 ± 0.13 | 5.60 ± 0.78 ab |

ANOVA: treatment: $p = 0.021$; depth: $p < 0.001$; interaction treatment:depth: $p = 0.041$. Letters in columns indicate homogenous groups according to Tukey post-hoc comparisons per depth layer if significant pairwise comparisons were found, $n = 3$ per cell

The online version of the original article can be found under doi:[10.1007/s00442-011-2092-z](https://doi.org/10.1007/s00442-011-2092-z).

J. Kreyling (✉)
Biogeography, University of Bayreuth, Bayreuth, Germany
e-mail: juergen.kreyling@uni-bayreuth.de

M. Haei
Department of Ecology and Environmental Science,
Umeå University, Umeå, Sweden

H. Laudon
Department of Forest Ecology and Management,
Swedish University of Agricultural Sciences, Umeå, Sweden