



Correction: Seed dispersal function of the brown bear *Ursus arctos* on Hokkaido Island in northern Japan: gut passage time, dispersal distance, germination, and effects of remaining pulp

Yoshihiro Tsunamoto¹ · Hifumi Tsuruga¹ · Konomi Kobayashi² · Takeshi Sukegawa² · Takuya Asakura²

Published online: 17 May 2024

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2024

Correction: Oecologia (2024) 204:505–515

<https://doi.org/10.1007/s00442-024-05510-5>

In the sentence beginning ‘Each plant’s average simulated seed dispersal distance was 202–512 m’ of the abstract of this article, the seed dispersal distance value was displayed incorrectly as ‘202–512 m’. Instead it should have been ‘181–345 m’.

The original article can be found online at <https://doi.org/10.1007/s00442-024-05510-5>.

✉ Yoshihiro Tsunamoto
tsunamoto-yoshihiro@hro.or.jp

¹ Hokkaido Research Organization, Research Institute of Energy, Environment and Geology, Kita 19 Nishi 12, Kita-ku, Sapporo, Hokkaido 060-0819, Japan

² Sapporo Maruyama Zoo, 3-1 Miyagaoka, Chuo-ku, Sapporo, Hokkaido, Japan