



# Correction to: The influence of cross-sectional channel geometry on rheology and flux estimates for active lava flows

Einat Lev<sup>1</sup> · Mike R. James<sup>2</sup>

Published online: 4 February 2020

© International Association of Volcanology & Chemistry of the Earth's Interior 2020

**Correction to: Bull Volcanol (2014) 76:829**

<https://doi.org/10.1007/s00445-014-0829-3>

The original version of this article unfortunately contained a mistake. The text accompanying Eq. 3 in the published paper should be corrected.

Currently:

$$\gamma = 1 - \frac{384}{\pi^5} \sum_{n=1,3,5,\dots}^{\infty} \frac{1}{n^5} \tanh \frac{2an\pi}{4h}$$

Should be:

$$\gamma = 1 - \frac{384}{\pi^5} \sum_{n=1,3,5,\dots}^{\infty} \frac{1}{n^5} \frac{h}{2a} \tanh \frac{2an\pi}{4h}$$

---

The online version of the original article can be found at <https://doi.org/10.1007/s00445-014-0829-3>

---

✉ Einat Lev  
[einatlev@ldeo.columbia.edu](mailto:einatlev@ldeo.columbia.edu)

<sup>1</sup> Lamont-Doherty Earth Observatory, Columbia University, New York, NY, USA

<sup>2</sup> Lancaster Environment Centre, Lancaster University, Lancaster, UK