

The Sylvester–Chvátal Theorem*

Xiaomin Chen

Department of Computer Science, Rutgers University,
110 Frelinghuysen Road, Piscataway, NJ 08854-8019, USA
xiaomin@paul.rutgers.edu

Abstract. The Sylvester–Gallai theorem asserts that every finite set S of points in two-dimensional Euclidean space includes two points, a and b , such that either there is no other point in S on the line ab , or the line ab contains all the points in S . Chvátal extended the notion of *lines* to arbitrary metric spaces and made a conjecture that generalizes the Sylvester–Gallai theorem. In the present article we prove this conjecture.

1. Introduction

The following celebrated geometric problem was proposed by Sylvester in 1893 [12].

Prove that it is not possible to arrange any finite number of real points so that a right line through every two of them shall pass through a third, unless they all lie in the same right line.

In the twentieth century several proofs were found; the first one, published some 50 years after the problem was proposed and 1 year after Erdős revived it, was due to Gallai [9]. A very short proof was given by Kelly; this proof uses the notion of Euclidean distance and can be found in [4], Section 4.7 of [5], and Chapter 8 of [8]. Additional information on the Sylvester–Gallai theorem can be found in [1], [3], [7], and [11].

In an arbitrary metric space (V, ρ) , Menger [10] defined a ternary relation $\mathcal{B}(\rho)$ of *metric betweenness* on V by

$$(u, v, w) \in \mathcal{B} \Leftrightarrow u, v, w \text{ are all distinct and } \rho(u, w) = \rho(u, v) + \rho(v, w). \quad (1)$$

* This research was supported by (theory of computation) NSF Grant 0105692.

We follow the tradition—adopted by Menger [10]—of writing $[uvw]$ for $(u, v, w) \in \mathcal{B}$. As Chvátal pointed out in [3], if we define the line ab as

$$\{x : [xab]\} \cup \{a\} \cup \{x : [axb]\} \cup \{b\} \cup \{x : [abx]\}, \quad (2)$$

then a straightforward generalization of the Sylvester–Gallai theorem in arbitrary finite metric space is no longer true. Chvátal’s [3] definition of a *line* in an arbitrary metric space is a recursive closure of (2); details are as follows.

First, the ternary relation $\mathcal{B}(\rho)$ in (1) is transformed into a set $\mathcal{H}(\mathcal{B}(\rho))$ of three-point subsets of V :

$$\mathcal{H}(\mathcal{B}(\rho)) = \{(a, b, c) : (a, b, c) \in \mathcal{B}(\rho)\}. \quad (3)$$

Then a family $\mathcal{A}(\mathcal{H}(\mathcal{B}(\rho)))$ of subsets of V is defined by

$$\mathcal{A}(\mathcal{H}(\mathcal{B}(\rho))) = \{S \subseteq V : \text{no } T \text{ in } \mathcal{H}(\mathcal{B}(\rho)) \text{ has } |T \cap S| = 2\}. \quad (4)$$

(It is clear that $\mathcal{A}(\mathcal{H}(\mathcal{B}(\rho)))$ is closed under arbitrary intersection.) For any subset W of V , define $\mathcal{C}(W)$ as

$$\mathcal{C}(W) = \bigcap_{W \subseteq S \in \mathcal{A}} S. \quad (5)$$

Equivalently, if W is finite, then $\mathcal{C}(W)$ is the return value of the program

```

S = W ;
while some T in H(B(rho)) has |T ∩ S| = 2 do S = S ∪ T end ;
return S ;

```

Finally, for any two distinct points a and b of V , the *line* ab is defined as $\mathcal{C}(\{a, b\})$. Note that, according to the definition of a line, there may be more than one line containing two given points; the definition coincides the usual definition of lines in Euclidean spaces. With this definition, Chvátal made the following conjecture in [3], which is made into a theorem by the present paper.

Theorem 1. *If (V, ρ) is a metric space such that $1 < |V| < \infty$, then V contains two distinct points a and b such that the line ab is $\{a, b\}$ or V .*

The original Sylvester–Gallai Theorem is a special case of Theorem 1, where (V, ρ) is a finite subspace of the Euclidean plane.

For a finite, undirected, connected graph $G = (V, E, w)$ with positive weights, there is naturally an induced metric space (V, ρ) , where $\rho(x, y)$ is defined to be the distance between x and y in G . Note that every finite metric space is induced by such a graph. So it suffices to prove the conjecture for finite metric spaces induced by graphs. Although the proof does not need the notations in graphs, this was the setting where we worked on the conjecture.

Chvátal [3] proved this conjecture for all metric spaces with at most nine points and for all (finite) metric spaces induced by connected graphs with unit weights. Chen [2] proved the conjecture for all (finite) subspaces of ℓ_1^2 , the two-dimensional space with

the ℓ_1 -metric. In prior work we developed techniques that allowed us to give a much simpler proof for the ℓ_1^2 case, as well as proofs for metric spaces induced by graphs with some special weights. Based on similar techniques, just one week before we found the proof, Bin Tian proved the conjecture for all finite subspaces of ℓ_1^3 by using a computer program. Finally, our proof here was built based on these techniques and some other crucial observations.

2. The Sylvester–Chvátal Theorem

We reserve the letter V for the ground-set of a finite metric space with at least two points and we reserve the letter ρ for the metric of this space. Our proof of Theorem 1 splits into two parts.

Theorem 2. *If every three points of V are contained in some line, then some line consists of all points of V .*

Theorem 3. *If some three points of V are contained in no line, then some line consists of precisely two points.*

Proof of Theorem 2. Consider a line, L , which is maximal with respect to set-inclusion; we claim that $L = V$. To justify this claim, assume to the contrary that: some point, c , of V lies outside L . Line L is generated by two points, a and b , of V . By assumption, a, b, c are contained in some line; this line contains $L \cup \{c\}$, contradicting the maximality of L .

Proof of Theorem 3. By a *simple edge*, we mean any edge— ab —of the complete graph with vertex-set V such that no point x of V satisfies $[axb]$. By a *simple triangle*, we mean any three points— a, b, c —of V such that all of ab, bc, ca are simple edges. Now consider the following statements:

- (i) some three points of V are contained in no line,
- (ii) some simple triangle is contained in no line,
- (iii) some line consists of precisely two points.

(i) \Rightarrow (ii) By (i), there are three points— a, b, c —of V such that

$$\text{no line contains } \{a, b, c\}; \tag{6}$$

among all such triples, choose one that minimizes $\rho(a, b) + \rho(b, c) + \rho(a, c)$; we claim that a, b, c is a simple triangle.

To justify this claim, assume to the contrary that, without loss of generality, there is a point d such that $[adb]$. Note first that $d \neq c$ (else (6) is contradicted by $[acb]$) and then that $\rho(d, c) < \rho(d, b) + \rho(b, c)$ (else (6) is contradicted by $[dbc]$ and $[adb]$). It follows that

$$\begin{aligned} \rho(a, d) + \rho(d, c) + \rho(a, c) &< \rho(a, d) + \rho(d, b) + \rho(b, c) + \rho(a, c) \\ &= \rho(a, b) + \rho(b, c) + \rho(a, c); \end{aligned}$$




Fig. 1. If line $L(a, c)$ consists of at least three points, the point d will cause infinite trouble. This point guarantees the existence of special paths. As we shall prove, we can always find shorter and shorter special paths.

now minimality of a, b, c implies that some line contains $\{a, d, c\}$; by $[adb]$, the same line contains $\{a, b, c\}$, contradicting (6).

(ii) \Rightarrow (iii) For each ordered triple u, v, w of points of V , we write

$$\Delta(u, v, w) = \rho(u, v) + \rho(v, w) - \rho(u, w).$$

By (ii), some simple triangle— a, b, c —satisfies (6); among all such simple triangles, choose one that minimizes $\Delta(a, b, c)$; we claim that line $L(a, c)$ consists of precisely two points.

To justify this claim, assume to the contrary that line $L(a, c)$ consists of at least three points. This means that some point d satisfies $[dac]$ or $[adc]$ or $[acd]$; since ac is a simple edge, $[adc]$ is excluded; now symmetry allows us to assume $[acd]$. Among all such points d , we choose one that minimizes $\rho(c, d)$; this property of d guarantees that cd is a simple edge (Fig. 1).

Let us show that

$$bd \text{ is not a simple edge.} \quad (7)$$

If (7) is false, then (b, c, d) is a simple triangle; $[acd]$ and (6) guarantee that this simple triangle is contained in no line. It follows that $\Delta(b, c, d) \geq \Delta(a, b, c)$, which means $\rho(c, d) - \rho(b, d) \geq \rho(a, b) - \rho(a, c)$; since $[acd]$, we conclude $[abd]$; but then (6) is contradicted.

In addition, observe that

$$\rho(a, b) + \rho(b, d) < \rho(a, d) + \Delta(a, b, c); \quad (8)$$

if (8) is false, then $[acd]$ guarantees $\rho(b, d) \geq \rho(b, c) + \rho(c, d)$, and so $[bcd]$; but then (6) is contradicted.

By a *path*, we mean any sequence— (a_1, a_2, \dots, a_k) —of points of V ; we define its *length* as

$$\sum_{i=1}^{k-1} \rho(a_i, a_{i+1});$$

if the path is denoted P , then we denote its length $\ell(P)$. By a *special path*, we mean a path (a_1, a_2, \dots, a_k) such that $a_1 = a, k \geq 3, a_k = d$,

- (α) no line contains $\{a_1, a_2, a_3\}$, and
- (β) at least one of a_1a_2 and a_2a_3 is not a simple edge.




Fig. 2. Three cases of the special path P .

Note that (a, b, d) is a special path: here, (α) follows from $[acd]$ with (6) and (β) follows from (7). Now we can choose a shortest special path, (a_1, a_2, \dots, a_k) ; we denote it P . Since (a, b, d) is a special path, (8) guarantees that

$$\ell(P) < \rho(a, d) + \Delta(a, b, c). \tag{9}$$

To complete the proof of (ii) \Rightarrow (iii), we distinguish between three cases (Fig. 2).

Case 1: Neither a_1a_2 nor a_2a_3 is a simple edge. By assumption of this case, there is a point b_{12} such that $[a_1b_{12}a_2]$. Among all such points b_{12} , we choose one that minimizes $\rho(b_{12}, a_2)$; this property of b_{12} guarantees that $b_{12}a_2$ is a simple edge. Similarly, there is a point b_{23} such that $[a_2b_{23}a_3]$ and such that a_2b_{23} is a simple edge. Note that

$$\text{no line contains } \{b_{12}, a_2, b_{23}\}; \tag{10}$$

else $[a_1b_{12}a_2]$, $[a_2b_{23}a_3]$ would guarantee that the same line contains $\{a_1, a_2, a_3\}$, contradicting property (α) of P . Let P' denote the path $(a_1, b_{12}, b_{23}, a_3, \dots, a_k)$. From $[a_1b_{12}a_2]$ and $[a_2b_{23}a_3]$, we have

$$\ell(P) - \ell(P') = \Delta(b_{12}, a_2, b_{23});$$

(10) guarantees that $\Delta(b_{12}, a_2, b_{23}) > 0$, and so P' is shorter than P ; now the minimality of P implies that P' is not special. Since no line contains $\{a_1, b_{12}, b_{23}\}$ (else $[a_1b_{12}a_2]$ would guarantee that the same line contains $\{b_{12}, a_2, b_{23}\}$, contradicting (10)) and yet P' is not special, both a_1b_{12} and $b_{12}b_{23}$ are simple edges. Since $b_{12}b_{23}$ is a simple edge, b_{12}, a_2, b_{23} is a simple triangle, and so (10) implies $\Delta(b_{12}, a_2, b_{23}) \geq \Delta(a, b, c)$. However, then

$$\ell(P) = \ell(P') + \Delta(b_{12}, a_2, b_{23}) \geq \rho(a, d) + \Delta(a, b, c),$$

contradicting (9).

Case 2: a_1a_2 is a simple edge and a_2a_3 is not. As in Case 1, there is a point b_{23} such that $[a_2b_{23}a_3]$ and such that a_2b_{23} is a simple edge. Note that

$$\text{no line contains } \{a_1, a_2, b_{23}\}; \tag{11}$$

else $[a_2b_{23}a_3]$ would guarantee that the same line contains $\{a_1, a_2, a_3\}$, contradicting property (α) of P . Let P' denote the path $(a_1, b_{23}, a_3, \dots, a_k)$. From $[a_2b_{23}a_3]$, we have

$$\ell(P) - \ell(P') = \Delta(a_1, a_2, b_{23});$$

(11) guarantees that $\Delta(a_1, a_2, b_{23}) > 0$, and so P' is shorter than P ; now minimality of P implies that P' is not special. Since no line contains $\{a_1, b_{23}, a_3\}$ (else $[a_2b_{23}a_3]$ would guarantee that the same line contains $\{a_1, a_2, b_{23}\}$, contradicting (11)) and yet P' is not special, both a_1b_{23} and $b_{23}a_3$ are simple edges. Since a_1b_{23} is a simple edge, a_1, a_2, b_{23} is a simple triangle, and so (11) implies $\Delta(a_1, a_2, b_{23}) \geq \Delta(a, b, c)$. However, then

$$\ell(P) = \ell(P') + \Delta(a_1, a_2, b_{23}) \geq \rho(a, d) + \Delta(a, b, c),$$

contradicting (9).

Case 3: a_2a_3 is a simple edge and a_1a_2 is not. As in Case 1, there is a point b_{12} such that $[a_1b_{12}a_2]$ and such that $b_{12}a_2$ is a simple edge. Note that

$$\text{no line contains } \{b_{12}, a_2, a_3\}; \quad (12)$$

else $[a_1b_{12}a_2]$ would guarantee that the same line contains $\{a_1, a_2, a_3\}$, contradicting property (α) of P . Let P' denote the path $(a_1, b_{12}, a_3, \dots, a_k)$. From $[a_1b_{12}a_2]$, we have

$$\ell(P) - \ell(P') = \Delta(b_{12}, a_2, a_3);$$

(12) guarantees that $\Delta(b_{12}, a_2, a_3) > 0$, and so P' is shorter than P ; now minimality of P implies that P' is not special. Since no line contains $\{a_1, b_{12}, a_3\}$ (else $[a_1b_{12}a_2]$ would guarantee that the same line contains $\{b_{12}, a_2, a_3\}$, contradicting (12)) and yet P' is not special, both a_1b_{12} and $b_{12}a_3$ are simple edges. Since $b_{12}a_3$ is a simple edge, b_{12}, a_2, a_3 is a simple triangle, and so (12) implies $\Delta(b_{12}, a_2, a_3) \geq \Delta(a, b, c)$. However, then

$$\ell(P) = \ell(P') + \Delta(b_{12}, a_2, a_3) \geq \rho(a, d) + \Delta(a, b, c),$$

contradicting (9). □

3. An Application in Block Designs

Recall that a (v, k, λ) design is a hypergraph on a set V of v points with the property that any pair of two points is contained in exactly λ edges with k points in each edge. We say a block design on V is *realizable as a metric space* if there is a metric space (V, ρ) such that for any three points $a, b, c \in V$ we have

$$\{a, b, c\} \in \mathcal{H}(\mathcal{B}(\rho)) \quad \text{if and only if } \{a, b, c\} \text{ is contained in some edge of the design.}$$

Recall that a finite projective plane of order n is an $(n^2 + n + 1, n + 1, 1)$ design, a finite affine plane of order n is an $(n^2, n, 1)$ design, and a Steiner triple system is a $(v, 3, 1)$ design. If a $(v, k, 1)$ design is realizable as a metric space, then every line in the metric space contains exactly k points. Therefore, immediately following Theorem 1 we have

Corollary 4. *No $(v, k, 1)$ design with $k \geq 3$ and $v > k$ is realizable as a metric space. In particular, no projective plane of order higher than 1, nor any affine plane of order higher than 2, nor any Steiner triple system with more than three points is realizable as a metric space.*

Mario Szegedy, in a discussion of this subject, first asked the question whether there is a short proof of the fact that no projective plane of order higher than 1 is realizable as a metric space. To the best of the author's knowledge, no simple proof is available; even the question is new. The special case of the Fano plane was solved by Chvátal in [3]; the proof was not simple.

Acknowledgments

I thank Vašek Chvátal for having introduced his nice conjecture to me. The way we present the proof of the main theorem here is also due to Vašek. In addition, I thank József Beck, Jeff Kahn, Bill Steiger, Mario Szegedy, and two anonymous referees for their helpful comments.

References

1. P. Borwein and W. O. J. Moser, A survey of Sylvester's problem and its generalizations, *Aequationes Mathematicae* **40** (1990), 111–135.
2. X. Chen, On a Conjecture of Chvátal, DIMACS Technical Report 2003-07, 2003.
3. V. Chvátal, Sylvester–Gallai theorem and metric betweenness, *Discrete & Computational Geometry* **31** (2004), 175–195.
4. H. S. M. Coxeter, A problem of collinear points, *American Mathematical Monthly* **55** (1948), 26–28.
5. H. S. M. Coxeter, *Introduction to Geometry*, Wiley, New York, 1961.
6. M. Deza and M. Laurent, *Geometry of Cuts and Metrics*, Springer-Verlag, Berlin, 1997.
7. P. Erdős and G. Purdy, Some extremal problems in combinatorial geometry, in *Handbook of Combinatorics* (R. L. Graham, M. Grötschel, and L. Lovász, eds.), pp. 809–874, North-Holland, Amsterdam, 1991.
8. D. Gale, *Tracking the Automatic Ant and Other Mathematical Explorations*, Springer-Verlag, New York, 1998.
9. T. Gallai, Solution to problem no. 4065, *American Mathematical Monthly* **51** (1944), 169–171.
10. K. Menger, Untersuchungen über allgemeine metrik, *Mathematische Annalen* **100** (1928), 75–163.
11. J. Pach and P. K. Agarwal, *Combinatorial Geometry*, Wiley-Interscience Series in Discrete Mathematics and Optimization, Wiley, New York, 1995.
12. J. J. Sylvester, Mathematical Question 11851, *Educational Times* **59** (1893), 98.

Received November 5, 2003, and in revised form March 13, 2004. Online publication December 30, 2005.