

Evaluating an optimal gastric closure method for transgastric surgery

M. Ryou · R. D. Pai · J. S. Sauer · D. W. Rattner ·
C. C. Thompson

Published online: 22 June 2007
© Springer Science+Business Media, LLC 2007

Erratum to: Surg Endosc
DOI 10.1007/s00464-006-9075-x

This article was published in Vol. 21, No. 4, pp. 677–680, and the following Key words were omitted: NOTES, Transluminal, Translumenal, Transcolonic, Transvaginal, Access, and Transvisceral.

Also omitted were the middle initials of the following authors: R. D. Pai, J. S. Sauer, D. W. Rattner, and C. C. Thompson.

Springer regrets the error.

The online version of the original article can be found under
doi:[10.1007/s00464-006-9075-x](https://doi.org/10.1007/s00464-006-9075-x)

M. Ryou · R. D. Pai · J. S. Sauer · C. C. Thompson (✉)
Department of Gastroenterology, Brigham & Women's Hospital,
75 Francis St., Boston, Massachusetts 02115, USA
e-mail: ccthompson@partners.org

J. S. Sauer
LSI Solutions, 7796 Victor-Mendon Rd., Victor,
New York 14564, USA

D. W. Rattner
Department of Surgery, Massachusetts General Hospital,
55 Fruit St., Boston, Massachusetts 02114, USA