Video-assisted breast surgery and sentinel lymph node biopsy guided by three-dimensional computed tomographic lymphography

K. Yamashita · K. Shimizu

Published online: 1 September 2007
© Springer Science+Business Media, LLC 2007

Erratum to: Surg Endosc
DOI 10.1007/s00464-007-9407-5

The correct version of Fig. 2(a) is hereby reproduced.

Fig. 2 Three-dimensional computed tomographic (3D-CT) lymphography. Iopamidol (2 ml) was injected intracutaneously into the periareolar skin and the skin above the tumor. Lymph ducts and the sentinel lymph node (SLN) are visualized with 3D-CT lymphography. The lymph ducts are draining lymph each from periareola and the tumor into a single SLN

The online version of the original article can be found under doi:
[10.1007/s00464-007-9407-5](https://doi.org/10.1007/s00464-007-9407-5)

K. Yamashita (✉)
Department of Surgery, Nippon Medical School Musashikosugi
Hospital, 1-396 Kosugi-cho, Nakahara-ku, Kawasaki,
Kanagawa 211-8533, Japan
e-mail: yamasita@nms.ac.jp

K. Shimizu
Division of Endocrine Surgery, Department of Surgery,
Nippon Medical School, 1-5, Sendagi-1, Bunkyo-ku,
Tokyo 113-8602, Japan