

Erratum to: Capnographic monitoring for carbon dioxide insufflation during endoscopic submucosal dissection: comparison of transcutaneous and end-tidal capnometers

Ryusaku Kusunoki · Yuji Amano · Takafumi Yuki · Akihiko Oka ·
Mayumi Okada · Yasumasa Tada · Goichi Uno · Ichiro Moriyama ·
Noriyoshi Ishimura · Shunji Ishihara · Yoshikazu Kinoshita

Published online: 13 December 2011
© Springer Science+Business Media, LLC 2011

Erratum to: Surg Endosc
DOI 10.1007/s00464-011-1908-6

In the original publication, “mucosal” should be “submucosal” in the title. This has been corrected in this erratum.

The online version of the original article can be found under
doi:[10.1007/s00464-011-1908-6](https://doi.org/10.1007/s00464-011-1908-6).

R. Kusunoki · A. Oka · M. Okada · Y. Tada · N. Ishimura ·
S. Ishihara · Y. Kinoshita
Second Department of Internal Medicine, Faculty of Medicine,
Shimane University, Izumo, Japan

Y. Amano (✉) · T. Yuki · G. Uno
Division of Endoscopy, Shimane University Hospital,
89-1, Enya-cho, Izumo-shi, Shimane 693-8501, Japan
e-mail: amano@med.shimane-u.ac.jp

I. Moriyama
Division of Cancer Center, Shimane University Hospital,
89-1, Enya-cho, Izumo-shi, Shimane 693-8501, Japan