

Erratum to: The use of self-gripping (ProgridTM) mesh during laparoscopic total extraperitoneal (TEP) inguinal hernia repair: a prospective feasibility and long-term outcomes study

Erin Bresnahan · Andrew Bates · Andrew Wu ·
Mark Reiner · Brian Jacob

Published online: 20 February 2015
© Springer Science+Business Media New York 2015

Erratum to: Surg Endosc
DOI 10.1007/s00464-014-3991-y

In Table 1, the Total # of patients should be 62. The seroma rate is mentioned as 13.2 % on pages 4 and 5, but should be 14.3 % as listed in Table 3. Springer regrets the error.

The online version of the original article can be found under doi:[10.1007/s00464-014-3991-y](https://doi.org/10.1007/s00464-014-3991-y).

E. Bresnahan · A. Wu · M. Reiner · B. Jacob
Department of Surgery, Mount Sinai Medical Center, New York,
NY, USA

A. Bates (✉) · B. Jacob
Department of Surgery, Mount Sinai Hospital, New York, NY,
USA
e-mail: andrewtbates@gmail.com

A. Wu
Metabolic, Endocrine and Minimally Invasive Surgery, Icahn
Mount Sinai School of Medicine, New York, NY, USA

B. Jacob
Department of Surgery, Laparoscopic Surgical Center of New
York, New York, NY, USA